

Electro-Magnetic Field Equation and Lorentz gauge in Rindler spacetime

Sangwha-Yi

Department of Math , Taejon University 300-716

ABSTRACT

In the general relativity theory, we find the electro-magnetic field transformation and the electro-magnetic field equation (Maxwell equation) in Rindler spacetime. We treat Lorentz gauge transformation in Rindler spacetime. We prove the electro-magnetic wave function cannot exist in Rindler spacetime. Specially, this article say the uniqueness of the accelerated frame because the accelerated frame can treat electro-magnetic field equation.

PACS Number:04,04.90.+e,03.30, 41.20

Key words:General relativity theory,

Rindler spacetime,

Electro-magnetic field transformation,

Electro-magnetic field equation

Lorentz gauge transformation

e-mail address:sangwha1@nate.com

Tel:051-624-3953

1. Introduction

In 2007 year, G.F.Torres del Castillo and C.I.Perez Sanchez already discovered Maxwell equations in uniformly accelerated frame in vacuum (see Ref [13]). In 2011 year, J.W.Maluf and F.F.Faria discovered electro-magnetic field transformation in Rindler space-time on ArXiv preprint(see also Ref[11]). But they did mistake they used Maxwell equations of gravity field. Maxwell equations of uniformly accelerated frame have to treat in flat Minkowski space-time not in gravity space-time.

Our theory's aim is that we find electro-magnetic field equation in Rindler space-time in vacuum also not in vacuum in the general relativity theory. In Section 2, we prepare for finding electro-magnetic field equation in Rindler space-time. In this section, we discover Lorentz gauge transformation and Lorentz fixing condition, transformation of the electro-magnetic 4-vector potential in Rindler space-time. In Section 3, we define the electro-magnetic field in Rindler space-time and we find the transformation of the electro-magnetic field. In Section 4, we obtain the electro-magnetic field equation in Rindler space-time and we apply the gauge theory to Maxwell equations (discovered by us) in Rindler space-time for viewing invariant about the gauge transformation.

We think seriously electro-magnetic wave function (radiation) in Rindler space-time but we know it doesn't satisfy electro-magnetic wave equation in mathematically. (See APPENDIX A)Hence, in 2007 year and in 2011 year, all researchers mistake calculation of electro-magnetic wave function.

We understand electro-magnetic wave function can exist in inertial frame by J.C.Maxwell or A. Einstein.

2. Transformation of the electro-magnetic 4-vector potential, Lorentz gauge transformation and Lorentz gauge fixing condition

The Rindler coordinate transformation is

$$ct = \left(\frac{c^2}{a_0} + \xi^1\right) \sinh\left(\frac{a_0 \xi^0}{c}\right)$$
$$x = \left(\frac{c^2}{a_0} + \xi^1\right) \cosh\left(\frac{a_0 \xi^0}{c}\right) - \frac{c^2}{a_0}, y = \xi^2, z = \xi^3 \quad (1)$$

In this time, the tetrad $e^a{}_\mu$ is (see Ref [12])

$$d\tau^2 = dt^2 - \frac{1}{c^2}[dx^2 + dy^2 + dz^2]$$
$$= -\frac{1}{c^2} \eta_{ab} \frac{\partial x^a}{\partial \xi^\mu} \frac{\partial x^b}{\partial \xi^\nu} d\xi^\mu d\xi^\nu$$
$$= -\frac{1}{c^2} \eta_{ab} e^a{}_\mu e^b{}_\nu d\xi^\mu d\xi^\nu = -\frac{1}{c^2} g_{\mu\nu} d\xi^\mu d\xi^\nu, \quad e^a{}_\mu = \frac{\partial x^a}{\partial \xi^\mu} \quad (2)$$

$$e^{\alpha}_0(\xi^0) = \frac{\partial X^{\alpha}}{\partial \xi^0} = \left(\left(1 + \frac{a_0}{c^2} \xi^1\right) \cosh\left(\frac{a_0 \xi^0}{c}\right), \left(1 + \frac{a_0}{c^2} \xi^1\right) \sinh\left(\frac{a_0 \xi^0}{c}\right), 0, 0 \right) \quad (3)$$

About y -axis's and z -axis's orientation

$$e^{\alpha}_2(\xi^0) = \frac{\partial X^{\alpha}}{\partial \xi^2} = (0, 0, 1, 0) \quad , \quad e^{\alpha}_3(\xi^0) = \frac{\partial X^{\alpha}}{\partial \xi^3} = (0, 0, 0, 1) \quad (4)$$

The other unit vector $e^{\alpha}_1(\xi^0)$ is

$$e^{\alpha}_1(\xi^0) = \frac{\partial X^{\alpha}}{\partial \xi^1} = \left(\sinh\left(\frac{a_0 \xi^0}{c}\right), \cosh\left(\frac{a_0 \xi^0}{c}\right), 0, 0 \right) \quad (5)$$

Therefore,

$$\begin{aligned} c dt &= c \cosh\left(\frac{a_0 \xi^0}{c}\right) d\xi^0 \left(1 + \frac{a_0}{c^2} \xi^1\right) + \sinh\left(\frac{a_0 \xi^0}{c}\right) d\xi^1 \\ dx &= c \sinh\left(\frac{a_0 \xi^0}{c}\right) d\xi^0 \left(1 + \frac{a_0}{c^2} \xi^1\right) + \cosh\left(\frac{a_0 \xi^0}{c}\right) d\xi^1, \quad dy = d\xi^2, \quad dz = d\xi^3 \end{aligned} \quad (6)$$

The vector transformation is

$$V'^{\mu} = \frac{\partial X'^{\mu}}{\partial X^{\alpha}} V^{\alpha}, \quad U'_{\mu} = \frac{\partial X^{\alpha}}{\partial X'^{\mu}} U_{\alpha} \quad (7)$$

Therefore, the transformation of the electro-magnetic 4-vector potential $(\phi, \vec{A}) = A^{\alpha}$ is

$$\begin{aligned} A^{\alpha} &= \frac{\partial X^{\alpha}}{\partial X'^{\mu}} A'^{\mu} = \frac{\partial X^{\alpha}}{\partial \xi^{\mu}} A_{\xi}^{\mu} = e^{\alpha}_{\mu} A_{\xi}^{\mu}, \quad e^{\alpha}_{\mu} = \frac{\partial X^{\alpha}}{\partial \xi^{\mu}} \\ dx^{\alpha} &= \frac{\partial X^{\alpha}}{\partial X'^{\mu}} dx'^{\mu} = \frac{\partial X^{\alpha}}{\partial \xi^{\mu}} d\xi^{\mu} = e^{\alpha}_{\mu} d\xi^{\mu}, \quad e^{\alpha}_{\mu} = \frac{\partial X^{\alpha}}{\partial \xi^{\mu}} \end{aligned} \quad (8)$$

$$\left(\frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \nabla^2 \right) \phi = 4\pi\rho$$

$$\left(\frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \nabla^2 \right) \vec{A} = \frac{4\pi}{c} \vec{j}$$

$$\text{4-vector } (c\rho, \vec{j}) = \rho_0 \frac{dx^{\alpha}}{d\tau} \quad (9)$$

Lorentz gauge transformation is in Rindler space-time,

$$A^{\mu} \rightarrow A^{\mu} + \partial^{\mu} \Lambda = A^{\mu} + g^{\mu\nu} \partial_{\nu} \Lambda \quad , \quad \Lambda \text{ is a scalar function.}$$

$$g^{00} = -\frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)^2}, g^{11} = g^{22} = g^{33} = 1$$

Hence,

$$\phi_\xi \rightarrow \phi_\xi - \frac{1}{c} \frac{\partial \Lambda}{\partial \xi^0} \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)^2}, \bar{A}_\xi \rightarrow \bar{A}_\xi + \bar{\nabla}_\xi \Lambda, \Lambda \text{ is a scalar function.} \quad (10)$$

Lorentz gauge fixing condition is in Rindler space-time,

$$A^\mu{}_{;\mu} = \frac{\partial A^\mu}{\partial \xi^\mu} + \Gamma^\mu{}_{\mu\rho} A^\rho,$$

$$\Gamma^\mu{}_{\mu\rho} = \Gamma^0{}_{01} = \frac{1}{2} g^{00} \left(\frac{\partial g_{00}}{\partial \xi^1} \right) = \frac{a_0}{c^2} \frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)}$$

$$g^{00} = -\frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)^2}, g^{11} = g^{22} = g^{33} = 1 \quad (11)$$

$$0 = \frac{\partial \phi}{\partial t} + \bar{\nabla} \cdot \bar{A} = \frac{\partial \phi_\xi}{\partial \xi^0} + \bar{\nabla}_\xi \cdot \bar{A}_\xi + \frac{A_{\xi^1} a_0}{c^2} \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)}$$

$$A^\mu \rightarrow A^\mu + \partial^\mu \Lambda = A^\mu + g^{\mu\nu} \partial_\nu \Lambda$$

$$\begin{aligned} A^\mu{}_{;\mu} &= \frac{\partial A^\mu}{\partial \xi^\mu} + \Gamma^\mu{}_{\mu\rho} A^\rho \rightarrow \partial_\mu (A^\mu + g^{\mu\nu} \partial_\nu \Lambda) + \Gamma^0{}_{01} \left(A^1 + \frac{\partial \Lambda}{\partial \xi^1} \right) \\ &= \partial_\mu A^\mu + (g^{\mu\nu} \partial_\mu \partial_\nu) \Lambda + \Gamma^0{}_{01} \left(A^1 + \frac{\partial \Lambda}{\partial \xi^1} \right) \end{aligned}$$

$$\begin{aligned} 0 &= \frac{\partial \phi_\xi}{\partial \xi^0} + \bar{\nabla}_\xi \cdot \bar{A}_\xi + \frac{A_{\xi^1} a_0}{c^2} \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \\ &\rightarrow \frac{1}{c} \frac{\partial \phi_\xi}{\partial \xi^0} + \bar{\nabla}_\xi \cdot \bar{A}_\xi - \left[\frac{1}{c^2} \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)^2} \left(\frac{\partial}{\partial \xi^0} \right)^2 - \nabla_\xi^2 \right] \Lambda \\ &+ \frac{A_{\xi^1} a_0}{c^2} \frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} + \frac{\partial \Lambda}{\partial \xi^1} \frac{a_0}{c^2} \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} = 0 \end{aligned}$$

$$\left[\frac{1}{c^2} \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)^2} \left(\frac{\partial}{\partial \xi^0}\right)^2 - \nabla_{\xi^2}^2 \right] \Lambda - \frac{\partial \Lambda}{\partial \xi^1} \frac{a_0}{c^2} \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} = 0$$

(12)

Hence, the transformation of the electro-magnetic 4-vector potential (ϕ, \vec{A}) in inertial frame and the

electro-magnetic 4-vector potential $(\phi_{\xi}, \vec{A}_{\xi})$ in uniformly accelerated frame is

$$\begin{aligned} \phi &= \cosh\left(\frac{a_0 \xi^0}{c}\right) \left(1 + \frac{a_0}{c^2} \xi^1\right) \phi_{\xi} + \sinh\left(\frac{a_0 \xi^0}{c}\right) A_{\xi^1} \\ A_x &= \sinh\left(\frac{a_0 \xi^0}{c}\right) \left(1 + \frac{a_0}{c^2} \xi^1\right) \phi_{\xi} + \cosh\left(\frac{a_0 \xi^0}{c}\right) A_{\xi^1} \\ A_y &= A_{\xi^2}, A_z = A_{\xi^3} \end{aligned}$$

(13)

$$g = \begin{pmatrix} -\left(1 + \frac{a_0 \xi^1}{c^2}\right)^2 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \eta = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

(See Ref [12])

$$e^a_{\mu} e_b^{\mu} = \delta^a_b, \quad e^a_{\mu} e_a^{\nu} = \delta_{\mu}^{\nu}$$

$$e^a_{\mu} e_b^{\nu} \eta_{ab} = g_{\mu\nu} \rightarrow A^T \eta A = g$$

$$e_a^{\mu} e_b^{\nu} g_{\mu\nu} = \eta_{ab} \rightarrow (A^T)^{-1} g A^{-1} = (A^T)^{-1} A^T \eta A A^{-1} = \eta$$

$$e^a_{\mu} = \eta^{ab} g_{\mu\nu} e_b^{\nu} \rightarrow \eta^{-1} (A^T)^{-1} A^T \eta A = A = \eta^{-1} (A^T)^{-1} g \quad (14)$$

$$\begin{pmatrix} cdt \\ dx \\ dy \\ dz \end{pmatrix} = \begin{pmatrix} \cosh\left(\frac{a_0 \xi^0}{c}\right) \left(1 + \frac{a_0 \xi^1}{c^2}\right) & \sinh\left(\frac{a_0 \xi^0}{c}\right) & 0 & 0 \\ \sinh\left(\frac{a_0 \xi^0}{c}\right) \left(1 + \frac{a_0 \xi^1}{c^2}\right) & \cosh\left(\frac{a_0 \xi^0}{c}\right) & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} cd\xi^0 \\ d\xi^1 \\ d\xi^2 \\ d\xi^3 \end{pmatrix}$$

$$\begin{aligned}
&= A \begin{pmatrix} cd\xi^0 \\ d\xi^1 \\ d\xi^2 \\ d\xi^3 \end{pmatrix} \\
&= \begin{pmatrix} \cosh(\frac{a_0\xi^0}{c}) & \sinh(\frac{a_0\xi^0}{c}) & 0 & 0 \\ \sinh(\frac{a_0\xi^0}{c}) & \cosh(\frac{a_0\xi^0}{c}) & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} cd\xi^0 \\ d\xi^1 \\ d\xi^2 \\ d\xi^3 \end{pmatrix}
\end{aligned}$$

(15)

$$e_{\mu}^{\alpha} = \frac{\partial \xi^{\alpha}}{\partial x^{\mu}} = A^{-1} = \begin{pmatrix} \frac{\partial \xi^0}{\partial t} & \frac{\partial \xi^0}{\partial x} & \frac{\partial \xi^0}{\partial y} & \frac{\partial \xi^0}{\partial z} \\ \frac{\partial \xi^1}{\partial t} & \frac{\partial \xi^1}{\partial x} & \frac{\partial \xi^1}{\partial y} & \frac{\partial \xi^1}{\partial z} \\ \frac{\partial \xi^2}{\partial t} & \frac{\partial \xi^2}{\partial x} & \frac{\partial \xi^2}{\partial y} & \frac{\partial \xi^2}{\partial z} \\ \frac{\partial \xi^3}{\partial t} & \frac{\partial \xi^3}{\partial x} & \frac{\partial \xi^3}{\partial y} & \frac{\partial \xi^3}{\partial z} \end{pmatrix}$$

$$= \begin{pmatrix} \frac{\cosh(\frac{a_0\xi^0}{c})}{(1 + \frac{a_0\xi^1}{c^2})} & -\frac{\sinh(\frac{a_0\xi^0}{c})}{(1 + \frac{a_0\xi^1}{c^2})} & 0 & 0 \\ -\sinh(\frac{a_0\xi^0}{c}) & \cosh(\frac{a_0\xi^0}{c}) & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

(16)

$$\begin{pmatrix} \frac{1}{c} \frac{\partial}{\partial t} \\ \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \\ \frac{\partial}{\partial z} \end{pmatrix} = (A^{-1})^T \begin{pmatrix} \frac{1}{c} \frac{\partial}{\partial \xi^0} \\ \frac{\partial}{\partial \xi^1} \\ \frac{\partial}{\partial \xi^2} \\ \frac{\partial}{\partial \xi^3} \end{pmatrix} = (A^T)^{-1} \begin{pmatrix} \frac{1}{c} \frac{\partial}{\partial \xi^0} \\ \frac{\partial}{\partial \xi^1} \\ \frac{\partial}{\partial \xi^2} \\ \frac{\partial}{\partial \xi^3} \end{pmatrix}$$

$$= \begin{pmatrix} \frac{\cosh(\frac{a_0 \xi^0}{c})}{(1 + \frac{a_0 \xi^1}{c^2})} & -\sinh(\frac{a_0 \xi^0}{c}) & 0 & 0 \\ -\frac{\sinh(\frac{a_0 \xi^0}{c})}{(1 + \frac{a_0 \xi^1}{c^2})} & \cosh(\frac{a_0 \xi^0}{c}) & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \frac{1}{c} \frac{\partial}{\partial \xi^0} \\ \frac{\partial}{\partial \xi^1} \\ \frac{\partial}{\partial \xi^2} \\ \frac{\partial}{\partial \xi^3} \end{pmatrix}$$

$$= \begin{pmatrix} \cosh(\frac{a_0 \xi^0}{c}) & -\sinh(\frac{a_0 \xi^0}{c}) & 0 & 0 \\ -\sinh(\frac{a_0 \xi^0}{c}) & \cosh(\frac{a_0 \xi^0}{c}) & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \frac{1}{c} \frac{\partial}{\partial \hat{\xi}^0} \\ \frac{\partial}{\partial \hat{\xi}^1} \\ \frac{\partial}{\partial \hat{\xi}^2} \\ \frac{\partial}{\partial \hat{\xi}^3} \end{pmatrix} \quad (17)$$

$$\begin{aligned} \frac{1}{c} \frac{\partial}{\partial t} &= \frac{\partial \xi^0}{\partial t} \frac{1}{c} \frac{\partial}{\partial \xi^0} + \frac{\partial \xi^1}{\partial t} \frac{\partial}{\partial \xi^1} \\ &= \frac{\cosh(\frac{a_0 \xi^0}{c})}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{\partial \xi^0} - \sinh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \xi^1} \\ \frac{\partial}{\partial x} &= \frac{\partial \xi^0}{\partial x} \frac{1}{c} \frac{\partial}{\partial \xi^0} + \frac{\partial \xi^1}{\partial x} \frac{\partial}{\partial \xi^1} \end{aligned}$$

$$\begin{aligned}
&= -\frac{\sinh(\frac{a_0 \xi^0}{c})}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{\partial \xi^0} + \cosh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \xi^1} \\
\frac{\partial}{\partial y} &= \frac{\partial}{\partial \xi^2}, \quad \frac{\partial}{\partial z} = \frac{\partial}{\partial \xi^3} \\
\frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \nabla^2 &= \frac{1}{c^2 (1 + \frac{a_0 \xi^1}{c^2})^2} (\frac{\partial}{\partial \xi^0})^2 - \nabla_{\xi}^2 \\
\vec{\nabla} &= (\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z}), \quad \vec{\nabla}_{\xi} = (\frac{\partial}{\partial \xi^1}, \frac{\partial}{\partial \xi^2}, \frac{\partial}{\partial \xi^3})
\end{aligned} \tag{18}$$

3. Electro-magnetic Field in the Rindler space-time

The electro-magnetic field (\vec{E}, \vec{B}) is in the inertial frame,

$$\vec{E} = -\vec{\nabla} \phi - \frac{\partial \vec{A}}{\partial t}, \quad \vec{B} = \vec{\nabla} \times \vec{A} \tag{19}$$

We have to calculate for define electro-magnetic field in Rindler space-time.

$$\begin{aligned}
E_x &= -\frac{\partial \phi}{\partial x} - \frac{\partial A_x}{\partial t} \\
&= -\left[-\frac{\sinh(\frac{a_0 \xi^0}{c})}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{\partial \xi^0} + \cosh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \xi^1} \right] \cdot \left[\cosh(\frac{a_0 \xi^0}{c}) (1 + \frac{a_0 \xi^1}{c^2}) \phi_{\xi} + \sinh(\frac{a_0 \xi^0}{c}) A_{\xi^1} \right] \\
&\quad - \left[\frac{\cosh(\frac{a_0 \xi^0}{c})}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{\partial \xi^0} - \sinh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \xi^1} \right] \cdot \left[\sinh(\frac{a_0 \xi^0}{c}) (1 + \frac{a_0 \xi^1}{c^2}) \phi_{\xi} + \cosh(\frac{a_0 \xi^0}{c}) A_{\xi^1} \right] \\
&= -\frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial A_{\xi^1}}{\partial \xi^0} - (1 + \frac{a_0 \xi^1}{c^2}) \frac{\partial \phi_{\xi}}{\partial \xi^1} - 2\phi_{\xi} \frac{a_0}{c^2} \\
&= -\frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{\partial \xi^1} \left[(1 + \frac{a_0 \xi^1}{c^2})^2 \phi_{\xi} \right] - \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial A_{\xi^1}}{\partial \xi^0} \tag{20}
\end{aligned}$$

$$E_y = -\frac{\partial \phi}{\partial y} - \frac{\partial A_y}{\partial t} = -\frac{\partial}{\partial \xi^2} \left[\cosh(\frac{a_0 \xi^0}{c}) (1 + \frac{a_0 \xi^1}{c^2}) \phi_{\xi} + \sinh(\frac{a_0 \xi^0}{c}) A_{\xi^1} \right]$$

$$\begin{aligned}
& - \left[\frac{\cosh\left(\frac{a_0 \xi^0}{c}\right)}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial}{\partial \xi^0} - \sinh\left(\frac{a_0 \xi^0}{c}\right) \frac{\partial}{\partial \xi^1} \right] A_{\xi^2} \\
= & - \left(1 + \frac{a_0 \xi^1}{c^2}\right) \cosh\left(\frac{a_0 \xi^0}{c}\right) \frac{\partial \phi_\xi}{\partial \xi^2} - \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \cosh\left(\frac{a_0 \xi^0}{c}\right) \frac{\partial A_{\xi^2}}{\partial \xi^0} \\
& + \sinh\left(\frac{a_0}{c} \xi^0\right) \left[\frac{\partial A_{\xi^2}}{\partial \xi^1} - \frac{\partial A_{\xi^1}}{\partial \xi^2} \right] \\
= & \cosh\left(\frac{a_0}{c} \xi^0\right) \left[- \frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} \frac{\partial}{\partial \xi^2} \left[\phi_\xi \left(1 + \frac{a_0 \xi^1}{c^2}\right)^2 \right] - \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial A_{\xi^2}}{\partial \xi^0} \right] \\
& + \sinh\left(\frac{a_0}{c} \xi^0\right) \left[\frac{\partial A_{\xi^2}}{\partial \xi^1} - \frac{\partial A_{\xi^1}}{\partial \xi^2} \right] \tag{21}
\end{aligned}$$

$$\begin{aligned}
E_z = -\frac{\partial \phi}{\partial z} - \frac{\partial A_z}{\partial t} = & - \frac{\partial}{\partial \xi^3} \left[\cosh\left(\frac{a_0 \xi^0}{c}\right) \left(1 + \frac{a_0}{c^2} \xi^1\right) \phi_\xi + \sinh\left(\frac{a_0 \xi^0}{c}\right) A_{\xi^1} \right] \\
& - \left[\frac{\cosh\left(\frac{a_0 \xi^0}{c}\right)}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial}{\partial \xi^0} - \sinh\left(\frac{a_0 \xi^0}{c}\right) \frac{\partial}{\partial \xi^1} \right] A_{\xi^3} \\
= & - \left(1 + \frac{a_0 \xi^1}{c^2}\right) \cosh\left(\frac{a_0 \xi^0}{c}\right) \frac{\partial \phi_\xi}{\partial \xi^3} - \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \cosh\left(\frac{a_0 \xi^0}{c}\right) \frac{\partial A_{\xi^3}}{\partial \xi^0} \\
& + \sinh\left(\frac{a_0}{c} \xi^0\right) \left[\frac{\partial A_{\xi^3}}{\partial \xi^1} - \frac{\partial A_{\xi^1}}{\partial \xi^3} \right] \\
= & \cosh\left(\frac{a_0}{c} \xi^0\right) \left[- \frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} \frac{\partial}{\partial \xi^3} \left[\phi_\xi \left(1 + \frac{a_0 \xi^1}{c^2}\right)^2 \right] - \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial A_{\xi^3}}{\partial \xi^0} \right] \\
& + \sinh\left(\frac{a_0}{c} \xi^0\right) \left[\frac{\partial A_{\xi^3}}{\partial \xi^1} - \frac{\partial A_{\xi^1}}{\partial \xi^3} \right]
\end{aligned} \tag{22}$$

$$B_x = \frac{\partial A_z}{\partial y} - \frac{\partial A_y}{\partial z} = \frac{\partial A_{\xi^3}}{\partial \xi^2} - \frac{\partial A_{\xi^2}}{\partial \xi^3} \quad (23)$$

$$\begin{aligned} B_y &= \frac{\partial A_x}{\partial z} - \frac{\partial A_z}{\partial x} = \frac{\partial A_x}{\partial \xi^3} - \frac{\partial A_{\xi^3}}{\partial x} \\ &= \frac{\partial}{\partial \xi^3} \left[\sinh\left(\frac{a_0 \xi^0}{c}\right) \left(1 + \frac{a_0}{c^2} \xi^1\right) \phi_\xi + \cosh\left(\frac{a_0 \xi^0}{c}\right) A_{\xi^1} \right] \\ &\quad - \left[-\frac{\sinh\left(\frac{a_0 \xi^0}{c}\right)}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial}{\partial \xi^0} + \cosh\left(\frac{a_0 \xi^0}{c}\right) \frac{\partial}{\partial \xi^1} \right] A_{\xi^3} \\ &= \cosh\left(\frac{a_0}{c} \xi^0\right) \left[\frac{\partial A_{\xi^1}}{\partial \xi^3} - \frac{\partial A_{\xi^3}}{\partial \xi^1} \right] \\ &\quad - \sinh\left(\frac{a_0}{c} \xi^0\right) \left[-\frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} \frac{\partial}{\partial \xi^3} \left[\phi_\xi \left(1 + \frac{a_0 \xi^1}{c^2}\right)^2 \right] - \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial A_{\xi^3}}{\partial \xi^0} \right] \end{aligned} \quad (24)$$

$$\begin{aligned} B_z &= \frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y} = \frac{\partial A_{\xi^2}}{\partial x} - \frac{\partial A_x}{\partial \xi^2} \\ &= \left[-\frac{\sinh\left(\frac{a_0 \xi^0}{c}\right)}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial}{\partial \xi^0} + \cosh\left(\frac{a_0 \xi^0}{c}\right) \frac{\partial}{\partial \xi^1} \right] A_{\xi^2} \\ &\quad - \frac{\partial}{\partial \xi^2} \left[\sinh\left(\frac{a_0 \xi^0}{c}\right) \left(1 + \frac{a_0}{c^2} \xi^1\right) \phi_\xi + \cosh\left(\frac{a_0 \xi^0}{c}\right) A_{\xi^1} \right] \\ &= \cosh\left(\frac{a_0}{c} \xi^0\right) \left[\frac{\partial A_{\xi^2}}{\partial \xi^1} - \frac{\partial A_{\xi^1}}{\partial \xi^2} \right] \\ &\quad + \sinh\left(\frac{a_0}{c} \xi^0\right) \left[-\frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} \frac{\partial}{\partial \xi^2} \left[\phi_\xi \left(1 + \frac{a_0 \xi^1}{c^2}\right)^2 \right] - \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial A_{\xi^2}}{\partial \xi^0} \right] \end{aligned} \quad (25)$$

Hence, we can define the electro-magnetic field $(\vec{E}_\xi, \vec{B}_\xi)$ in Rindler space-time.

$$\vec{E}_\xi = -\frac{1}{(1+\frac{a_0\xi^1}{c^2})} \vec{\nabla}_\xi \left\{ \phi_\xi \left(1 + \frac{a_0\xi^1}{c^2}\right)^2 \right\} - \frac{1}{(1+\frac{a_0\xi^1}{c^2})} \frac{\partial \vec{A}_\xi}{c\partial\xi^0}$$

$$\vec{B}_\xi = \vec{\nabla}_\xi \times \vec{A}_\xi$$

In this time, $\vec{\nabla}_\xi = \left(\frac{\partial}{\partial\xi^1}, \frac{\partial}{\partial\xi^2}, \frac{\partial}{\partial\xi^3} \right), \vec{A}_\xi = (A_{\xi^1}, A_{\xi^2}, A_{\xi^3})$ (26)

Lorentz gauge transformation is in Rindler space-time,

$$\phi_\xi \rightarrow \phi_\xi - \frac{1}{c} \frac{\partial\Lambda}{\partial\xi^0} \frac{1}{(1+\frac{a_0\xi^1}{c^2})}, \vec{A}_\xi \rightarrow \vec{A}_\xi + \vec{\nabla}_\xi \Lambda, \Lambda \text{ is a scalar function.} \quad (27)$$

$$\begin{aligned} \vec{E}_\xi &= -\frac{1}{(1+\frac{a_0\xi^1}{c^2})} \vec{\nabla}_\xi \left\{ \phi_\xi \left(1 + \frac{a_0\xi^1}{c^2}\right)^2 \right\} + \frac{1}{(1+\frac{a_0\xi^1}{c^2})} \vec{\nabla}_\xi \frac{\partial\Lambda}{c\partial\xi^0} \\ &\quad - \frac{1}{(1+\frac{a_0\xi^1}{c^2})} \frac{\partial \vec{A}_\xi}{c\partial\xi^0} - \frac{1}{(1+\frac{a_0\xi^1}{c^2})} \frac{\partial}{c\partial\xi^0} \vec{\nabla}_\xi \Lambda \\ &= -\frac{1}{(1+\frac{a_0\xi^1}{c^2})} \vec{\nabla}_\xi \left\{ \phi_\xi \left(1 + \frac{a_0\xi^1}{c^2}\right)^2 \right\} - \frac{1}{(1+\frac{a_0\xi^1}{c^2})} \frac{\partial \vec{A}_\xi}{c\partial\xi^0} \end{aligned}$$

$$\vec{B}_\xi = \vec{\nabla}_\xi \times \vec{A}_\xi + \vec{\nabla}_\xi \times \vec{\nabla}_\xi \Lambda = \vec{\nabla}_\xi \times \vec{A}_\xi \quad (28)$$

Lorentz gauge fixing condition is in Rindler space-time,

$$\begin{aligned} 0 &= \frac{\partial\phi_\xi}{c\partial\xi^0} + \vec{\nabla}_\xi \cdot \vec{A}_\xi + \frac{A_{\xi^1} a_0}{c^2} \frac{1}{(1+\frac{a_0\xi^1}{c^2})} \\ &\rightarrow \frac{1}{c} \frac{\partial\phi_\xi}{\partial\xi^0} + \vec{\nabla}_\xi \cdot \vec{A}_\xi - \left[\frac{1}{c^2} \frac{1}{(1+\frac{a_0\xi^1}{c^2})^2} \left(\frac{\partial}{\partial\xi^0} \right)^2 - \nabla_\xi^2 \right] \Lambda \end{aligned}$$

$$\begin{aligned}
& + \frac{A_{\xi^1} a_0}{c^2} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} + \frac{\partial \Lambda}{\partial \xi^1} \frac{a_0}{c^2} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} = 0 \\
& \left[\frac{1}{c^2} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} \left(\frac{\partial}{\partial \xi^0} \right)^2 - \nabla_{\xi}^2 \right] \Lambda - \frac{\partial \Lambda}{\partial \xi^1} \frac{a_0}{c^2} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} = 0 \\
& \phi - \frac{1}{c} \frac{\partial \Lambda}{\partial t} = \cosh\left(\frac{a_0 \xi^0}{c}\right) \left(1 + \frac{a_0}{c^2} \xi^1\right) \left\{ \phi_{\xi} - \frac{1}{c} \frac{\partial \Lambda}{\partial \xi^0} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} \right\} \\
& + \sinh\left(\frac{a_0 \xi^0}{c}\right) \left(A_{\xi^1} + \frac{\partial \Lambda}{\partial \xi^1} \right) \\
& A_x + \frac{\partial \Lambda}{\partial x} = \sinh\left(\frac{a_0 \xi^0}{c}\right) \left(1 + \frac{a_0}{c^2} \xi^1\right) \left\{ \phi_{\xi} - \frac{1}{c} \frac{\partial \Lambda}{\partial \xi^0} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} \right\} \\
& + \cosh\left(\frac{a_0 \xi^0}{c}\right) \left(A_{\xi^1} + \frac{\partial \Lambda}{\partial \xi^1} \right) \\
& A_y + \frac{\partial \Lambda}{\partial y} = A_{\xi^2} + \frac{\partial \Lambda}{\partial \xi^2}, \quad A_z + \frac{\partial \Lambda}{\partial z} = A_{\xi^3} + \frac{\partial \Lambda}{\partial \xi^3} \tag{29}
\end{aligned}$$

We obtain the transformation of the electro-magnetic field.

$$\begin{aligned}
E_x &= - \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial}{\partial \xi^1} \left\{ \phi_{\xi} \left(1 + \frac{a_0}{c^2} \xi^1\right)^2 \right\} - \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial A_{\xi^1}}{\partial \xi^0} = E_{\xi^1}, \\
E_y &= E_{\xi^2} \cosh\left(\frac{a_0 \xi^0}{c}\right) + B_{\xi^3} \sinh\left(\frac{a_0 \xi^0}{c}\right), \\
E_z &= E_{\xi^3} \cosh\left(\frac{a_0 \xi^0}{c}\right) - B_{\xi^2} \sinh\left(\frac{a_0 \xi^0}{c}\right) \\
B_x &= B_{\xi^1}, \\
B_y &= B_{\xi^2} \cosh\left(\frac{a_0 \xi^0}{c}\right) - E_{\xi^3} \sinh\left(\frac{a_0 \xi^0}{c}\right) \\
B_z &= B_{\xi^3} \cosh\left(\frac{a_0 \xi^0}{c}\right) + E_{\xi^2} \sinh\left(\frac{a_0 \xi^0}{c}\right) \tag{30}
\end{aligned}$$

Hence,

$$E_x = E_{\xi^1}, B_x = B_{\xi^1},$$

$$\begin{pmatrix} E_y \\ B_y \\ E_z \\ B_z \end{pmatrix} = H \begin{pmatrix} E_{\xi^2} \\ B_{\xi^2} \\ E_{\xi^3} \\ B_{\xi^3} \end{pmatrix}$$

$$H = \begin{pmatrix} \cosh\left(\frac{a_0 \xi^0}{c}\right) & 0 & 0 & \sinh\left(\frac{a_0 \xi^0}{c}\right) \\ 0 & \cosh\left(\frac{a_0 \xi^0}{c}\right) & -\sinh\left(\frac{a_0 \xi^0}{c}\right) & 0 \\ 0 & -\sinh\left(\frac{a_0 \xi^0}{c}\right) & \cosh\left(\frac{a_0 \xi^0}{c}\right) & 0 \\ \sinh\left(\frac{a_0 \xi^0}{c}\right) & 0 & 0 & \cosh\left(\frac{a_0 \xi^0}{c}\right) \end{pmatrix} \quad (31)$$

The inverse-transformation of the electro-magnetic field is

$$E_{\xi^1} = E_x, B_{\xi^1} = B_x$$

$$\begin{pmatrix} E_{\xi^2} \\ B_{\xi^2} \\ E_{\xi^3} \\ B_{\xi^3} \end{pmatrix} = H^{-1} \begin{pmatrix} E_y \\ B_y \\ E_z \\ B_z \end{pmatrix}$$

$$H^{-1} = \begin{pmatrix} \cosh\left(\frac{a_0 \xi^0}{c}\right) & 0 & 0 & -\sinh\left(\frac{a_0 \xi^0}{c}\right) \\ 0 & \cosh\left(\frac{a_0 \xi^0}{c}\right) & \sinh\left(\frac{a_0 \xi^0}{c}\right) & 0 \\ 0 & \sinh\left(\frac{a_0 \xi^0}{c}\right) & \cosh\left(\frac{a_0 \xi^0}{c}\right) & 0 \\ -\sinh\left(\frac{a_0 \xi^0}{c}\right) & 0 & 0 & \cosh\left(\frac{a_0 \xi^0}{c}\right) \end{pmatrix} \quad (32)$$

(See also Ref [11])

$$E_{\xi^1} = E_x, B_{\xi^1} = B_x$$

$$E_{\xi^2} = E_y \cosh\left(\frac{a_0 \xi^0}{c}\right) - B_z \sinh\left(\frac{a_0 \xi^0}{c}\right),$$

$$\begin{aligned}
B_{\xi^2} &= B_y \cosh\left(\frac{a_0 \xi^0}{c}\right) + E_z \sinh\left(\frac{a_0 \xi^0}{c}\right) \\
E_{\xi^3} &= E_z \cosh\left(\frac{a_0 \xi^0}{c}\right) + B_y \sinh\left(\frac{a_0 \xi^0}{c}\right) \\
B_{\xi^3} &= B_z \cosh\left(\frac{a_0 \xi^0}{c}\right) - E_y \sinh\left(\frac{a_0 \xi^0}{c}\right)
\end{aligned} \tag{33}$$

4. Electro-magnetic Field Equation (Maxwell Equation) in the Rindler space-time

Maxwell equation is

$$\vec{\nabla} \cdot \vec{E} = 4\pi\rho \tag{34-i}$$

$$\vec{\nabla} \times \vec{B} = \frac{\partial \vec{E}}{c \partial t} + 4\pi \vec{j} \tag{34-ii}$$

$$\vec{\nabla} \cdot \vec{B} = 0 \tag{34-iii}$$

$$\vec{\nabla} \times \vec{E} = -\frac{\partial \vec{B}}{c \partial t} \tag{34-iv}$$

We continue boring calculation for discovering Maxwell equations in Rindler space-time..

$$1. \vec{\nabla} \cdot \vec{E} = 4\pi\rho$$

$$E_x = E_{\xi^1},$$

$$E_y = E_{\xi^2} \cosh\left(\frac{a_0 \xi^0}{c}\right) + B_{\xi^3} \sinh\left(\frac{a_0 \xi^0}{c}\right),$$

$$E_z = E_{\xi^3} \cosh\left(\frac{a_0 \xi^0}{c}\right) - B_{\xi^2} \sinh\left(\frac{a_0 \xi^0}{c}\right)$$

$$\begin{aligned}
4\pi\rho &= \frac{\partial E_x}{\partial x} + \frac{\partial E_y}{\partial y} + \frac{\partial E_z}{\partial z} \\
&= \left[-\frac{\sinh\left(\frac{a_0 \xi^0}{c}\right)}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial}{c \partial \xi^0} + \cosh\left(\frac{a_0 \xi^0}{c}\right) \frac{\partial}{\partial \xi^1} \right] E_{\xi^1} \\
&\quad + \frac{\partial}{\partial \xi^2} \left[E_{\xi^2} \cosh\left(\frac{a_0 \xi^0}{c}\right) + B_{\xi^3} \sinh\left(\frac{a_0 \xi^0}{c}\right) \right] \\
&\quad + \frac{\partial}{\partial \xi^3} \left[E_{\xi^3} \cosh\left(\frac{a_0 \xi^0}{c}\right) - B_{\xi^2} \sinh\left(\frac{a_0 \xi^0}{c}\right) \right]
\end{aligned}$$

$$= \cosh\left(\frac{a_0}{c} \xi^0\right) (\vec{\nabla}_\xi \cdot \vec{E}_\xi) + \sinh\left(\frac{a_0}{c} \xi^0\right) \left[\frac{\partial B_{\xi^3}}{\partial \xi^2} - \frac{\partial B_{\xi^2}}{\partial \xi^3} - \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial E_{\xi^1}}{\partial \xi^0} \right] \quad (35)$$

$$2. \vec{\nabla} \times \vec{B} = \frac{\partial \vec{E}}{\partial t} + \frac{4\pi}{c} \vec{j}$$

$$B_x = B_{\xi^1}$$

$$B_y = B_{\xi^2} \cosh\left(\frac{a_0 \xi^0}{c}\right) - E_{\xi^3} \sinh\left(\frac{a_0 \xi^0}{c}\right)$$

$$B_z = B_{\xi^3} \cosh\left(\frac{a_0 \xi^0}{c}\right) + E_{\xi^2} \sinh\left(\frac{a_0 \xi^0}{c}\right)$$

$$\text{X-component) } \frac{\partial B_z}{\partial y} - \frac{\partial B_y}{\partial z}$$

$$= \frac{\partial}{\partial \xi^2} \left[B_{\xi^3} \cosh\left(\frac{a_0 \xi^0}{c}\right) + E_{\xi^2} \sinh\left(\frac{a_0 \xi^0}{c}\right) \right]$$

$$- \frac{\partial}{\partial \xi^3} \left[B_{\xi^2} \cosh\left(\frac{a_0 \xi^0}{c}\right) - E_{\xi^3} \sinh\left(\frac{a_0 \xi^0}{c}\right) \right]$$

$$= \cosh\left(\frac{a_0}{c} \xi^0\right) \left[\frac{\partial B_{\xi^3}}{\partial \xi^2} - \frac{\partial B_{\xi^2}}{\partial \xi^3} \right] + \sinh\left(\frac{a_0 \xi^0}{c}\right) \left[\frac{\partial E_{\xi^2}}{\partial \xi^2} + \frac{\partial E_{\xi^3}}{\partial \xi^3} \right]$$

$$= \frac{\partial E_x}{\partial t} + \frac{4\pi}{c} j_x$$

$$= \left[\frac{\cosh\left(\frac{a_0 \xi^0}{c}\right)}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial}{\partial \xi^0} - \sinh\left(\frac{a_0 \xi^0}{c}\right) \frac{\partial}{\partial \xi^1} \right] E_{\xi^1} + \frac{4\pi}{c} j_x$$

Hence,

$$\frac{4\pi}{c} j_x$$

$$= \sinh\left(\frac{a_0 \xi^0}{c}\right) (\vec{\nabla}_\xi \cdot \vec{E}_\xi) + \cosh\left(\frac{a_0 \xi^0}{c}\right) \left[\frac{\partial B_{\xi^3}}{\partial \xi^2} - \frac{\partial B_{\xi^2}}{\partial \xi^3} - \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial E_{\xi^1}}{\partial \xi^0} \right] \quad (36)$$

$$\begin{aligned}
& \text{Y-component) } \frac{\partial B_x}{\partial z} - \frac{\partial B_z}{\partial x} \\
&= \frac{\partial B_{\xi^1}}{\partial \xi^3} \\
&= \left[-\frac{\sinh(\frac{a_0 \xi^0}{c})}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{\partial \xi^0} + \cosh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \xi^1} \right] \cdot [B_{\xi^3} \cosh(\frac{a_0 \xi^0}{c}) + E_{\xi^2} \sinh(\frac{a_0 \xi^0}{c})] \\
&= \frac{\partial E_y}{\partial t} + \frac{4\pi}{c} j_y \\
&= \left[\frac{\cosh(\frac{a_0 \xi^0}{c})}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{\partial \xi^0} - \sinh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \xi^1} \right] \cdot [E_{\xi^2} \cosh(\frac{a_0 \xi^0}{c}) + B_{\xi^3} \sinh(\frac{a_0 \xi^0}{c})] \\
&\quad + \frac{4\pi}{c} j_y \\
\frac{4\pi}{c} j_y &= \frac{\partial B_{\xi^1}}{\partial \xi^3} - \frac{\partial B_{\xi^3}}{\partial \xi^1} - \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{a_0}{c^2} B_{\xi^3} - \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial E_{\xi^2}}{\partial \xi^0} \\
&= \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial}{\partial \xi^3} \{B_{\xi^1} (1 + \frac{a_0}{c^2} \xi^1)\} - \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial}{\partial \xi^1} \{B_{\xi^3} (1 + \frac{a_0}{c^2} \xi^1)\} - \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial E_{\xi^2}}{\partial \xi^0}
\end{aligned} \tag{37}$$

$$\begin{aligned}
& \text{Z-component) } \frac{\partial B_y}{\partial x} - \frac{\partial B_x}{\partial y} \\
&= \left[-\frac{\sinh(\frac{a_0 \xi^0}{c})}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{\partial \xi^0} + \cosh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \xi^1} \right] \cdot [B_{\xi^2} \cosh(\frac{a_0 \xi^0}{c}) - E_{\xi^3} \sinh(\frac{a_0 \xi^0}{c})] \\
&\quad - \frac{\partial B_{\xi^1}}{\partial \xi^2}
\end{aligned}$$

$$\begin{aligned}
&= \frac{\partial E_z}{\partial t} + \frac{4\pi}{c} j_z \\
&= \left[\frac{\cosh\left(\frac{a_0 \xi^0}{c}\right)}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial}{\partial \xi^0} - \sinh\left(\frac{a_0 \xi^0}{c}\right) \frac{\partial}{\partial \xi^1} \right] \cdot [E_{\xi^3} \cosh\left(\frac{a_0 \xi^0}{c}\right) - B_{\xi^2} \sinh\left(\frac{a_0 \xi^0}{c}\right)] \\
&\quad + \frac{4\pi}{c} j_z \\
\frac{4\pi}{c} j_z &= \frac{\partial B_{\xi^2}}{\partial \xi^1} - \frac{\partial B_{\xi^1}}{\partial \xi^2} + \frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} \frac{a_0}{c^2} B_{\xi^2} - \frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} \frac{\partial E_{\xi^3}}{\partial \xi^0} \\
&= \frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} \frac{\partial}{\partial \xi^1} \{B_{\xi^2} (1 + \frac{a_0}{c^2} \xi^1)\} - \frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} \frac{\partial}{\partial \xi^2} \{B_{\xi^1} (1 + \frac{a_0}{c^2} \xi^1)\} - \frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} \frac{\partial E_{\xi^3}}{\partial \xi^0}
\end{aligned}$$

(38)

$$3. \vec{\nabla} \cdot \vec{B} = 0$$

$$\begin{aligned}
\vec{\nabla} \cdot \vec{B} &= \frac{\partial B_x}{\partial x} + \frac{\partial B_y}{\partial y} + \frac{\partial B_z}{\partial z} \\
&= \left[-\frac{\sinh\left(\frac{a_0 \xi^0}{c}\right)}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial}{\partial \xi^0} + \cosh\left(\frac{a_0 \xi^0}{c}\right) \frac{\partial}{\partial \xi^1} \right] B_{\xi^1} \\
&\quad + \frac{\partial}{\partial \xi^2} [B_{\xi^2} \cosh\left(\frac{a_0 \xi^0}{c}\right) - E_{\xi^3} \sinh\left(\frac{a_0 \xi^0}{c}\right)] \\
&\quad + \frac{\partial}{\partial \xi^3} [B_{\xi^3} \cosh\left(\frac{a_0 \xi^0}{c}\right) + E_{\xi^2} \sinh\left(\frac{a_0 \xi^0}{c}\right)] \\
&= \cosh\left(\frac{a_0 \xi^0}{c}\right) (\vec{\nabla}_{\xi} \cdot \vec{B}_{\xi}) + \sinh\left(\frac{a_0 \xi^0}{c}\right) \left[-\left(\frac{\partial E_{\xi^2}}{\partial \xi^3} + \frac{\partial E_{\xi^3}}{\partial \xi^2} \right) - \frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} \frac{\partial B_{\xi^1}}{\partial \xi^0} \right] = 0
\end{aligned}$$

(39)

$$4. \vec{\nabla} \times \vec{E} = -\frac{\partial \vec{B}}{\partial t}$$

$$E_x = E_{\xi^1} ,$$

$$E_y = E_{\xi^2} \cosh\left(\frac{a_0 \xi^0}{c}\right) + B_{\xi^3} \sinh\left(\frac{a_0 \xi^0}{c}\right),$$

$$E_z = E_{\xi^3} \cosh\left(\frac{a_0 \xi^0}{c}\right) - B_{\xi^2} \sinh\left(\frac{a_0 \xi^0}{c}\right)$$

$$\text{X-component) } \frac{\partial E_z}{\partial y} - \frac{\partial E_y}{\partial z}$$

$$= \frac{\partial}{\partial \xi^2} [E_{\xi^3} \cosh\left(\frac{a_0 \xi^0}{c}\right) - B_{\xi^2} \sinh\left(\frac{a_0 \xi^0}{c}\right)]$$

$$- \frac{\partial}{\partial \xi^3} [E_{\xi^2} \cosh\left(\frac{a_0 \xi^0}{c}\right) + B_{\xi^3} \sinh\left(\frac{a_0 \xi^0}{c}\right)]$$

$$= \cosh\left(\frac{a_0}{c} \xi^0\right) \left[\frac{\partial E_{\xi^3}}{\partial \xi^2} - \frac{\partial E_{\xi^2}}{\partial \xi^3} \right] - \sinh\left(\frac{a_0 \xi^0}{c}\right) \left[\frac{\partial B_{\xi^2}}{\partial \xi^2} + \frac{\partial B_{\xi^3}}{\partial \xi^3} \right]$$

$$= - \frac{\partial B_x}{\partial t}$$

$$= - \left[\frac{\cosh\left(\frac{a_0 \xi^0}{c}\right)}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial}{\partial \xi^0} - \sinh\left(\frac{a_0 \xi^0}{c}\right) \frac{\partial}{\partial \xi^1} \right] B_{\xi^1}$$

Hence,

$$- \sinh\left(\frac{a_0 \xi^0}{c}\right) (\vec{\nabla}_\xi \cdot \vec{B}_\xi) + \cosh\left(\frac{a_0 \xi^0}{c}\right) \left[\left(\frac{\partial E_{\xi^3}}{\partial \xi^2} - \frac{\partial E_{\xi^2}}{\partial \xi^3} \right) + \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial B_{\xi^1}}{\partial \xi^0} \right] = 0 \quad (40)$$

$$\text{Y-component) } \frac{\partial E_x}{\partial z} - \frac{\partial E_z}{\partial x}$$

$$= \frac{\partial E_{\xi^1}}{\partial \xi^3}$$

$$- \left[- \frac{\sinh\left(\frac{a_0 \xi^0}{c}\right)}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial}{\partial \xi^0} + \cosh\left(\frac{a_0 \xi^0}{c}\right) \frac{\partial}{\partial \xi^1} \right] \cdot [E_{\xi^3} \cosh\left(\frac{a_0 \xi^0}{c}\right) - B_{\xi^2} \sinh\left(\frac{a_0 \xi^0}{c}\right)]$$

$$\begin{aligned}
&= -\frac{\partial B_y}{\partial t} \\
&= -\left[\frac{\cosh\left(\frac{a_0 \xi^0}{c}\right)}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial}{\partial \xi^0} - \sinh\left(\frac{a_0 \xi^0}{c}\right) \frac{\partial}{\partial \xi^1} \right] \cdot [B_{\xi^2} \cosh\left(\frac{a_0 \xi^0}{c}\right) - E_{\xi^3} \sinh\left(\frac{a_0 \xi^0}{c}\right)] \\
&\quad \frac{\partial E_{\xi^1}}{\partial \xi^3} - \frac{\partial E_{\xi^3}}{\partial \xi^1} - \frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} \frac{a_0}{c^2} E_{\xi^3} + \frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} \frac{\partial B_{\xi^2}}{\partial \xi^0} \\
&= \frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} \frac{\partial}{\partial \xi^3} \left\{ E_{\xi^1} \left(1 + \frac{a_0}{c^2} \xi^1\right) \right\} - \frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} \frac{\partial}{\partial \xi^1} \left\{ E_{\xi^3} \left(1 + \frac{a_0 \xi^1}{c^2}\right) \right\} + \frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} \frac{\partial B_{\xi^2}}{\partial \xi^0} \\
&= 0
\end{aligned} \tag{41}$$

$$\begin{aligned}
&\text{Z-component) } \frac{\partial E_y}{\partial x} - \frac{\partial E_x}{\partial y} \\
&= \left[-\frac{\sinh\left(\frac{a_0 \xi^0}{c}\right)}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial}{\partial \xi^0} + \cosh\left(\frac{a_0 \xi^0}{c}\right) \frac{\partial}{\partial \xi^1} \right] \cdot [E_{\xi^2} \cosh\left(\frac{a_0 \xi^0}{c}\right) + B_{\xi^3} \sinh\left(\frac{a_0 \xi^0}{c}\right)] \\
&\quad - \frac{\partial E_{\xi^1}}{\partial \xi^2} \\
&= -\frac{\partial B_z}{\partial t} \\
&= -\left[\frac{\cosh\left(\frac{a_0 \xi^0}{c}\right)}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial}{\partial \xi^0} - \sinh\left(\frac{a_0 \xi^0}{c}\right) \frac{\partial}{\partial \xi^1} \right] \cdot [B_{\xi^3} \cosh\left(\frac{a_0 \xi^0}{c}\right) + E_{\xi^2} \sinh\left(\frac{a_0 \xi^0}{c}\right)] \\
&\quad \frac{\partial E_{\xi^2}}{\partial \xi^1} - \frac{\partial E_{\xi^1}}{\partial \xi^2} + \frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} \frac{a_0}{c^2} E_{\xi^2} + \frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} \frac{\partial B_{\xi^3}}{\partial \xi^0} \\
&= \frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} \frac{\partial}{\partial \xi^1} \left\{ E_{\xi^2} \left(1 + \frac{a_0}{c^2} \xi^1\right) \right\} - \frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} \frac{\partial}{\partial \xi^2} \left\{ E_{\xi^1} \left(1 + \frac{a_0 \xi^1}{c^2}\right) \right\} + \frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} \frac{\partial B_{\xi^3}}{\partial \xi^0}
\end{aligned}$$

= 0

(42)

Therefore, we obtain the electro-magnetic field equation by Eq (35)-Eq(42) in Rindler space-time. (See also Ref [13])

$$\vec{\nabla}_\xi \cdot \vec{E}_\xi = 4\pi\rho_\xi \left(1 + \frac{a_0\xi^1}{c^2}\right) \quad (43-i)$$

$$\frac{1}{\left(1 + \frac{a_0\xi^1}{c^2}\right)} \vec{\nabla}_\xi \times \left\{ \vec{B}_\xi \left(1 + \frac{a_0\xi^1}{c^2}\right) \right\} = \frac{1}{\left(1 + \frac{a_0\xi^1}{c^2}\right)} \frac{\partial \vec{E}_\xi}{\partial \xi^0} + \frac{4\pi}{c} \vec{j}_\xi \quad (43-ii)$$

$$\vec{\nabla}_\xi \cdot \vec{B}_\xi = 0 \quad (43-iii)$$

$$\frac{1}{\left(1 + \frac{a_0\xi^1}{c^2}\right)} \vec{\nabla}_\xi \times \left\{ \vec{E}_\xi \left(1 + \frac{a_0\xi^1}{c^2}\right) \right\} = -\frac{1}{\left(1 + \frac{a_0\xi^1}{c^2}\right)} \frac{\partial \vec{B}_\xi}{\partial \xi^0} \quad (43-iv)$$

$$\vec{E}_\xi = (E_{\xi^1}, E_{\xi^2}, E_{\xi^3}), \vec{B}_\xi = (B_{\xi^1}, B_{\xi^2}, B_{\xi^3}),$$

$$\vec{\nabla}_\xi = \left(\frac{\partial}{\partial \xi^1}, \frac{\partial}{\partial \xi^2}, \frac{\partial}{\partial \xi^3} \right)$$

Hence, the transformation of 4-vector $(c\rho, \vec{j}) = \rho_0 \frac{dx^\alpha}{d\tau}$ is

$$\begin{aligned} \rho &= \rho_\xi \left(1 + \frac{a_0\xi^1}{c^2}\right) \cosh\left(\frac{a_0\xi^0}{c}\right) + \frac{j_{\xi^1}}{c} \sinh\left(\frac{a_0\xi^0}{c}\right) \\ j_x &= j_{\xi^1} \cosh\left(\frac{a_0\xi^0}{c}\right) + c\rho_\xi \left(1 + \frac{a_0}{c^2} \xi^1\right) \sinh\left(\frac{a_0\xi^0}{c}\right), \quad j_y = j_{\xi^2}, j_z = j_{\xi^3} \end{aligned} \quad (44-i)$$

In this time, 4-vector $(c\rho_\xi, \vec{j}_\xi) = \rho_0 \frac{d\xi^\alpha}{d\tau}$

For instant, the spherical charge density ρ_ξ of a stationary accelerated frame is in a charged huge sphere.

$$t = \xi^0 = 0, 0 \leq x = \xi^1, y = \xi^2, z = \xi^3 \leq R,$$

$$\vec{E} = \vec{E}_\xi = \frac{Q}{R^3} \vec{r}, \quad \rho = \rho_0 = \frac{Q}{V} = \frac{3Q}{4\pi R^3}, \quad \vec{j} = \vec{j}_\xi = \vec{0}$$

$$\frac{3Q}{4\pi R^3} \frac{1}{\left(1 + \frac{a_0 R}{c^2}\right)} \leq \rho_\xi = \frac{\rho}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} = \frac{3Q}{4\pi R^3} \frac{1}{\left(1 + \frac{a_0 X}{c^2}\right)} \leq \frac{3Q}{4\pi R^3}$$

(44-ii)

Generally, the continuity equation is in Rindler space-time,

$$0 = j^\mu{}_{;\mu} = \frac{\partial j^\mu}{\partial \xi^\mu} + \Gamma^\mu{}_{\mu\rho} j^\rho,$$

$$\Gamma^\mu{}_{\mu\rho} = \Gamma^0{}_{01} = \frac{1}{2} g^{00} \left(\frac{\partial g_{00}}{\partial \xi^1} \right) = \frac{a_0}{c^2} \frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)}$$

$$g^{00} = -\frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)^2}, g^{11} = g^{22} = g^{33} = 1$$

$$0 = \frac{\partial \rho}{\partial t} + \vec{\nabla} \cdot \vec{j} = \frac{\partial \rho_\xi}{\partial \xi^0} + \vec{\nabla}_\xi \cdot \vec{j}_\xi + \frac{j_{\xi^1} a_0}{c^2} \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \quad (45)$$

We treat Lorentz gauge transformation about the electro-magnetic field equation in Rindler space-time.

Eq(43-i) is

$$\begin{aligned} \vec{\nabla}_\xi \cdot \vec{E}_\xi &= \vec{\nabla}_\xi \cdot \left\{ -\frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \vec{\nabla}_\xi \left\{ \phi_\xi \left(1 + \frac{a_0 \xi^1}{c^2}\right)^2 \right\} - \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial \vec{A}_\xi}{\partial \xi^0} \right\} \\ &= -\vec{\nabla}_\xi \left\{ \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \right\} \cdot \left[\vec{\nabla}_\xi \left\{ \phi_\xi \left(1 + \frac{a_0 \xi^1}{c^2}\right)^2 \right\} + \frac{\partial \vec{A}_\xi}{\partial \xi^0} \right] \\ &\quad - \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \left[\nabla_\xi^2 \left\{ \phi_\xi \left(1 + \frac{a_0 \xi^1}{c^2}\right)^2 \right\} + \frac{\partial}{\partial \xi^0} (\vec{\nabla}_\xi \cdot \vec{A}_\xi) \right] \\ &= \frac{a_0}{c^2} \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)^2} \left[\frac{\partial}{\partial \xi^1} \left\{ \phi_\xi \left(1 + \frac{a_0 \xi^1}{c^2}\right)^2 \right\} + \frac{\partial A_{\xi^1}}{\partial \xi^0} \right] \end{aligned}$$

$$\begin{aligned}
& -\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)}\left[\nabla_\xi^2-\frac{1}{c^2}\frac{1}{\left(1+\frac{a_0}{c^2}\xi^1\right)^2}\left(\frac{\partial}{\partial\xi^0}\right)^2\right]\left\{\phi_\xi\left(1+\frac{a_0\xi^1}{c^2}\right)^2\right\} \\
& -\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)}\frac{\partial}{\partial\xi^0}\left[-\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)}\frac{A_{\xi^1}a_0}{c^2}\right] \\
& \frac{1}{c}\frac{\partial\phi_\xi}{\partial\xi^0}+\vec{\nabla}_\xi\cdot\vec{A}_\xi=-\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)}\frac{a_0}{c^2}A_{\xi^1} \\
= & -\frac{a_0}{c^2}\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)}E_{\xi^1}-\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)}\left[\nabla_\xi^2-\frac{1}{c^2}\frac{1}{\left(1+\frac{a_0}{c^2}\xi^1\right)^2}\left(\frac{\partial}{\partial\xi^0}\right)^2\right]\left\{\phi_\xi\left(1+\frac{a_0\xi^1}{c^2}\right)^2\right\} \\
& +\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)^2}\frac{\partial A_{\xi^1}}{\partial\xi^0}\frac{a_0}{c^2} \\
& =4\pi\rho_\xi\left(1+\frac{a_0\xi^1}{c^2}\right) \tag{46}
\end{aligned}$$

If we apply Lorentz gauge transformation to Eq (46),

$$\begin{aligned}
\phi_\xi & \rightarrow \phi_\xi - \frac{1}{c}\frac{\partial\Lambda}{\partial\xi^0}\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)}, \quad \vec{A}_\xi \rightarrow \vec{A}_\xi + \vec{\nabla}_\xi\Lambda, \quad \Lambda \text{ is a scalar function.} \\
= & -\frac{a_0}{c^2}\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)}E_{\xi^1}-\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)}\left[\nabla_\xi^2-\frac{1}{c^2}\frac{1}{\left(1+\frac{a_0}{c^2}\xi^1\right)^2}\left(\frac{\partial}{\partial\xi^0}\right)^2\right]\left\{\phi_\xi\left(1+\frac{a_0\xi^1}{c^2}\right)^2\right\} \\
& +\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)}\left[\nabla_\xi^2-\frac{1}{c^2}\frac{1}{\left(1+\frac{a_0}{c^2}\xi^1\right)^2}\left(\frac{\partial}{\partial\xi^0}\right)^2\right]\frac{1}{c}\frac{\partial\Lambda}{\partial\xi^0} \\
& +\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)^2}\frac{\partial A_{\xi^1}}{\partial\xi^0}\frac{a_0}{c^2}+\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)^2}\frac{a_0}{c^2}\frac{\partial}{\partial\xi^0}\frac{\partial\Lambda}{\partial\xi^1} \\
= & -\frac{a_0}{c^2}\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)}E_{\xi^1}-\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)}\left[\nabla_\xi^2-\frac{1}{c^2}\frac{1}{\left(1+\frac{a_0}{c^2}\xi^1\right)^2}\left(\frac{\partial}{\partial\xi^0}\right)^2\right]\left\{\phi_\xi\left(1+\frac{a_0\xi^1}{c^2}\right)^2\right\} \\
& +\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)}\frac{\partial}{\partial\xi^0}\left\{\left[\nabla_\xi^2-\frac{1}{c^2}\frac{1}{\left(1+\frac{a_0}{c^2}\xi^1\right)^2}\left(\frac{\partial}{\partial\xi^0}\right)^2\right]\Lambda+\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)}\frac{a_0}{c^2}\frac{\partial\Lambda}{\partial\xi^1}\right\}
\end{aligned}$$

$$+ \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)^2} \frac{\partial A_{\xi^1}}{c \partial \xi^0} \frac{a_0}{c^2} \quad (47)$$

In this time,

$$\left[\frac{1}{c^2} \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)^2} \left(\frac{\partial}{\partial \xi^0}\right)^2 - \nabla_{\xi}^2 \right] \Lambda - \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{a_0}{c^2} \frac{\partial \Lambda}{\partial \xi^1} = 0 \quad (48)$$

Hence, Eq(43-i) is

$$\begin{aligned} & \vec{\nabla}_{\xi} \cdot \vec{E}_{\xi} \\ &= -\frac{a_0}{c^2} \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} E_{\xi^1} - \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \left[\nabla_{\xi}^2 - \frac{1}{c^2} \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)^2} \left(\frac{\partial}{\partial \xi^0}\right)^2 \right] \left\{ \phi_{\xi} \left(1 + \frac{a_0 \xi^1}{c^2}\right)^2 \right\} \\ &+ \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)^2} \frac{\partial A_{\xi^1}}{c \partial \xi^0} \frac{a_0}{c^2} \\ &= 4\pi\rho_{\xi} \left(1 + \frac{a_0 \xi^1}{c^2}\right) \end{aligned} \quad (49)$$

Eq(43-i) is invariant about Lorentz gauge transformation in Rindler space-time.

Eq (43-ii) is

$$\begin{aligned} & \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \vec{\nabla}_{\xi} \times \left\{ \vec{B}_{\xi} \left(1 + \frac{a_0 \xi^1}{c^2}\right) \right\} \\ &= \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \vec{\nabla}_{\xi} \times \left\{ \vec{\nabla}_{\xi} \times \vec{A}_{\xi} \left(1 + \frac{a_0 \xi^1}{c^2}\right) \right\} \\ &= \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \vec{\nabla}_{\xi} \left(1 + \frac{a_0 \xi^1}{c^2}\right) \times \left\{ \vec{\nabla}_{\xi} \times \vec{A}_{\xi} \right\} + \vec{\nabla}_{\xi} \times \vec{\nabla}_{\xi} \times \vec{A}_{\xi} \\ &= \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{a_0}{c^2} (1,0,0) \times \vec{B}_{\xi} + \left\{ -\nabla_{\xi}^2 \vec{A}_{\xi} + \vec{\nabla}_{\xi} (\vec{\nabla}_{\xi} \cdot \vec{A}_{\xi}) \right\} \\ &= \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{a_0}{c^2} (0, -B_{\xi^3}, B_{\xi^2}) + \left\{ -\nabla_{\xi}^2 \vec{A}_{\xi} + \vec{\nabla}_{\xi} (\vec{\nabla}_{\xi} \cdot \vec{A}_{\xi}) \right\} \end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)}\frac{\partial\vec{E}_\xi}{\partial\xi^0}+\frac{4\pi}{c}\vec{j}_\xi \\
&= -\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)^2}\frac{\partial}{\partial\xi^0}[\vec{\nabla}_\xi\{\phi_\xi(1+\frac{a_0\xi^1}{c^2})^2\}]-\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)^2}\frac{1}{c^2}\left(\frac{\partial}{\partial\xi^0}\right)^2\vec{A}_\xi+\frac{4\pi\vec{j}_\xi}{c} \\
&= -\frac{\partial}{\partial\xi^0}\vec{\nabla}_\xi\phi_\xi-\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)}\frac{2a_0}{c^2}\frac{\partial\phi_\xi}{\partial\xi^0}(1,0,0)-\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)^2}\frac{1}{c^2}\left(\frac{\partial}{\partial\xi^0}\right)^2\vec{A}_\xi+\frac{4\pi\vec{j}_\xi}{c}
\end{aligned} \tag{50}$$

Therefore,

$$\begin{aligned}
&\frac{4\pi}{c}\vec{j}_\xi \\
&= \frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)}\frac{a_0}{c^2}(0,-B_{\xi^3},B_{\xi^2})+\{-\nabla_\xi^2\vec{A}_\xi+\vec{\nabla}_\xi(\vec{\nabla}_\xi\cdot\vec{A}_\xi)\} \\
&+ \frac{\partial}{\partial\xi^0}\vec{\nabla}_\xi\phi_\xi+\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)}\frac{2a_0}{c^2}\frac{\partial\phi_\xi}{\partial\xi^0}(1,0,0)+\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)^2}\frac{1}{c^2}\left(\frac{\partial}{\partial\xi^0}\right)^2\vec{A}_\xi \\
&= \frac{a_0}{c^2}\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)}(0,-B_{\xi^3},B_{\xi^2})+\frac{1}{\left(1+\frac{a_0}{c^2}\xi^1\right)}\frac{2a_0}{c^2}\frac{\partial\phi_\xi}{\partial\xi^0}(1,0,0) \\
&+ [-\nabla_\xi^2+\frac{1}{c^2}\frac{1}{\left(1+\frac{a_0}{c^2}\xi^1\right)^2}\left(\frac{\partial}{\partial\xi^0}\right)^2]\vec{A}_\xi+\vec{\nabla}_\xi[-\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)}\frac{a_0}{c^2}A_{\xi^1}] \\
&\quad \frac{1}{c}\frac{\partial\phi_\xi}{\partial\xi^0}+\vec{\nabla}_\xi\cdot\vec{A}_\xi=-\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)}\frac{a_0}{c^2}A_{\xi^1}
\end{aligned} \tag{51}$$

$$\begin{aligned}
&\frac{4\pi}{c}\vec{j}_\xi \\
&= \frac{a_0}{c^2}\frac{1}{\left(1+\frac{a_0\xi^1}{c^2}\right)}(0,-B_{\xi^3},B_{\xi^2})+\frac{1}{\left(1+\frac{a_0}{c^2}\xi^1\right)}\frac{2a_0}{c^2}\frac{\partial\phi_\xi}{\partial\xi^0}(1,0,0)
\end{aligned}$$

$$\begin{aligned}
& + [-\nabla_{\xi}^2 + \frac{1}{c^2} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} (\frac{\partial}{\partial \xi^0})^2] \bar{A}_{\xi} \\
& + \bar{\nabla}_{\xi} [-\frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} \frac{a_0}{c^2} A_{\xi^1}] \tag{52}
\end{aligned}$$

If we apply Lorentz gauge transformation to Eq (52),

$$\phi_{\xi} \rightarrow \phi_{\xi} - \frac{1}{c} \frac{\partial \Lambda}{\partial \xi^0} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2}, \quad \bar{A}_{\xi} \rightarrow \bar{A}_{\xi} + \bar{\nabla}_{\xi} \Lambda, \quad \Lambda \text{ is a scalar function.}$$

$$\begin{aligned}
& \frac{4\pi}{c} \vec{j}_{\xi} \\
& = \frac{a_0}{c^2} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} (0, -B_{\xi^3}, B_{\xi^2}) + \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{2a_0}{c^2} \frac{\partial \phi_{\xi}}{c \partial \xi^0} (1, 0, 0) \\
& \quad - \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^3} \frac{2a_0}{c^2} \frac{1}{c^2} (\frac{\partial}{\partial \xi^0})^2 \Lambda (1, 0, 0) \\
& + [-\nabla_{\xi}^2 + \frac{1}{c^2} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} (\frac{\partial}{\partial \xi^0})^2] \bar{A}_{\xi} + [-\nabla_{\xi}^2 + \frac{1}{c^2} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} (\frac{\partial}{\partial \xi^0})^2] \bar{\nabla}_{\xi} \Lambda \\
& \quad + \bar{\nabla}_{\xi} [-\frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} \frac{a_0}{c^2} A_{\xi^1}] + \bar{\nabla}_{\xi} [-\frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} \frac{a_0}{c^2} \frac{\partial \Lambda}{\partial \xi^1}] \\
& = \frac{a_0}{c^2} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} (0, -B_{\xi^3}, B_{\xi^2}) + \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{2a_0}{c^2} \frac{\partial \phi_{\xi}}{c \partial \xi^0} (1, 0, 0) \\
& \quad - \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^3} \frac{2a_0}{c^2} \frac{1}{c^2} (\frac{\partial}{\partial \xi^0})^2 \Lambda (1, 0, 0) \\
& + [-\nabla_{\xi}^2 + \frac{1}{c^2} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} (\frac{\partial}{\partial \xi^0})^2] \bar{A}_{\xi} + [-\nabla_{\xi}^2 + \frac{1}{c^2} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} (\frac{\partial}{\partial \xi^0})^2] \bar{\nabla}_{\xi} \Lambda \\
& \quad + \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} \frac{a_0^2}{c^4} A_{\xi^1} (1, 0, 0) - \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} \frac{a_0}{c^2} \bar{\nabla}_{\xi} A_{\xi^1}
\end{aligned}$$

$$+ \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})^2} \frac{a_0^2}{c^4} \frac{\partial \Lambda}{\partial \xi^1} (1,0,0) - \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{a_0}{c^2} \frac{\partial}{\partial \xi^1} \vec{\nabla}_\xi \Lambda$$

(53)

In this time,

$$\begin{aligned} & \left[\frac{1}{c^2} \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})^2} \left(\frac{\partial}{\partial \xi^0} \right)^2 - \nabla_\xi^2 \right] \Lambda - \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{a_0}{c^2} \frac{\partial \Lambda}{\partial \xi^1} = 0 \\ 0 &= \vec{\nabla}_\xi \left[\left\{ -\nabla_\xi^2 + \frac{1}{c^2} \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})^2} \left(\frac{\partial}{\partial \xi^0} \right)^2 - \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{a_0}{c^2} \frac{\partial}{\partial \xi^1} \right\} \Lambda \right] \\ &= \vec{\nabla}_\xi \left\{ \frac{1}{c^2} \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})^2} \right\} \left(\frac{\partial}{\partial \xi^0} \right)^2 \Lambda + \left[-\nabla_\xi^2 + \frac{1}{c^2} \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})^2} \left(\frac{\partial}{\partial \xi^0} \right)^2 \right] \vec{\nabla}_\xi \Lambda \\ &\quad - \vec{\nabla}_\xi \left\{ \frac{a_0}{c^2} \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \right\} \frac{\partial \Lambda}{\partial \xi^1} - \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{a_0}{c^2} \frac{\partial}{\partial \xi^1} \vec{\nabla}_\xi \Lambda \\ &= -\frac{2}{c^2} \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})^3} \frac{a_0}{c^2} \left(\frac{\partial}{\partial \xi^0} \right)^2 \Lambda (1,0,0) + \left[-\nabla_\xi^2 + \frac{1}{c^2} \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})^2} \left(\frac{\partial}{\partial \xi^0} \right)^2 \right] \vec{\nabla}_\xi \Lambda \\ &\quad + \frac{a_0^2}{c^4} \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})^2} \frac{\partial \Lambda}{\partial \xi^1} (1,0,0) - \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{a_0}{c^2} \frac{\partial}{\partial \xi^1} \vec{\nabla}_\xi \Lambda \end{aligned} \tag{54}$$

Therefore,

$$\begin{aligned} & \frac{4\pi}{c} \vec{j}_\xi \\ &= \frac{a_0}{c^2} \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} (0, -B_{\xi^3}, B_{\xi^2}) + \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{2a_0}{c^2} \frac{\partial \phi_\xi}{\partial \xi^0} (1,0,0) \\ &\quad - \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})^3} \frac{2a_0}{c^2} \frac{1}{c^2} \left(\frac{\partial}{\partial \xi^0} \right)^2 \Lambda (1,0,0) \\ &\quad + \left[-\nabla_\xi^2 + \frac{1}{c^2} \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})^2} \left(\frac{\partial}{\partial \xi^0} \right)^2 \right] \vec{A}_\xi + \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})^2} \frac{a_0^2}{c^4} A_{\xi^1} (1,0,0) \end{aligned}$$

$$\begin{aligned}
& - \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{a_0}{c^2} \vec{\nabla}_\xi A_{\xi^1} + \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})^2} \frac{a_0^2}{c^4} \frac{\partial \Lambda}{\partial \xi^1} (1,0,0) - \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{a_0}{c^2} \frac{\partial}{\partial \xi^1} \vec{\nabla}_\xi \Lambda \\
& + \vec{\nabla}_\xi \left[\left\{ -\nabla_\xi^2 + \frac{1}{c^2} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} \left(\frac{\partial}{\partial \xi^0} \right)^2 - \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{a_0}{c^2} \frac{\partial}{\partial \xi^1} \right\} \Lambda \right] \\
& + \frac{2}{c^2} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^3} \frac{a_0}{c^2} \left(\frac{\partial}{\partial \xi^0} \right)^2 \Lambda (1,0,0) - \frac{a_0^2}{c^4} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} \frac{\partial \Lambda}{\partial \xi^1} (1,0,0) \\
& + \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{a_0}{c^2} \frac{\partial}{\partial \xi^1} \vec{\nabla}_\xi \Lambda \\
& = \frac{a_0}{c^2} \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} (0, -B_{\xi^3}, B_{\xi^2}) + \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{2a_0}{c^2} \frac{\partial \phi_\xi}{\partial \xi^0} (1,0,0) \\
& \quad + \left[-\nabla_\xi^2 + \frac{1}{c^2} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} \left(\frac{\partial}{\partial \xi^0} \right)^2 \right] \vec{A}_\xi \\
& + \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})^2} \frac{a_0^2}{c^4} A_{\xi^1} (1,0,0) - \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{a_0}{c^2} \vec{\nabla}_\xi A_{\xi^1} \tag{55}
\end{aligned}$$

Hence, Eq(43-ii) is invariant about Lorentz gauge transformation in Rindler space-time.

Eq (43-iii) is

$$\vec{\nabla}_\xi \cdot \vec{B}_\xi = \vec{\nabla}_\xi \cdot (\vec{\nabla}_\xi \times \vec{A}_\xi + \vec{\nabla}_\xi \times \vec{\nabla}_\xi \Lambda) = \vec{\nabla}_\xi \times \vec{\nabla}_\xi \cdot \vec{A}_\xi = 0 \tag{56}$$

Eq (43-iv) is

$$\begin{aligned}
& \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \vec{\nabla}_\xi \times \left\{ \vec{E}_\xi \left(1 + \frac{a_0 \xi^1}{c^2} \right) \right\} \\
& = - \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \vec{\nabla}_\xi \times \left[\vec{\nabla}_\xi \left\{ \phi_\xi \left(1 + \frac{a_0 \xi^1}{c^2} \right)^2 \right\} - \vec{\nabla}_\xi \left(\frac{\partial \Lambda}{\partial \xi^0} \right) + \frac{\partial \vec{A}_\xi}{\partial \xi^0} + \frac{\partial}{\partial \xi^0} (\vec{\nabla}_\xi \Lambda) \right]
\end{aligned}$$

$$= -\frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \vec{\nabla}_\xi \times \frac{\partial \vec{A}_\xi}{\partial \xi^0} = -\frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial (\vec{\nabla}_\xi \times \vec{A}_\xi)}{\partial \xi^0} = -\frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial \vec{B}_\xi}{\partial \xi^0} \quad (57)$$

Hence, Eq (43-iii), Eq (43-iv) are invariant about Lorentz gauge transformation in Rindler space-time.

Hence, the electro-magnetic field equations (Maxwell Equations) in Rindler space-time are invariant about Lorentz gauge transformation.

5. Conclusion

We find the electro-magnetic field transformation and the electro-magnetic equation in uniformly accelerated frame in one theory.

Generally, the coordinate transformation of accelerated frame is (see Ref [9])

$$(I) \quad ct = \left(\frac{c^2}{a_0} + \xi^1\right) \sinh\left(\frac{a_0 \xi^0}{c}\right)$$

$$x = \left(\frac{c^2}{a_0} + \xi^1\right) \cosh\left(\frac{a_0 \xi^0}{c}\right) - \frac{c^2}{a_0}, y = \xi^2, z = \xi^3 \quad (58)$$

$$(II) \quad ct = \frac{c^2}{a_0} \exp\left(\frac{a_0}{c^2} \xi^1\right) \sinh\left(\frac{a_0 \xi^0}{c}\right)$$

$$x = \frac{c^2}{a_0} \exp\left(\frac{a_0}{c^2} \xi^1\right) \cosh\left(\frac{a_0 \xi^0}{c}\right) - \frac{c^2}{a_0}, y = \xi^2, z = \xi^3 \quad (59)$$

If you try to use Eq(59) for make Maxwell equation in Rindler space-time, You have to fail it.

In A.Einstein's article (see Ref [10]), Einstein obtain Lorenz transformation by Maxwell equation in inertial frame, Einstein give up Galilei transformation in inertial frame. In accelerated frame, we think our article's choice is Rindler coordinate (I) can treat electro-magnetic field equation likely Einstein's election.

APPENDIX A

In 2-Dimension Rindler space-time, if we mistake calculation, we think the electro-magnetic wave function.

$$E_{\xi^1} = E_x = E_{x_0} \sin \omega \left(t - \frac{x}{c}\right)$$

$$= E_{x_0} \sin \omega \left[-\left\{\frac{c}{a_0} + \frac{\xi^1}{c}\right\} \exp\left(-\frac{a_0}{c} \xi^0\right) + \frac{c}{a_0}\right] = E_{x_0} \sin \Phi \quad (A-1)$$

$$ct = \left(\frac{c^2}{a_0} + \xi^1\right) \sinh\left(\frac{a_0 \xi^0}{c}\right), \quad x = \left(\frac{c^2}{a_0} + \xi^1\right) \cosh\left(\frac{a_0 \xi^0}{c}\right) - \frac{c^2}{a_0}$$

In this time,

$$-\exp\left(-\frac{a_0}{c} \xi^0\right) = \sinh\left(\frac{a_0 \xi^0}{c}\right) - \cosh\left(\frac{a_0 \xi^0}{c}\right) \quad (\text{A-2})$$

(A-1) have to satisfy the following equation.

$$\frac{1}{\left(1 + \frac{a_0 \xi^1}{c}\right)^2} \frac{1}{c^2} \left(\frac{\partial}{\partial \xi^0}\right)^2 \sin \Phi = \left(\frac{\partial}{\partial \xi^1}\right)^2 \sin \Phi \quad (\text{A-3})$$

$$\begin{aligned} \left(\frac{\partial}{\partial \xi^1}\right)^2 \sin \Phi &= \frac{\partial}{\partial \xi^1} \left\{ \frac{\partial}{\partial \xi^1} \sin \omega \left[-\left\{ \frac{c}{a_0} + \frac{\xi^1}{c} \right\} \exp\left(-\frac{a_0}{c} \xi^0\right) + \frac{c}{a_0} \right] \right\} \\ &= \frac{\partial}{\partial \xi^1} \left\{ (\cos \Phi) \cdot -\omega \frac{1}{c} \exp\left(-\frac{a_0}{c} \xi^0\right) \right\} = (-\sin \Phi) \cdot \frac{\omega^2}{c^2} \exp\left(-2 \frac{a_0}{c} \xi^0\right) \end{aligned} \quad (\text{A-4})$$

But this calculation situation is different.

$$\begin{aligned} &\frac{1}{\left(1 + \frac{a_0 \xi^1}{c}\right)^2} \frac{1}{c^2} \left(\frac{\partial}{\partial \xi^0}\right)^2 \sin \Phi \\ &= \frac{1}{\left(1 + \frac{a_0 \xi^1}{c}\right)^2} \frac{1}{c^2} \frac{\partial}{\partial \xi^0} \left\{ \frac{\partial}{\partial \xi^0} \sin \omega \left[-\left\{ \frac{c}{a_0} + \frac{\xi^1}{c} \right\} \exp\left(-\frac{a_0}{c} \xi^0\right) + \frac{c}{a_0} \right] \right\} \\ &= \frac{1}{\left(1 + \frac{a_0 \xi^1}{c}\right)^2} \frac{1}{c^2} \frac{\partial}{\partial \xi^0} \left\{ (\cos \Phi) \cdot \omega \left\{ \frac{c}{a_0} + \frac{\xi^1}{c} \right\} \frac{a_0}{c} \exp\left(-\frac{a_0}{c} \xi^0\right) \right\} \end{aligned}$$

In this time, $D(\alpha\beta) = \beta D\alpha + \alpha D\beta$

$$\begin{aligned} &= \frac{1}{\left(1 + \frac{a_0 \xi^1}{c}\right)^2} \frac{1}{c^2} \left\{ \frac{\partial}{\partial \xi^0} (\cos \Phi) \cdot \omega \left\{ \frac{c}{a_0} + \frac{\xi^1}{c} \right\} \frac{a_0}{c} \exp\left(-\frac{a_0}{c} \xi^0\right) \right. \\ &\quad \left. + (\cos \Phi) \cdot \omega \left\{ \frac{c}{a_0} + \frac{\xi^1}{c} \right\} \frac{a_0}{c} \frac{\partial}{\partial \xi^0} \left(\exp\left(-\frac{a_0}{c} \xi^0\right) \right) \right\} \\ &= \frac{1}{\left(1 + \frac{a_0 \xi^1}{c}\right)^2} \frac{1}{c^2} \left\{ -\sin \Phi \cdot \omega^2 \left\{ \frac{c}{a_0} + \frac{\xi^1}{c} \right\}^2 \frac{a_0^2}{c^2} \exp\left(-2 \frac{a_0}{c} \xi^0\right) \right. \end{aligned}$$

$$-(\cos \Phi) \cdot \omega \left\{ \frac{c}{a_0} + \frac{\xi^1}{c} \right\} \frac{a_0^2}{c^2} \exp\left(-\frac{a_0}{c} \xi^0\right) \quad (\text{A-5})$$

Hence, if we compare Eq(A-4) and Eq(A-5),

$$\begin{aligned} \left(\frac{\partial}{\partial \xi^1}\right)^2 \sin \Phi &= (-\sin \Phi) \cdot \frac{\omega^2}{c^2} \exp\left(-2 \frac{a_0}{c} \xi^0\right) \\ &\neq -\sin \Phi \cdot \omega^2 \frac{1}{c^2} \exp\left(-2 \frac{a_0}{c} \xi^0\right) \end{aligned}$$

$$-\frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} (\cos \Phi) \cdot \omega \frac{a_0}{c} \frac{1}{c^2} \exp\left(-\frac{a_0}{c} \xi^0\right) = \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)^2} \frac{1}{c^2} \left(\frac{\partial}{\partial \xi^0}\right)^2 \sin \Phi \quad (\text{A-6})$$

Therefore, we think it cannot exist electro-magnetic wave function in Rindler space-time.

Reference

- [1]S.Weinberg,Gravitation and Cosmology(John wiley & Sons,Inc,1972)
- [2]W.Rindler, Am.J.Phys.**34**.1174(1966)
- [3]P.Bergman,Introduction to the Theory of Relativity(Dover Pub. Co.,Inc., New York,1976),Chapter V
- [4]C.Misner, K,Thorne and J. Wheeler, Gravitation(W.H.Freedman & Co.,1973)
- [5]S.Hawking and G. Ellis,The Large Scale Structure of Space-Time(Cam-bridge University Press,1973)
- [6]R.Adler,M.Bazin and M.Schiffer,Introduction to General Relativity(McGraw-Hill,Inc.,1965)
- [7]A.Miller, Albert Einstein's Special Theory of Relativity(Addison-Wesley Publishing Co., Inc., 1981)
- [8]W.Rindler, Special Relativity(2nd ed., Oliver and Boyd, Edinburg,1966)
- [9][Massimo Pauri](#), [Michele Vallisner](#), "Marzke-Wheeler coordinates for accelerated observers in special relativity":Arxiv:gr-qc/0006095(2000)
- [10]A. Einstein, " Zur Elektrodynamik bewegter K"orper", Annalen der Physik. 17:891(1905)
- [11]J.W.Maluf and F.F.Faria,"The electromagnetic field in accelerated frames":Arxiv:gr-qc/1110.5367v1(2011)
- [12]S.Yi, "Expansion of Rindler Coordinate Theory And Light's Doppler Effect", The African Review of Physics,8:37(2013)
- [13]G.F.Torres del Castillio and C.I.Perez Sanchez,"The Maxwell equations in a uniformly accelerated frame", Revista Mexicana De Fisica,53,1,4-9(2007)