

THE REAL-ZEROS OF JONES POLYNOMIAL OF TORUS

CHANG LI

Dedicated to My Parents.

ABSTRACT. This article proved two theorems and presented one conjecture about the real-zeros of Jones Polynomial of Torus. The distribution of zeros of Jones Polynomial is an interesting topic in knots theory of math and physics.

1. THE REAL-ZEROS OF JONES POLYNOMIAL OF TORUS

The real-zeros of Jones polynomial of torus $V_t(p, q)$ are distributed on x axis. In physics, since real-zeros of an equation usually represent observable values, it is interested to investigate them in advanced.

$$(1.1) \quad V_t(p, q) = 1 - t^{p+1} - t^{q+1} + t^{p+q}, t \in \mathbb{C}$$

Theorem 1. *For all Jones polynomials of Torus $V(p, q)$ there are two positive real-zeros with one is $+1$ and another is inside $1 < r < 2$.*

Proof. Suppose a positive real-zero of $V_t(p, q)$ is r where $0 < r < +\infty$, then there is a r so that $V_r(p, q) = 0$. By Cauchy theorem, $\|r\| < 1 + \max(\|a_i\|) \leq 2$. We can represent $V_r(p, q) = 0$ as $(1 - r^p)(1 - r^q) = (r - 1)(r^p + r^q)$. If we suppose $0 < r < 1$, then $0 < r^p < 1$ and $0 < r^q < 1$, the left side of equation is positive, but since $r - 1 < 0$, the right side is negative. So there is $1 \leq r < 2$.

By Descartes' rule of signs, there are two signs change in the coefficients of equation (6), so there are two positive zeros. Obviously, $+1$ is a zero, another positive real-zero is in $1 < r < 2$. \square

Theorem 2. *For all Jones polynomials of Torus $V(p, q)$, for both p and q are positive even integers, there are no negative zeros; for one of p or q is odd and another is even, there is one negative zero; for both p and q are odd, there are two negative zeros.*

Proof. Replace $-t$ in Jones polynomial $V_t(p, q)$ as $V_{-t}(p, q)$ Descartes's rule can be applied to decide all the negative real-zeros.

* Suppose $p = 2m, q = 2n$, then $V_{-t}(p, q) = 1 + t^{p+1} + t^{q+1} + t^{p+q}$. Because no signs change, there are no negative real-zeros.

* Suppose $p = 2m + 1, q = 2n$, $V_{-t}(p, q) = 1 - t^{p+1} + t^{q+1} - t^{p+q}$. Because $V_{-t}(p, q) = V_{-t}(q, p)$, it is also true for $p = 2m, q = 2n+1$.

Date: January 25, 2017.

Key words and phrases. real zeros, jones polynomial, torus.

* Suppose $p = 2m + 1, q = 2n + 1$, then $V_{-t}(p, q) = 1 - t^{p+1} - t^{q+1} + t^{p+q}$. It is the case of positive real-zeros, so there are two negative real-zeros -1 and r with $-1 < r < 0$. \square

Conjecture 1. For $p \geq 2, q \geq 3$, $Ze(p, q)$ is the positive real-zero of $V_t(p, q)$ which is not equal to $+1$, then there is approximation formula below

$$(1.2) \quad \frac{\ln(Ze(p, q) - 1)}{\ln(Ze(p, q + 1) - 1)} \approx \frac{\ln(p + q - 3)}{\ln(p + q - 2)}$$

This conjecture shows the relationship of the real-zeros distribution of Jones Polynomial of Torus. Computer software has partially verified this conjecture.

REFERENCES

- [1] Rolfsen, Dale, *Knots and Links*, Providence, R.I. : AMS Chelsea Pub., 2003.
- [2] F. Y. Wu, J. Wang, *Zeroes of the Jones polynomial*, Physica A 296, 483-494 (2001), <https://arxiv.org/abs/cond-mat/0105013>, DOI: 10.1016/S0378-4371(01)00189-3.
- [3] Georgios Giasemidis, Miguel Tierz, *Torus knot polynomials and susy Wilson loops*, Lett. Math. Phys. 104, 1535-1556 (2014), <https://arxiv.org/abs/1401.8171>, DOI: 10.1007/s11005-014-0724-z.

NEATWARE, 2900 WARDEN AVE, P.O.Box 92012, TORONTO, ONTARIO M1W3Y8, CANADA
E-mail address: changli@neatware.com