

Analysis of the HRM spectra in plasma glow discharge

Dr. Stoyan Sarg Sargoytchev

World Institute for Scientific Exploration

Abstract:

The HRM effect (Heterodyne Resonance Mechanism) is predicted in the Basic Structures of Matter – Supergravitation Unified Theory (BSM-SG). In properly activated neutral plasma a discrete frequency spectrum is observed. The term heterodyne means that the process is activated in the kHz range, while the observed spectrum is in the MHz range. The clear spectrum has spatial features and could be observed only in a small volume at fast sweep time. The spectrum characteristics depend on the type of gas and conditions of the transient process. Hydrogen and inert gases are the most suitable for the HRM effect. According to analysis, the HRM spectrum is a signature of synchronized spin-flips of the electrons involved in ion-electron pairs. HRM effect involves quantum mechanical interactions in which the electrons access the zeropoint energy of the physical vacuum. It is predicted that the HRM effect takes place in the transient process of lightning. The signature of HRM effect in the lightning is a subject of another article.

Keywords: *neutral plasma, glow discharge spectrum, Rydberg matter, electron spin flip, zeropoint energy*

1. Analysis of the experimental results published in [6,7]

Fig 1. shows the waveform of a single burst recorded by oscilloscope Rigol DS1102E (the scope lacks of high display resolution). The waveforms are measured by using the vacuum cell with air at pressure of 11.5 mbars. (Fig. 10 of [7]).




Fig. 1. Waveforms of glow discharge of air at 11.5 mbars, a. – burst rate with the HRM oscillations, b. expanded view of the HRM oscillations

The burst frequency is in the range of 400 kHz to 800 kHz (depending on high voltage). The burst is followed by oscillations in MHz range that are signature of the HRM effect.

Fig. 2 shows the measured burst rate dependence on the external inductance for two HV values.




Fig. 2. Burst rate dependence on external inductance L_I for 2 kV and 2.7kV. Vacuum cell with air at 11 mbars.

The waveform in Fig. 1.a shows the burst rate (frequency) in KHz range and the RF frequency of HRM in MHz range. Fig. 4.b shows the HRM in MHz range with higher resolution. The gas was air at 12 mbars. It was observed that there is always a delay between the burst pulse in KHz range and the HRM. The operating HV was between 2 and 3 kV. For many runs the delay is between 20 and 24 μs and the HRM duration between 2 and 4 μs . The burst frequency depends on the external inductance L_I and the HV as shown in Fig. 2. but the delay and the duration of HRM are not much dependent on L_I and HV. The delay exists also for external activation by electrical pulse as shown in Fig. 3 [7].




Fig. 3. HRM patch appears after some delay from the switching pulse

Let us analyze the hydrogen spectrum shown in Fig. 4 [6,7]. The four bands of HRM spectrum probably correspond to ion-electron pair clusters with electron velocities corresponding

to 0.85 eV, 1.51 eV, 3.4 eV and 13.6 eV. The higher intensity band at 11.6 MHz is probably for pairs with electron energy of 13.6 eV (the optimal confined velocity of the electron [4]).




Fig. 4. Four bands of HRM spectrum of hydrogen with identified central frequencies

Fig. 5. shows the HRM spectrum of the strongest band at 11.6 MHz by enhanced spectral resolution.




Fig. 5. RF spectral signature of a supercluster from HRM effect in EM activated plasma of hydrogen at 12 mbars.

Fig. 6. shows oscillations with one and a same frequency of 5.17 MHz obtained by using a tunable resonant circuit [7]. The shape of waveform is like signal with amplitude modulation. It is clearly visible that at the nodes the oscillations change the phase by 180 deg. This could be from electron spin change. The signal intensity is so high over the noise that we may conclude that electron spin change occurs simultaneously in a huge number of ion-electron pairs.





Fig. 6. Expanded section of HRM oscillations of hydrogen gas measured by a digital scope.

The amplitude modulation of the waveform in Fig. 7. indicates that the intrinsic electron energy is not constant. It changes periodically that causes a periodical spin flip of the electrons. This conclusion comes from the waveforms shown in Fig. 6 where the oscillations are externally manipulated by a positive feedback. We see that at strong feedback a gap between patches occurs without oscillations. This means that the electrons exhaust their internal energy due to external manipulation, so they need some in order to restore it to the optimal level. The needed energy they obtain from the ZPE-D energy of CL space.



a.

b.

Fig. 7. Envelope waveforms for different strength of the positive feedback; a. – weak, b. strong

Now let us imagine the spatial arrangement of one supercluster. Its configuration is illustrated in Fig. 8. It is formed by one central cluster and three layers of clusters. Each layers contains six clusters. Clusters of any layer are at the same radius in respect to the central axis. At the same time they are twisted like the threads of a rope. The ion-electron of the whole supercluster move with one and same velocity, corresponding to one of the quantum energies 13,6 eV, 3.41 eV, 1.51 eV, 0.85 eV etc.




Fig. 8. Spatial arrangement of clusters in a supercluster. The clusters of same order are at one and a same radius in respect to the central axis.

The analogy of this configuration is like a solenoid with multiple layers of windings. The magnetic field of such solenoid is illustrated in Fig. 9. The field strength decreases with the radius in respect to the central axis.




Fig. 9. Magnetic field of the single layer solenoid.

The magnetic fields created by ion-electron pairs in the configuration shown in Fig. 8 could be regarded as a multi-layer solenoid shown in Fig. 9 but with some particular differences. The external magnetic field decreases with the radius, but we must consider one additional option: The electrons in the clusters of one and a same layer (see Fig. 8) may not have one and a same rotational direction. In three of the clusters, for example, the rotation could be clockwise, while in another three – a counter clockwise.

Assuming that the observed spectra are result of spin flip of the electrons we will use the expression (1) and (2) of the spin magnetic moments of the electron, μ_{es}

$$\mu_{es} = \pm \frac{eh}{4\pi m_e} \quad (1)$$

$$W = \mu_{es} B = \Delta E \quad (2)$$

where: B is the magnetic field in which the electron moves, W – is the work between two spin energies equal to the energy difference ΔE between the up and down electron spin. From well-known expression $E = hf$ we have the frequency shift proportional to ΔE

$$\Delta f \sim \Delta E \quad (3)$$

The graphical illustration shown in Fig. 10 helps understanding the observed spectra.




Fig. 10. Graphical illustration of the physical parameters behind the observed frequency spectrum due to electron spin flip. The upper coordinates are in the frequency domain, while bottom coordinates $+B$ and $-B$ are shown as a function of r , but r for $+B$ and $-B$ are shown in opposite direction in order to match the spectrum domain.

The upper part in Fig. 9 is the spectrum of one supercluster, while the bottom part shows the dependence of the frequencies on the radius of the clusters with one and a same radius referenced to the central axis of the supercluster. The magnetic field shown by gray lines decreases with the cluster radius (in analogy with the magnetic field solenoid shown in Fig. 8). According to (2) and (3) the frequency shift is proportional to the strength B of the common magnetic field. For this reason B decreases with the radius r of the cluster. (). When referenced to the central frequency f_c , the positive frequency shifts corresponds to a magnetic field with a direction $+B$, while the negative shifts - to the magnetic field with an opposite direction $-B$. These means that in one and a same cluster half of the electrons may have, for example, a clockwise rotation, while the other half - a counter-clockwise rotation. The central frequency should be due to at largest radius of the cluster where B is small. To test this a serial resonance frequency was included in a serial connection with the circuit shown in [7, Fig 9]. By tuning the resonant circuit the central peak was separated with higher resolution, while other frequencies are suppressed. The observed high resolution spectrum of the central frequency is shown in Fig. 11.




Fig. 11. Central frequency with high spectral resolution showing two close frequencies

One additional question should be answered: Why all frequency shifts are referenced to the central line? The reasonable answer is: The central frequency is stable because the magnetic field creating by the rotating electron is much stronger than the magnetic field from the electron spin, so it serves as a base for the frequency shifts. The difference between frequency intensities in the supercluster could be explained by the following consideration. The central fc frequency emitted from the central cluster must pass through the external clusters and it is known that EM waves are attenuated when passing through a plasma. The EM waves from external cluster will not exhibit such attenuation and therefore they will appear with stronger intensity. Note that the stronger magnetic field provides a larger frequency shift, not a stronger intensity of the frequencies.

All above mentioned considerations explain the symmetry of the observed spectra shown in Figures 4 and 5. However there are some additional features of the ion-electron superclusters that are apparent from the spectra shown in Figures 12,13 and 14. These spectra of hydrogen are made at one and a same, HV and pressure values and the same sweep time. However, we see that the position of the central frequency is the same but some frequencies are missing or shifted at the consecutive spectra runs. If ignoring the missing frequencies the positions in the frequency domain are still symmetrical. The possible explanation is follows: The missing spectral lines could be a result of the way the magnetic fields of clusters at one and a same radius are interconnected. They could be paired either to the opposite clusters or to the nearby cluster (lying at one and a same radius in respect to the supercluster axis). Their interconnections will influence the common magnetic field configuration that will become more complex than the simple analogy with the multilayer solenoid.




Fig. 12. Hydrogen spectrum, 20 ms sweep time [6]




Fig. 13..Hydrogen spectrum, 50 ms sweep time [6]




Fig. 14. Hydrogen spectrum, 50 ms sweep time [6]

2. Conclusions from the experimental analysis of the HRM effect invoked in a plasma glow discharge

- a. The HRM effect takes place after some delay from activation electrical pulse (in order of 20 uS for air at 13 mbars). This time is necessary for ionization and formation of ion-electron pair.
- b. The duration of the HRM oscillations is about 2 to 4.5 uS (for air at 13 mbars).
- c. During HRM mechanism electron tends to move with one of optimal quantum velocity, while the common magnetic field forces a synchronization that affects the internal energies of the electron, so they could replenish their energy by spin flipping.
- d. The spin flipping of a huge number of electrons in oscillating ion-electron pairs is synchronized due to the common magnetic field.
- e. In spin flipping the electron interacts with the superhigh Compton frequency taking that is a quantum mechanical feature of the electron
- f. The synchronized spin flip of enormous number of electrons takes a fraction of the zeropoint energy of physical vacuum denoted as ZPE-D in BSM-SG theory (Dyamic type of Zero Point energy) [8].

The main questions are: where is the fraction of the accessed ZPE-D energy and how it could be obtain by external means?

The ZPE-D accessed by electrons is in the oscillating ion electron pairs. In order to obtain this energy by external means the ion-electron pairs must be destroyed. The released energy will be taken from the free ions, so they will have larger momentums in comparison to their momentums at the time of ionization. Their integral momentum containing the accessed ZPE-D energy could be used either for obtaining of electrical potential or used directly for expansion of the gas. For the second option the possibility for directing the axis of the superclusters with external magnetic field could be used. Obviously it will lead to a push force different from the explosion force caused by the chemical reaction. Consequently:

- g. The oscillating ion-electron pair after a specific time of their creation must be destroyed in order to free the accessed zeropoint energy.
- h. The expansion of the gas due to a forced destructuin of the ion-electron pairs is not a thermodynamical process.

3. Analysis of prior art research on Rydberg matter showing the signature of HRM effect.

Extensive research on Rydberg matter was made by Leif Holmlid. In his article [9] he says Excitation levels n , nB with shorter interatomic distances exists for hydrogen clusters and distances down to 140 pm are now measured with this light-scattering method. The clusters studied have maximum dimentions from 0.3 nm up to tens of nm, often beenig planar. In another articl Leif Holmlid detects energetic ions [10].

In another article by F. Olofson, P. Andersson and L. Holmlid, about Rydberg matter clusters in polar mesospheric clouds it is found: Such experiments show that RM clusters interact strongly with radar frequencies: this explains the radio frequency heating and reflection studies of PMSE layers. [11]. In the same article it was also found that the time between collisions with other RM clusters is . 6 h. In other article S. Badiei and L. Lolmlid, presents s study of hydrogen Rydberg matter (RM) by pulsed-lase-time-of-flight. It was found that between four and six photons with a total energy of 8.8 – 13 eV take part in the RM fragments.

In his manuscript of “A problem in Plasma Science” Dr. Harold Apsden put emphasise on anomalous acceleration forces exerted by electrons on ions. He discussed this issue in his article in IEEE Transaction on Plasma Science [12].

In the article “Anomalouse Electron-Ion energy Transfer in a relativistic-Electron-Beam-Heated Plasma”, J. Sethian e al., describe an experiment to study the previous observations that the beam-to-plasma energy transfer is far faster than can be explained by classical processes. [13]. In a plasma column they inject an electron beam, with parameters $v = 350$ kV, $I = 22$ kA, $t = 60$ ns. They claim: experimental evidence is presented for an observed anomalous electron-ion energy transfer in a magnetized beam-heated plasma that is approximately 103 times faster than classical. They claim a rapid decrease in plasma density during the first 2 usec after beam injection. In fact this is anomalous process of plasma expansion not explainable by the rules of classical thermodynamics.

Yu Astrov et al. observed quite interesting formations of clusters [14]. They make a gas discharge device formed of two parallel perfectly flat silicon disks with diameter 20 mm doped with gold or zinc to behave as a spatially distributed resistive load. A photo resistive layer is deposited for changing the resistance by illumination. Transparent metal electrodes are deposited to which a DC voltage in order of a few kilovolts is applied. The gap between disks with size of 1 mm is filled by nitrogen at pressure of 120 hPa. The device is transparent permitted observing the discharge pattern while the resistivity can be controlled by illumination. The device is cooled to liquid nitrogen temperature of 90K. Applying a high voltage causes a discharge pattern of spatially separated clusters that are optically recorded. The most stable pattern is hexagonal. In Fig. 2 of the article they show two of the observed patters with the following description: The hexagon (initial pattern evolves from the homogeneous state by increasing the supply voltage, a further increase of the voltage is accompanied by the transformation of the hexagon into the stripe pattern. In another article of Astrov et al., with better optical resolution they observe a zigzag pattern instead of clear stripe [14]. The observaions matches the specific features of the superclusters from ion-electron pairs.

The prior art experimental data are in full agreement with the theoretical concept of the HRM effect and the obtained experimental results. Generally they confirm the theoretical concept of HRM and additionally show some spatial features:

4. Conclusions:

- a. The observed pattern in [14] is a signature of superclusters formations
- b. The superclusters are aligned along the sight of observation and evenly separated. The most stable pattern is a hexagonal. Evidently the complex individual magnetic field of the superclusters keeps them separated.
- c. The appearance of zigzag pattern indicates that the magnetic field configuration of the superclusters is complex, but it is the same for all superclusters during their time of existence.
- d. The experiments with injected high current short pulse show an anomalous effect of plasma expansion [13].

The conclusions from §2 and §3 serve as a physical base for definition of the technical considerations for using of the HRM effect for accessing the zeropoint energy of the physical vacuum.

It is evident that further research on the HRM effect will be quite useful. It could provide more information about the HRM effect dependence of type of gas, HV activation, and gas pressure. I expect that if the vacuum cell is longer and the glow discharge appear separated into light and dark zone the spectrum could appear different. If these zones correspond to not synchronized superclusters the spectrum will not be of well separated discrete frequencies with equal space intervals. Therefore, it will be useful to study the spectral and spatial characteristics simultaneously. This will lead to deeper understanding the physics of lightning phenomena. In lightning between clouds and ground and enormous energy is released that many think does not come entirely from the accumulated electrical potential in clouds. This is a subject of another article.

References:

1. S. Sarg ©2001, Basic Structures of Matter, monograph, <http://www.nlc-bnc.ca/amicus/index-e.html> (First edition, ISBN 0973051507, 2002; Second edition, ISBN 0973051558, 2005), (AMICUS No. 27105955), LC Class no.: QC794.6*; Dewey: 530.14/2 21
2. S. Sarg, New approach for building of unified theory <https://arxiv.org/abs/physics/0205052> (2002)
3. Stoyan Sarg, Basic Structures of Matter –Supergravitation Unified Theory, Trafford Publishing, 2006, ISBN 1412083877 (books review in "Physics in Canada," 62, No. 4, July/Aug, 2006).
4. S. Sarg, A Physical Model of the Electron according to the Basic Structures of Matter Hypothesis, Physics Essays, vol. 16 No. 2, 180-195, (2003); <https://www.ingentaconnect.com/content/pe/pe>
5. Stoyan Sarg, Structural Physics of Nuclear Fusion with BSM-SG atomic models, ISBN 9781482620030, Amazon.com, (2013)

6. Stoyan Sarg, Atlas of HRM spectra (volume 1). <http://vixra.org/abs/1903.0523>
7. Stoyan Sarg, Heterodyne Resonance mechanism in a transient process in plasma. Experimental study and spectra. <http://vixra.org/abs/1903.0520>
8. Stoyan Sarg, Zero Point Energy from the Viewpoint of an Alternative Concept of Space According to the BSM-Supergravitation Unified Theory , <http://vixra.org/pdf/1110.0071v1.pdf>
9. L. Holmlid, Sub-nanometer distances and cluster shapes in dense hydrogen and in higher levels of hydrogen Rydberg matter by phase-delay spectroscopy, , J. Nanoparticle Research, (2011) 13:5536-5546
10. L. Holmlid, High-energy Coulomb explosions in ultra-dense deuterium: Time-of-flight-mass spectroscopy with variable energy and flight length, International journal of mass spectroscopy 282 (2009) 70-76.
11. L. Holmlid, Rydberg Matter clusters of alkali metal atoms: the link between meteoritic matter, polar mesosphere summer echoes (PMSE), sporadic sodium layers, polar mesospheric clouds (PMCs, NLCs), and ion chemistry in the mesosphere, (2010) <https://arxiv.org/abs/1002.1570>
12. H. Apsden, IEEE Transactions on Plasma Science, PS-5, 159 (1977).
13. J. D. Sethian, D. A. Hammer and C. B. Wharton, Anomalous Electron-Ion Energy Transfer in a Relativistic-Electron-Beam-Heated Plasma, Physical Review Letters, 40, Number 7, (1978).
14. Yu. Astrov, E. Ammelt, and H. –G. Purwins, Experimental Evidence for Zigzag Instability of Solitary Stripes in a Gas Discharge System, Physical Review Letters, 78, 16, (1997), 3129-3132.