

Affirmative resolve of the Riemann Hypothesis

T.Nakashima

Abstract

Riemann Hypothesis has been the unsolved conjecture for 170 years. This conjecture is the last one of conjectures without proof in (B.Riemann). The statement is the real part of the non-trivial zero points of the Riemann Zeta function is $1/2$. Very famous and difficult this conjecture has not been solved by many mathematicians for many years. In this paper, I try to solve the proposition about the Möbius function equivalent to the Riemann Hypothesis. First, the non-trivial formula for Möbius function is proved in theorem1. In theorem3, I think this formula into 2 parts. By calculation for the latter part, I get upper bound for the sum of the Möbius functions (for meaning of R.H. See theorem3).

1

Handles propositions equivalent to the Riemann Hypothesis. I express the Riemann Hypothesis as R.H, and the Möbius function as $\mu(n)$.

Next theorem is well-known

Theorem

$$\sum_{n=1}^m \mu(n) = O(m^{\frac{1}{2}+\epsilon}) \Leftrightarrow R.H$$

I will prove Left hand formula.

Lemma 1.

$$\sum_{n|m} \mu(n) = 1(m=1), \sum_{n|m} \mu(n) = 0(m \neq 1)$$

Proof. First, if $m = 1$, it is $\sum_{n|m} \mu(n) = \mu(1) = 1$. Second case. There is a little explanation for this. Let m 's prime factorization be $m = p_1^{n_1} p_2^{n_2} p_3^{n_3} \cdots p_k^{n_k}$.

Then it becomes $\sum_{n|m} \mu(n) =_k C_0 -_k C_1 +_k C_2 -_k C_3 + \cdots_k C_k = (1-1)^k = 0$. \square

Theorem 1.

$$\sum_{n \leq m} \mu(n) \left[\frac{m}{n} \right] = 1$$

Proof. $\sum_{m'=1}^m \sum_{n|m'} \mu(n) = 1$ is from Lemma1

$$1 = \sum_{m'=1}^m \sum_{n|m'} \mu(n) = (\mu(1)) + (\mu(1) + \mu(2)) + (\mu(1) + \mu(3)) \\ + (\mu(1) + \mu(2) + \mu(4)) + \cdots$$

See $\mu(n)$ in this expression as a character. $\mu(1)$ appears m times in the expression. $\mu(2)$ appears $\left[\frac{m}{2}\right]$ times that is a multiple of 2 less than m . In general, the number of occurrences of $\mu(n)(n < m)$ in this expression is the number $\left[\frac{m}{n}\right]$ that is a multiple of n below m . I get $\sum_{n \leq m} \mu(n) \left[\frac{m}{n} \right] = 1$. \square

example

$m = 10$ case, $10 - 5 - 3 - 2 + 1 - 1 + 1 = 1$. $m = 13$ case, $13 - 6 - 4 - 2 + 2 - 1 + 1 - 1 - 1 = 1$ etc..

Theorem 2.

$$\sum_{n=1}^x \mu(n) \text{ changes sign at } n_0 \in [m^{\frac{1}{2}(1-\epsilon')}, m^{\frac{1}{2}}] (m > \exists m_\epsilon)$$

$$\sum_{n=1}^x \frac{m}{n} \mu(n) \text{ changes sign at } n' \in [m'^{(1-\epsilon')}, m'] (m > m' > \exists m_{\epsilon'})$$

Proof. $\sum_{n=1}^x \mu(n)$ changes sign in the interval $[m^{\frac{1}{2}(1-\epsilon')}, m^{\frac{1}{2}}]$, $m > \exists m_{\epsilon'}$ ([11]). $\sum_{n=1}^x mn\mu(n)$ changes sign in the interval $[m'^{(1-\epsilon')}, m']$, $m > m' > \exists m_{\epsilon'}$ ([11]). \square

Theorem 3.

$$\left| \sum_{n=1}^m \mu(n) \right| < K m^{\frac{1}{2}+\epsilon}$$

R.H. is got.

Proof. From theorem1

$$\sum_{n \leq n_0} \mu(n) \left[\frac{m}{n} \right] + \sum_{n_0 < n \leq m} \mu(n) \left[\frac{m}{n} \right] = 1$$

(By theorem2 $m^{\frac{1}{2}(1-\epsilon')} < n_0 < m^{\frac{1}{2}}$ is the point satisfies $\sum_{n \leq n_0} \mu(n) = 0$. I take ϵ as $\epsilon \approx \frac{1}{2}\epsilon'$.

The following is obtained by calculation for $\sum_{n_0 < n \leq m} \mu(n) \left[\frac{m}{n} \right]$. This represents the terms corresponds to $[\sqrt{m}]$ are \sqrt{m} to $m/(\sqrt{m} - 1)$, the terms corresponds to $[\sqrt{m}] - 1$ are $m/(\sqrt{m} - 1)$ to $m/(\sqrt{m} - 2)$ and the terms corresponds to 1 are $\frac{m}{2}$ to m .

$[\sqrt{m}]$ term is sum of all terms satisfy $\left[\frac{m}{n} \right] = [\sqrt{m}] - 1$, $m/\sqrt{m} = \sqrt{m} \geq [\sqrt{m}]$ and $m/(m/(\sqrt{m} - 1)) = \sqrt{m} - 1 \geq [\sqrt{m} - 1]$, $(m/(m/(\sqrt{m} - 1)) + 1) = m(\sqrt{m} - 1)/(m + \sqrt{m} - 1) < \sqrt{m} - 1$, so the range is \sqrt{m} to $m/(\sqrt{m} - 1)$. Next term is sum of all terms satisfy $\left[\frac{m}{n} \right] = [\sqrt{m}] - 2$, $m/(m/(\sqrt{m} - 2)) \geq [\sqrt{m} - 2]$. The range is $m/(\sqrt{m} - 1)$ to $m/(\sqrt{m} - 2)$. The last term satisfy $\left[\frac{m}{n} \right] = 1$, that is $\frac{m}{2}$ to m .

$$\begin{aligned} \sum_{n_0 < n \leq m} \mu(n) \left[\frac{m}{n} \right] &= ([m/(n_0)] - 1) \times \sum_{m/([m/(n_0)]) < n \leq m/([m/(n_0)] - 1)} \mu(n) + \dots + \\ &([[\sqrt{m}]] \times \sum_{m/(\sqrt{m}+1) < n \leq m/\sqrt{m}} \mu(n) + ([\sqrt{m}] - 1) \times \sum_{\sqrt{m} < n \leq m/(\sqrt{m} - 1)} \mu(n) + ([\sqrt{m}] - 2) \times \\ &\quad \sum_{m/(\sqrt{m} - 1) < n \leq m/(\sqrt{m} - 2)} \mu(n) + \dots + 1 \times \sum_{m/2 < n \leq m} \mu(n) \end{aligned}$$

By induction for Riemann Hypothesis, $|\sum_{n \leq m/N} \mu(n)| < K(\frac{m}{N})^{\frac{1}{2}+\epsilon}$. I get $|\sum_{m/(N+1) < n \leq m/N} \mu(n)| < K(\frac{m}{N+1})^{\frac{1}{2}+\epsilon} + K(\frac{m}{N})^{\frac{1}{2}+\epsilon}$,

I want to calculate some terms. $|\sum_{m/2 \leq n \leq m} \mu(n)| < K(m - 1)^{\frac{1}{2}+\epsilon} + 1 + K(\frac{m}{2})^{\frac{1}{2}+\epsilon}$, $|\sum_{m/3 \leq n \leq m/2} \mu(n)|$ is less than $K(\frac{m}{3})^{\frac{1}{2}+\epsilon} + K(\frac{m}{2})^{\frac{1}{2}+\epsilon}$. $|\sum_{\sqrt{m} < n \leq m/(\sqrt{m} - 1)} \mu(n)|$ is less than $K(\frac{m}{[\sqrt{m}]})^{\frac{1}{2}+\epsilon} + K(\frac{m}{[\sqrt{m}] - 1})^{\frac{1}{2}+\epsilon}$

Later, I calculate in the real examples.
example: $m = 100$ case.

$$-6 = 10 - 9 + 0 + 0 + 6 - 5 \times 2 - 4 - 3 + 2 + 1 \times 4$$

$1 \times 4 - 4 + 2 - 3, 4 - 1 + 1 - 1 = 3$ give the almost value of $\sum_{[100/n_0] < n \leq 100} \mu(n)$. Actually, $\sum_{9 < n \leq 100} \mu(n) = 2$. This gives $|\sum_{[100/n_0] < n \leq 100} \mu(n)| < K[100/n_0] =$

$K \times 10$,
example: $m = 10000$ case.


$$-95 = \dots + 3 \times 18 - 2 \times 15 - 1 \times 25$$

$-1 \times 25 + 3 \times 8 - 2 \times 15 + 3 \times 10, -25 + 8 - 15 + 10 = -22$ gives the almost value of $\sum_{94 < n \leq 10000} \mu(n) = -22$. This gives $|\sum_{[100/n_0] < n \leq 100} \mu(n)| < K[100/n_0] = K \times 107$.


Why does the value coincide? By theorem 2, $\sum_{n_0 \leq n \leq n_1} \mu(n) = 0 \Leftrightarrow \sum_{n_0 \leq n \leq n_1} [\frac{m}{n}] \mu(n) \approx 0, n_1 \ll m$.


3 terms case cover all pattern of all possible case about.

For example, $26 \times 1 - 10 \times 2 - 2 \times 3$ case. $+ - - = 0$ case.


 $(A < K(m-1)^{\frac{1}{2}+\epsilon} + 1, B < (\frac{m}{2})^{\frac{1}{2}+\epsilon})$
 $A+B = \frac{(2\alpha+3\beta)}{\alpha+\beta}B$ holds. This is $A = \frac{(\alpha+2\beta)}{\alpha+\beta}B$. I suppose $\alpha \neq 0$ and $A < \sqrt{3}B$.
 $A = \sum_{n_0 < n \leq m} \mu(n) \ll Km^{\frac{1}{2}+\epsilon}$. Later, I see $\alpha = 0$ case.

For example, $17 \times 1 + 5 \times 2 - 9 \times 3$ case. $+ + - = 0$ case.


 $(A+B) \frac{(\alpha+2\beta)}{\alpha+\beta} = 3B$ holds. This is $A = B(3 \frac{(\alpha+\beta)}{\alpha+2\beta} - 1)$. I suppose $\beta \neq 0$ and $A < \sqrt{3}B$.
 $A = \sum_{n_0 < n \leq m} \mu(n) \ll Km^{\frac{1}{2}+\epsilon}$.
 $\beta = 0$ or the former $\alpha = 0$ case.


 $\sum_{n \leq m} \mu(n) < 2K(\frac{m}{3})^{\frac{1}{2}+\epsilon}$ at best. But like later example, $+ - + = 0$ or $- + - = 0$ for 3,4,5 terms, (Later example, 2,3,4 terms into calculation. 2 term is 0.) I get $B < K((\frac{m}{4})^{\frac{1}{2}+\epsilon})$. $\sum_{n \leq m} \mu(n) < 2K(\frac{m}{4})^{\frac{1}{2}+\epsilon}$. I get $\sum_{n \leq m} \mu(n) < Km^{\frac{1}{2}+\epsilon}$

For example, $12 \times 1 - 12 \times 2 + 4 \times 3$ case. $+ - + = 0$ case.

$$\begin{aligned} \max\{ & (K(\frac{m}{2})^{\frac{1}{2}+\epsilon} + K(m-1)^{\frac{1}{2}+\epsilon} + 1)(1 - \frac{1}{2}), \\ & -(-K(\frac{m}{3})^{\frac{1}{2}+\epsilon} - K(\frac{m}{2})^{\frac{1}{2}+\epsilon})(1 - \frac{2}{3}) \} \\ & = (K(\frac{m}{2})^{\frac{1}{2}+\epsilon} + K(m-1)^{\frac{1}{2}+\epsilon} + 1)(1 - \frac{1}{2}) < Km^{\frac{1}{2}+\epsilon} \end{aligned}$$

I can take 4 or more terms.

example:m=15000

$$-25 + 13 \times 2 + 9 \times 3 - 1 \times 4 - 5 \times 5 \approx 0$$

But,

$$-25 + 13 \times 2 + 0 \times 3$$

give almost value $\sum_{n \leq m} \mu(n)$. And latter terms are enough small, so $\sum_{n \leq m} \mu(n) < K m^{\frac{1}{2}+\epsilon}$ is got.

By calculation result, 4 or more latter terms' influence are gradually small.
First some (≥ 3) terms decide all value.

The calculation key is $+ - + = 0$ or $- + - = 0$. By theorem2, $\sum_{n=1}^x \frac{m}{n} \mu(n)$ takes 0 frequently. This supports the calculation.

example

m=10000 case.

$$\begin{aligned} -95 &= 107 + 106 + 105 + 0 - 103 + 0 + 0 + 0 - 99 - 98 - 97 + 0 - 95 + 94 - 93 + 0 - 91 \\ &\quad + 0 + 0 - 88 - 87 + 86 + 0 + 0 + 84 \times 2 + 0 + 0 + 81 \times 2 + -0 + 0 - 78 + 77 \\ &\quad - 76 \times 2 + 75 + 74 + 0 - 72 \times 2 - 71 + 70 \times 2 + 69 + 68 \times 2 - 67 - 66 + 0 + 0 + 0 \\ &\quad + 62 \times 2 - 61 + 0 - 59 - 58 - 57 \times 2 + 56 \times 2 - 55 \times 2 + 54 + 0 - 52 \times 2 - 51 - 50 \times 2 \\ &\quad + 49 \times 3 + 48 \times 2 + 47 + 46 \times 4 + 45 \times 2 - 44 - 43 \times 3 + 0 - 41 \times 2 + 40 + 0 - 38 \\ &\quad + 0 - 36 \times 2 - 35 \times 3 - 34 + 33 \times 6 - 32 - 31 + 30 \times 4 + 29 \times 3 - 28 \times 4 - 27 + 0 \\ &\quad + 25 \times 3 + 0 - 23 \times 6 + 22 - 21 \times 2 + 0 + 19 \times 4 + 18 \times 7 + 17 - 16 \times 9 - 15 \times 9 + 14 \times 9 \\ &\quad + 0 + 0 + 11 \times 2 + 10 \times 3 - 9 \times 15 + 8 \times 10 + 7 \times 12 - 6 \times 20 + 5 \times 16 - 4 \times 6 + 3 \times 18 - 2 \times 15 \\ &\quad - 1 \times 25 \end{aligned}$$

By calculation, (mobius function's property, I vanish some terms.)

$$-1 \times 25 - 2 \times 15 + 3 \times 18$$

I get $-25 - 15 + 18 = -22$. $- + - = 0$ is well taken as,

$$-4 \times 6 + 5 \times 16 - 6 \times 10$$

This part's the sum of mobius function is $-6 + 16 - 10 = 0$. Next, $+ - + = 0$ is well taken as

$$7 \times 2 + 8 \times 10 - 9 \times 15 + 10 \times 4$$

This part's the sum of mobius function is $2 + 10 - 15 + 4 = 1$. Next $-+ - = 0$ is well taken as

$$-16 \times 8 + 17 + 18 \times 7 + 19 \times 4 - 21 \times 4$$

This part's the sum of mobius function is $-8 + 1 + 7 + 4 - 4 = 0$. Mobius function's partial sum is gradually small.

Mobius function's partial sum is $-22 + 1 = -21$. It is the 3 terms' case.

I get enough result.

$$\left| \sum_{n \leq m} \mu(n) \right| < 0 + Km^{\frac{1}{2} + \epsilon}$$

R.H. is got. □

Special thanks: I was very grateful to my friend H. Tokitu for translating in English. I would like to express my gratitude to him.

References

- [1] B. Riemann, "Ueber die Anzahl der Primzahlen unter einer gegebenen Grosse", Monatsberichte der Berliner Akademie. In Gesammelte Werke, Teubner, Leipzig (1892)
- [2] de Branges, Louis, "The convergence of Euler products", Journal of Functional Analysis 107 (1)(1992)122-210
- [3] Connes, Alain, "Trace formula in noncommutative geometry and the zeros of the Riemann zeta function", Selecta Mathematica. New Series 5 (1)(1999)29-106
- [4] Connes, Alain, "Noncommutative geometry and the Riemann zeta function", Mathematics: frontiers and perspectives, Providence, R.I.: American Mathematical Society, (2000)35–54
- [5] Conrey, J. B., J. Reine angew. More than two fifths of the zeros of the Riemann zeta function are on the critical line", Conrey, J. B., J. Reine angew. Math. 399(1989)1-16
- [6] Conrey, J. Brian, "The Riemann Hypothesis" Notices of the American Mathematical Society(2003)341-353
- [7] Conrey,J.B.; Li,Xian-Jin, "A note on some positivity conditions related to zeta and L-functions", International Mathematics Research Notices 2000 (18)(2000)929-940

- [8] Dyson, Freeman, "Birds and frogs", Notices of the American Mathematical Society 56 (2)(2009)212-223
- [9] Hardy, G. H. "Sur les Zéros de la Fonction $\zeta(s)$ de Riemann", C.R.Acad. Sci. Paris 158(1914) 1012-1014
- [10] Hardy,G.H.;Littlewood,J.E., "The zeros of Riemann's zeta-function on the critical line", Math. Z. 10 (3-4)(1921)283-317
- [11] I Kaczorowski and Pintz, Oscillatory properties of arithmetical functions. Acta Math. Hungar. 48 (1986)173-185
- [12] Littlewood,J.E., "The Riemann hypothesis", The scientist speculates: an anthology of partly baked idea, New York: Basic books(1962)
- [13] Monrtgomey,Hugh L., "The pair correlation of zeros of the zeta function", Analytic number theory, Proc. Sympos. Pure Math., XXIV, Providence, R.I.: American Mathematical Society(1973)181-193
- [14] Montgomery, Hugh L., "Zeros of approximations to the zeta function", in Erdős, Paul, Studies in pure mathematics. To the memory of Paul Turán, Basel, Boston, Berlin: Birkhäuser(1983)497-506
- [15] Sarnak, Peter, in Borwein, Peter; Choi, Stephen; Rooney, "Problems of the Millennium: The Riemann Hypothesis", Brendan et al. (PDF)(2008)