

# **A power sequence model as a bridge between fundamental natural constants and macroscopic liquid systems**

Otto G. Piringer

FABES Forschungs-GmbH, Schragenhofstr. 35, D-80992 München

Fax: 089 149009 80

e-mail: [otto.piringer@fabes-online.de](mailto:otto.piringer@fabes-online.de)

## **Abstract**

An interaction function, with the limit value  $w = e^{2\pi/e}$ , derived as basis for modelling emergent properties of nanoparticles, nanodrops and liquids, has been further developed. A correlation between physical systems of uniform particles and fundamental properties of natural numbers leads to a model, which acts as a bridge between properties of macroscopic systems and fundamental physical constants.

The four fundamental coupling constants, and the masses of fundamental elementary particles, including the Higgs- and Z-bosons, are derived. A proposal for the mass of the electron neutrino is given. Finally, the nuclear binding interaction is modeled with the interaction function. Only the experimental values of the *Planck* constant  $h$  and the speed of light  $c$  are needed for the model.

## **1. Introduction**

Recently an interaction function for liquid particles has been published [1,2]. For this interaction function a system of uniform particles with finite energies, as a basic assumption of quantum mechanics, has been correlated with the system of natural numbers and for  $x$  interacting particles the function

$$y = x^{q/x} \quad (1)$$

with the maximum values:

$$y_{max} = 3^{q/3} \text{ for } x = 3 \text{ and } Y_{max} = e^{q/e} > y_{max} \text{ for } x \gg 1 \text{ and } (1+1/x)^x = e = 2.71828\dots$$

results.

Using the de Broglie relation for linear momentum,  $p = h/\lambda$  of the particle, along with the Planck constant  $h$  and a wavelength  $\lambda$ , a second basic assumption of quantum mechanics has been applied and with a relative wavelength  $\lambda = 2\pi = q$ , the following power sequence is defined [2]:

with:

$$w_{n,e} = \left(1 + \frac{2\pi}{n}\right)^{\frac{n}{e}}, \quad (2)$$

with the limit value:

$$\lim_{n \rightarrow \infty} w_{n,e} = e^{2\pi/e} = 10.08909\dots \approx \frac{2\pi}{w_1 - w_{1,e}} = 10.0898\dots,$$

and

$$w_1 = (1+2\pi)^{1/2} = 2.6987\dots \text{ and } w_{1,e} = (1+2\pi)^{1/e} = 2.076\dots$$

The power sequence in Equation (2) is denoted as interaction function.

The functions  $y_{max}$  and  $Y_{max}$  in Equation (1) suggests two different kinds of interactions between physical particles.

As shown in [1], properties of liquids can be modelled with relations based on the interaction function in Equation (2). The interactions are all of electromagnetic nature.

Most properties of macroscopic systems are described with exponential functions. Positive exponents stand for high kinetic values, like for example diffusion or reaction constants. Negative exponents are responsible for strong attractive interactions in the matrix, like the activation energy of diffusion in polymers and of chemical reactions as two examples.

The maximum  $y_{max} = 3^{q/3}$  for 3 interacting particles in Equation (1), refers to the first term,  $w_I = (1+2\pi)^{1/2}$  with  $n=1$  in the power sequence:

$w_n = \left(1 + \frac{2\pi}{n}\right)^{\frac{n}{(1+1/n)^n}}$  with the same limit  $w$  as in Equation (2). The exponent  $n/(1+1/n)^n$  approaches to  $e = 2.718$  with increasing  $n$ .  $n/(1+1/n)^n = 2.59$  for  $n = 10$  and  $2.705$  for  $n = 100$ .

The value  $w_I$  represents the relative energy of one elementary particle. In conformity with Equation (1) a maximum of interaction results with a combination of 3 elementary particles and this leads to

$$y_{max} = w_I^3 = 19.6554 = 3^{q/3}, \text{ with a value } q = 8.1330397 \text{ and } q/3 = 2.711\dots \approx e.$$

As shown later, the strong force is responsible for the interaction between three elementary particles, with  $y_{max} = w_I^3$ .

The three interacting particles have a rest mass,  $m_0$  and in conformity with the Einstein Equation.  $E = mc^2$ , a rest energy  $E_0 = m_0c^2$ , with the speed of light  $c$ .

Now, with  $E_I^3 = 1/[exp(w_I^3) \cdot w_I^3] = 1.48 \cdot 10^{-10}$ , the relative rest energy is defined as coming from the three interacting particles  $w_I$ . The negative exponent in  $E_I^3$  shows the strong attractive interaction resulting from the three particles.

In order to transform the relative energy value  $w_I$  into a quantity expressed in IS units the limit value

$\lim_{n \rightarrow \infty} w_n = \lim_{n \rightarrow \infty} \left(1 + \frac{2\pi}{n}\right)^{\frac{n}{(1+1/n)^n}} = e^{2\pi/e}$  is related in this case to the decimal system with the basic number 10. In this way  $w_1 \cdot w/10$  represents the energy unit in J.

That means, the rest energy

$$E_0 = E_1^3 \cdot w/10 \text{ J} = 1.49326 \cdot 10^{-10} \text{ J}$$

and with  $c = 2.99792458 \cdot 10^8 \text{ ms}^{-1}$ , the atomic mass unit  $m_0 = u_0$  results:

$$u_0 = E_0/c^2 = (1/[\exp(w_1^3) \cdot w_1^3])w/10c^2 = 1.661475 \cdot 10^{-27} \text{ kg} \approx 1.66054 \cdot 10^{-27} \text{ kg} = u \text{ [3]}. \quad (3)$$

The interaction function is of fundamental importance for emergent properties in fluid phases and the first term  $w_1$  forms a bridge between the fundamental atomic mass unit and macroscopic properties.

With the Avogadro Constant  $N_A$  and the atomic mass unit, the atomic mass for hydrogen is:

$$A_H = 6.022 \cdot 10^{23} \cdot \text{mol}^{-1} \times 1.661 \cdot 10^{-27} \text{ kg} = 1 \text{ g mol}^{-1}. \text{ The relative atomic mass for hydrogen is } H = \underline{1}.$$

This connection explains the importance of the relative mass values  $M$  of particles in macroscopic systems and of subparticles (atomic groups) in the molecules of the system.

As shown later, Equation (1) should be taken into account for all individual particles, independent of their nature, with the gravitational interaction as an example.

## 2. A general interaction relation

Now a further development of the above model is presented. Initially only a finite energy of the uniform particles in the physical system is assumed, with the

relative value  $q = 1$ . Again, the system is correlated with the system of natural numbers.

The maximum value  $e^{1/e}$  with  $q = 1$  in Equation (1), resulted as a natural organization process with natural numbers in the direction of maximum interactions. This process can be presented formally as a result from the first two steps in the following algorithm:

$$\begin{aligned}
 y_1 &= \exp(q) \\
 y_2 &= \exp(q/y_1) \\
 y_3 &= \exp(q/y_2) \\
 &\dots\dots\dots \\
 y_i &= \exp(q/y_{i-1}).
 \end{aligned}
 \tag{4}$$

With  $q = 1$ ,  $y_1 = e$ ;  $y_2 = e^{1/e}$ . A continuation of the interaction process, towards maximum values, is now assumed in the form of the algorithm (4).

The algorithm provides an oscillating sequence. Starting with  $q = 1$ , one monotone decreasing sequence  $y_{2n-1}^I$  and one monotone increasing sequence  $y_{2n}^{II}$  results, which asymptotically reach the same limit value (Figure 1),

$$y_{\infty}^I = y_{\infty}^{II} = y_{\infty}(1) = 1.763222834351897\dots > e^{1/e} = 1.44466\dots$$




Figure 1. Graph (—) of the oscillating sequence  $y_n$  in the algorithm (4) for  $q = 1$ :  $y_{2n-1}^I$  (●) and  $y_{2n}^{II}$  (○) with the same limit value  $y_{\infty}^I = y_{\infty}^{II} = y_{\infty}(1) = 1.7632\dots$

The algorithm (4) based on the exponential function is used as backbone in this investigation, with  $y_{\infty}(1) = 1.763222834351897\dots$  as the first oscillation limit point.

The limit value  $y_{\infty}(1)$  obtained with  $q = 1$  can now be used again as starting value  $q_2 = y_{\infty}(1)$  in the algorithm (4) with the result of a new limit value  $y_{\infty}(q_2)$ , and so on. In this way a final limit value  $y_{\infty}(e) = e$  results asymptotically.

The above results are a consequence of the before mentioned organization process with natural numbers based on their basic properties. The correlation with a physical system based only on the assumption of uniform particles with finite relative energies,  $q = 1$ , leads in this way to a maximum relative interaction density of energy:

$$y_{\infty}(e) = \exp(e/y_{\infty}(e)) = e. \quad (5)$$

The value  $y_{\infty}(e) = e$  is understood as the dimensionless limit value for an interaction constant. As shown later, this limit is the coupling constant for the strong force.

With the introduction of the second fundamental quantum mechanical assumption for the physical system [1], in form of the relative wavelength  $\lambda = 2\pi$ , a special result is obtained with the algorithm (4), for  $q = w > e$ .

For an extension of the algorithm (4) into a region with  $q > e$ , the following invariant structure is postulated:

$$\exp(e \cdot A \cdot B / y_{\infty}(q)) = e \cdot A \cdot B. \quad \text{and} \quad B = (1/e \cdot A) \cdot e^{e \cdot A} \quad \text{for} \quad B = y_{\infty}(q). \quad (6)$$

The graph of the function  $B = f(A)$  from Equation (6) is shown in Figure 2.




Figure 2. Graph of  $B = (1/e \cdot A) \cdot e^{e \cdot A}$ .

In contrast to  $y_{\infty}(1)$ , with  $q=1$ , two different limit values,  $y_{\infty}^I(w)$  and  $y_{\infty}^{II}(w)$  are obtained for  $q=w$  and with the algorithm (4):

$y_{\infty}^I(w) = 23976.90006232883... > y_{\infty}^{II}(w) = 1.000420872319414...$  and their ratio  $Y_{\infty}(w) = y_{\infty}^I(w)/y_{\infty}^{II}(w) = 23966.81309411294...$

The value  $Y_{\infty}(w)$  is the second basic limit value in this investigation. Of special interest for the following relations is the ratio obtained from  $y_{\infty}(1)$  and  $Y_{\infty}(w)$  :

$$Y_{\infty}(1, w) = \exp\left(\frac{y_{\infty}(1)}{Y_{\infty}(w)}\right) = \mathbf{1.000073572055}.$$

Equation (6) is the fundamental relation for connections with natural constants.

### 3. Connection between the interaction function and fundamental natural constants

With  $B = w^2$ , the number  $A = A_0/2$  with  $A_0 = 1/136.9807$  results from Equation (6), Fig.2. This result with  $A_0$  indicates a connection between  $w$  and the fine structure constant [3]:

$$\alpha = \frac{e_q^2}{4\pi\epsilon_0} \cdot \frac{2\pi}{hc} = 1/137.035999084, \quad (7)$$

with the elementary charge,  $e_q$ , the vacuum permittivity  $\epsilon_0$ , speed of light,  $c = 2.99792458 \cdot 10^8 \text{ m}\cdot\text{s}^{-1}$ , and the Planck constant  $h = 6.62607015 \cdot 10^{-34} \text{ Js}$  [3].

$\alpha$  is a dimensionless fundamental constant of nature, derived in 1916 by Sommerfeld, representing the electromagnetic coupling constant.

The ratio

$$Y_0(A,B) = \exp(A_0/w^2) = 1.00007172196$$

from the initial values  $A_0$  and  $B = w^2$  in Equation (6) is similar to the above derived ratio  $Y_\infty(I,w) = 1.00007357206$  .

With the limits  $Y_\infty(w)$  and  $Y_\infty(I,w)$  the following number  $b$ , similar to  $w^2$ , can be defined:

$$\mathbf{b} = \left( \frac{Y_\infty(w) \cdot Y_\infty(1,w)}{(2\pi/e)} \right)^{1/2} = \pm 101.8306291... \quad (8)$$



With  $b = 101.8306291$  and Equation (6) the following iteration, for representing  $a_n$  values related to  $b_n$  values, is used:

$$\mathbf{a}_n = \left(\frac{2}{e \cdot b}\right) \cdot e^{\frac{e \cdot a_0}{2}} \cdot \frac{Y_\infty(1, w)}{Y_0(A, B)} \quad \text{and} \quad \mathbf{b}_n = b \cdot \frac{Y_n(A, B)}{Y_\infty(1, w)} \quad (9)$$

Starting with the initial values  $a_0 = A_0 = 1/136.98$  and  $b_0 = w^2$ , the iteration procedure (9) leads to limit values  $a_\infty$  and  $b_\infty$ :

$$a_1 = 1/137.035466, b_1 = 101.8304378; a_2 = 1/137.036002, b_2 = 101.830435; \\ a_3 = 1/137.0360069, b_3 = 101.8304348; a_4 = 1/137.03600696, b_4 = 101.8304348.$$

The limit values are:

$$a_\infty = 1/137.03600696 \text{ and } b_\infty = 101.8304348.$$

With the above iteration, the ratio  $a/a_\infty = 1.000000057$  results.

With  $A = a/2 = (137.035999084)/2$  and with Equation (6) a value  $b_a = 101.830429028$  results directly, with the ratio  $b_\infty/b_a = 1.000000057$ . However, in this way the limit property of  $b_\infty$  as result from the algorithm (4) is not apparent, because it is related directly to  $\alpha$  as an experimental value. Due to these small differences between experimental and calculated values,  $a_\infty$  is used in the following instead of  $\alpha$  as the electromagnetic coupling constant.

With Equation (6) and the iteration (9) the limit value  $\alpha_\infty$  results as a modelled value for the electromagnetic coupling constant. From this modelled value, instead of  $\alpha$ , in combination with the two experimental values for  $h$  and  $c$ , the value for  $e_q^2/4\pi\epsilon_0 = 2.3070775 \cdot 10^{-28}$  Jm results with Equation (7), instead of the experimental value found to be  $2.3070758 \cdot 10^{-28}$ .

The results obtained until now demonstrate the importance of the asymptotical approach introduced with the algorithm (4) for the corresponding maximal limit values of the different constants. That supports the assumption of a natural

evolution process which occurs in multiparticle systems, based on fundamental properties of natural numbers.

The model developed in this investigation [1,2] has no similarity with the well known Standard Model for interaction between fundamental natural constants [5].

The following derivations of coupling constants and masses of elementary particles emphasize the strong connection between different kinds of interactions, following from the same initial assumption. This can be seen in the following example which demonstrates the strong connection between the electromagnetic coupling constant  $\alpha_\infty$  and the limit value  $w$  of the interaction function (2). In the application of the interaction function presented in [1], with the vapor pressure of droplets as a special example, the interactions between the particles occurred in the form of strings. This behavior appeared as a characteristic property of electromagnetic interactions between the particles. The number  $B = w^2$  in Equation (6), similar with the limit value  $b_\infty$ , suggests a formal similarity between  $w$  and an electron as a multiparticle system as shown in Equation (1) for  $n \gg 1$ . With such a formal correlation, an electron may appear as a system of interacting subparticles in the form of strings along a linear arrangement, with the possibility of place exchanges between the linear arrangements of the interactions in space of the electron at a certain time. Only after an interaction between a string and another particle placed in the string direction, the exact position of the electron becomes observable [4].

Although very speculative, this discussion generates an idea about the structure of the electron, as similar with a liquid phase. The interactions between subparticles in the electron in form of strings, have as consequence the a priori unpredictable direction of the occurring interaction. This situation is fundamental

different from that in the nucleus, where the strong force is responsible for the interactions.

The electromagnetic coupling constant  $a_\infty$  results in this way as interaction of a pair of particles, corresponding to  $w^2$ , with

$$A_0 = 1/136.98, b_0 = w^2 \text{ and } Y_0(A,B) = \exp(A_0/w^2).$$

A formal similarity between  $w$  and an electron, as mentioned above, can request in conformity with Equation (1) a triple of interacting subparticles for representing a real electron, as shown later.

### The gravitational coupling constant

The limit value

$$\pm b_\infty \cdot (a_\infty \cdot e/2) \cdot e^{-a_\infty \cdot e/2} = \pm 1 \quad (10)$$

as a result from Equation (6) is obtained using algorithm (4), along with the limit value  $w > e$  in Equation (6). This result represents the two extremes obtained in the form of an extension and a contraction processes, as a consequence of interactions occurring in a physical system. As a total result, this leads to

$$+b_\infty \cdot (a_\infty \cdot e/2) \cdot e^{-a_\infty \cdot e/2} + (-b_\infty) \cdot (a_\infty \cdot e/2) \cdot e^{-a_\infty \cdot e/2} = 1 + e^{\pi i} = 0. \quad (11)$$

That means, despite the great changes in interaction energies during these processes, no additional building or destruction of parts of the total energy of the system occurs.

The small value of  $e^{-b_\infty} = 5.964912489 \cdot 10^{-45}$  indicates a connection with the dimensionless, gravitational coupling constant of nature.

The relation  $a_g = Gm_e^2 2\pi/hc$  contains the same reference group  $2\pi/hc$  like that in  $a_\infty$ .

$a_g = 1.75168850975 \cdot 10^{-45}$ , using the constant of gravitation

$$G = 6.67408(31) \cdot 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$$

and the electron mass

$$m_e = 9.1093837015 \cdot 10^{-31} \text{ kg} = 0.5109989461(31) \text{ MeV} [4]$$

represent real, experimentally measured constants, as well as the dimensionless ratio  $\alpha_g/a_\infty = 2.4004438 \cdot 10^{-43}$  [4].

If the limit value  $w$  is considered to represent a particle, than in conformity with Equation (1),  $w^3$  represents a particle (electron) having the highest interaction intensity and this leads to an interaction constant  $(w/\exp(w))^3$  for particle triplets.

By taking into account the limit value  $w_b = (\mathbf{b}_\infty)^{1/2} = 10.0911067$  from Equation (10), the following gravitational coupling constant is defined:

$$\mathbf{a}_{g,\infty} = \left[ w_b \left( \frac{w_b}{e^w} \right)^3 \right]^3 \frac{1}{e} \frac{1}{c^2} \cdot \mathbf{C}_{ag} = \mathbf{1.75169013} \cdot 10^{-45}$$

(12)

with

$$\mathbf{C}_{ag} = \frac{\exp(\mathbf{b}_\infty)}{\exp(w^2)} \left( \frac{w_b}{w} \right)^9 \left( \frac{w_b}{w} \right)^3 \cdot \left( \frac{w_b}{\ln Y_\infty(w)} \right)^{1/2} = \mathbf{1.04437119}.$$

This gives the ratio  $\mathbf{a}_g/\mathbf{a}_{g,\infty} = 1/1.00000092$ .

Neglecting the correction factor  $\mathbf{C}_{ag}$ , a value  $\mathbf{a}_{g,\infty} = \left[ w_b \left( \frac{w_b}{e^w} \right)^3 \right]^3 \frac{1}{e} \frac{1}{c^2} = 1.67727 \cdot 10^{-45}$  with the ratio  $\mathbf{a}_g/\mathbf{a}_{g,\infty} = 1.0444$  results.

The definition of  $\mathbf{a}_{g,\infty}$  in Equation (12) relates the interaction constant to the limit value  $w_b$  and the product  $\left[ w_b \left( \frac{w_b}{e^w} \right)^3 \right]^3$  to the limit  $e$  from the domain  $q \leq e$  and to  $c^2$ , in conformity with the Einstein relation,  $E = mc^2$ .

The limit values result as asymptotical limits from the algorithm (4) and this procedure has the necessity of considering corresponding correction factors as consequence, taken into account in  $\mathbf{C}_{ag}$ . The correction numbers result from the algorithm (4) with asymptotically reached limits and are related to well defined numbers. No pure empirical numbers to reach these results are used at all.

This procedure for correcting factors is applied for all following cases as well.

The dimensionless coupling constant (12) can be split into two factors,  $G_\infty$  and  $m_{e,\infty}^2$ :

$$\mathbf{G}_\infty = \left( \frac{w_b}{e^w} \right)^3 \frac{e}{3} \cdot \mathbf{C}_g = \mathbf{6.67404} \cdot \mathbf{10^{-11}} \text{ Nm}^2\text{kg}^{-2} \quad (13)$$

with

$$\mathbf{C}_g = \left( \frac{w_b}{w} \right)^3 \left( \frac{w_b}{\ln Y_\infty(w)} \right)^{1/6} = \mathbf{1.00071008}.$$

Where  $G/G_\infty = 1.000006$ . In this way the constant of gravitation results as a consequence of the extension of the algorithm into a region with  $q > e$ , with the corresponding limit  $b_\infty$ .

$$\mathbf{m}_{e,\infty}^2 = \mathbf{3(w_b)^3} \left[ \left( \frac{w_b}{e^w} \right)^3 \right]^2 \frac{\mathbf{1}}{e^2} \frac{\mathbf{1}}{c^2} \cdot \mathbf{C}_m / \left( \frac{\mathbf{2\pi}}{hc} \right) \text{ kg}^2 = \mathbf{8.29784557 \cdot 10^{-61}} \quad (14)$$

with

$$\mathbf{C}_m = \frac{\exp(b_\infty)}{\exp(w^2)} \left( \frac{w_b}{w} \right)^9 \cdot \left( \frac{w_b}{\ln Y_\infty(w)} \right)^{1/3} = \mathbf{1.0436301}.$$

With  $m_{e,\infty} = 9.1093837015 \cdot 10^{-31}$  kg,  $m_e/m_{e,\infty} = 1.000015$ .

As shown from the above relations, the mass of the electron results as a triple of the interaction value  $w$ . This behavior can be seen also from the 1/3 or 2/3 electrical charge of Quarks for the proton [5].

### The electromagnetic coupling constant

The ratio between  $a_g = Gm_e^2 2\pi/hc$  and  $a = (e_q^2/4\pi\epsilon_0)2\pi/hc$  gives the dimensionless value:

$$a_{g,\infty}/a_\infty = 2.4004438 \cdot 10^{-43} [4].$$

This relation allows a definition of the electromagnetic coupling constant with the interaction function:

$$\alpha_\infty = \frac{3w_b^2}{(ew)^3} \frac{1}{c^2} \frac{2\pi}{hc} C_{e,\epsilon} = 1/137.035954, \quad (15)$$

$$C_\alpha = \left( \frac{\exp(w^2)}{\exp(b_\infty)} \right)^{4/3} \cdot \frac{w_b^2}{w^2} = \mathbf{0.947575712913}.$$

$$\alpha/\alpha_\infty = 1/1.0000003.$$

The relation  $\frac{3b_\infty}{(ew)^3} \frac{1}{c^2}$  can be split into two factors,  $e_q^2$  and  $4\pi\epsilon_0$ :

$$e_{q,\infty}^2 = 3 \left[ \left( \frac{w_b}{ew} \right)^3 \right]^2 \frac{1}{e^2} \cdot \frac{1}{c^2} \cdot C_e = 2.566794 \cdot 10^{-38}. \quad (16)$$

$$C_e = \left( \frac{\exp(w_b)}{\exp(w^2)} \right)^{7/6} \cdot \left( \frac{w_b}{w} \right) \cdot (Y_\infty(1, w))^{-0.5} = 1.0487826$$

$$e_{q,exp} = 1.602176634 \cdot 10^{-19}, \quad e_{q,exp}/e_{q,\infty} = 1.00003$$

$$4\pi\epsilon_0 = w_b \left( \frac{w_b}{ew} \right)^3 \frac{1}{e^2} C_\epsilon = 1.1125734 \cdot 10^{-10}. \quad (17)$$

$$C_\epsilon = \frac{C_e}{C_\alpha} = 1.106806,$$

$$4\pi\epsilon_{0,exp} = 1.11265 \cdot 10^{-10}; \quad \frac{4\pi\epsilon_{0,exp}}{4\pi\epsilon_{0,\infty}} = 1.00007.$$

From all above derived relations results the importance of the initial Equation (1), with the maximum for three interacting particles.

### **The strong and weak coupling constants**

The coupling constant,  $a_s$  for the strong force and the constant for the weak coupling,  $a_w$  are not constant in reality [5]. Experimentally - the corresponding coupling constant for the strong force,  $a_s$  - as determined from the force between two protons, is greater than one [5]. But a decrease of the strong force with decreasing distances between the interacting quark particles occurs. This “running” coupling constant is a result of the phenomenon known as asymptotic freedom. The weak coupling  $a_w$ , also behaves similarly but at a slower rate [5].

With the four dimensionless coupling constants  $\alpha_g$ ,  $a_\infty$ ,  $a_w$ ,  $a_s$  and Equation (6) the following relation results:

$$\ln(\alpha_g \cdot a_w / (a_\infty \cdot a_s / 2)) \cdot (a_\infty \cdot a_s / 2) \cdot e^{-a_\infty \cdot a_s / 2} = -1 = -b_\infty \cdot (a_\infty \cdot e / 2) \cdot e^{-a_\infty \cdot e / 2} = e^{\pi \cdot i} = \cos(\pi) + i \cdot \sin(\pi)$$

(18)

The four fundamental constants of nature are connected to a fundamental mathematical relation between the transcendental numbers  $e$  and  $\pi$ . This relation is the result from the use of the limit value  $w$  of the interaction function (2) in the algorithm (4).

The structure of this relation and the limit value  $y_\infty(e) = e$  from Equation (5) suggests the value  $a_s = e$  for the strong coupling constant. With this value

$$a_w = \exp(-b_\infty) \cdot a_\infty \cdot a_s / \alpha_g \cdot 2 = 0.0337736 = 1/29.609, \quad (19)$$

results for a weak coupling constant  $a_w$ . The value  $a_w = 1/29.60895$  is similar to the published value  $1/29.5$  [5]. The value for the weak mixing Weinberg angle  $a_\infty/a_w = \sin^2(\Theta) = 0.216067$ , calculated from Equation (10) is 3 % smaller in comparison with the CODATA [3] value  $0.2229(30)$ . With this value the weak coupling constant is  $a_w = 1/30.545 = 0.03274$ . These results demonstrates the variability of the  $a_w$  and  $\alpha_s$  values. But from Equation (9) the following constant connection results:

$$a_s/a_w = 80.4856. \quad (20)$$

The gravimetric coupling constant  $a_g$  and the electromagnetic coupling constant  $\alpha_\infty$  appear only in connection with the Equation (6), beyond the limit  $q = e$ . But both domains,  $q \leq e$  and  $q \geq e$  must be considered for the strong coupling constant  $a_s$  and the weak coupling constant  $a_w$ . As a consequence this has a variability in the coupling constant values.

The weak and electromagnetic interactions can be treated as different manifestations of a single electroweak force [5]. The Weinberg angle  $\Theta$  shows the relation  $a_\infty/a_w = \sin^2(\Theta)$ . For neutrinos no electromagnetic interactions occur like those for electrons. Instead of using  $w$  as starting point in the algorithm (4), a starting value  $q_{v,0} = 2\pi/w = 0.62277$  is used.

The weak coupling appears as a consequence from the extension of Equation (5) to Equation (6), with  $q = w$  instead  $q = e$  as a starting point in the algorithm (4). The small value  $q_{v,0} = 2 \cdot \pi/w = 0.62277$  is inside of the range for the application of the algorithm (4) in conformity with Equation (5). This suggests the definition



of two limit values for  $a_w$ , based on  $e$  from Equation (5) and  $w$  from Equation (6). With  $e \cdot w = 27.425$  the maximum value  $a_{w,max} = 1/e \cdot w = 0.036463$  is defined. The limit value  $y_\infty(e) = e^1$  from Equation (5) corresponds to  $b_\infty = 101.8304348$  in Equation(6). If  $-1/y_\infty(e) = -e^{-1}$  is related to  $-b_\infty$ , then a minimum value  $a_{w,min} = -e^{-1} \cdot (-b_\infty) = 1/37.46132$  is defined.

Similar to the above two limit values for  $a_w$ , two limit values:

$$a_{s,max} = a_{w,max} \cdot 80.4856 = 2.93475 > a_s = e,$$

and

$$a_{s,min} = a_{w,min} \cdot 80.4856 = 2.1485$$

are obtained with Equation (20).

The coupling constant  $\alpha_g$  contains the mass  $m_e$  of an electron and therefore a connection between  $m_e$  and  $w$  can be established. From the approximation  $\exp(w^2)/(e^w)^2 \cdot m_e \approx e^w \cdot w$ , the following reference number  $M_e$  is defined:

$$M_e = \frac{e^w \cdot w_\infty}{Y_\infty(1,w)} \cdot \frac{1/137.036}{m_e} = \frac{1771.8992}{m_e} \quad (21)$$

The numbers  $w_\infty = \ln(y_\infty^I(w))$  and  $Y_\infty(I,w)$  take into account the asymptotical limits attained with the algorithm (4) at  $q=1$  and  $q = w$ , respectively.

With Equation (21) the masses of other elementary particles can be related to the electron mass, as shown in the following examples.

The mass of electron:

$$m_e = \frac{\frac{e^w \cdot w_\infty}{Y_\infty(1,w)} \cdot a_\infty}{M_e} = \frac{1771.8992 \cdot m_e}{1771.8992}$$

**The mass of the electron neutrino**

The weak and electromagnetic interactions can be treated as different manifestations of a single electroweak force [5]. The Weinberg angle  $\Theta$  shows the relation  $a_\infty/\alpha_w = \sin^2(\Theta)$ .

For neutrinos no electromagnetic interactions like those for electrons occur. Instead of using  $w$  as starting point in the algorithm (4), a starting value

$$q_{v,0} = 2 \cdot \pi/w = 0.62277 . \quad (22)$$

is used. With the corresponding values  $e^{q_{v,0}}$  and  $q_{v,0,\infty} = 0.412315$ , resulting with the algorithm (4), and with the weak coupling constant  $a_w$ , an expression for the neutrino mass,  $m_{v,0}$  results:

$$\mathbf{m}_{v,0} = \frac{e^{q_{v,0}} \cdot q_{v,0,\infty}}{M_e} \cdot \mathbf{a}_w . \quad (23)$$

A term corresponding to  $Y_\infty(l,w)$  in Equation (21) is not necessary in Equation (23), because  $2\pi/w < e$  and only one limit value from the algorithm (4) results:

$$q_{v,0,\infty} = \ln(y_\infty(q_{v,0}))$$

With  $a_w = 1/29.609$  from Equation (19),  $m_{v,0} = 0.05436$  eV.

With different  $a_w$  values, different values for  $m_{v,0}$  are obtained. But given the connection discussed below between the neutrino and the Z-boson [5], only the minimum value  $a_{w,min} = 1/37.46132 = 0.02669$  is selected and the mass  $m_{v,0} = 0.0432$  eV is calculated with Equation (23).

Whereas  $w$  results as the limit value from electromagnetic interactions in conformity with Equation (2),  $w_{l,e} = (1+2\pi)^{1/e} = 2.076$  can be understood as a limit value resulting from Equation (2) if no such interactions occur between the

$n$  particles. The three terms,  $w_I$ ,  $w_{I,e}$  and  $w_I - w_{I,e} = 0.6227 \approx 2\pi/w$  are then used representing three different neutrino masses with Equation (2).

Starting with  $q_{v,I} = w_{I,e}$  in algorithm (4),  $q_{v,I,\infty} = 0.87$  is obtained and with a relation similar to Equation(23),  $m_{v,I} = 0.388$  eV is obtained.

With  $q_{v,2} = w_I$  in the algorithm (4),  $q_{v,2,\infty} = 0.896$  and with a relation similar to Equation (23),  $m_{v,2} = 0.745$  eV results for  $a_{w,min}$ .

It is interesting to note an additional connection between fundamental elementary particles and macroscopic systems. For one mol of neutrino particles, Equation (23), leads to an energy of  $4168 \text{ Jmol}^{-1}$ , with the minimum value for  $\alpha_w$  and the Einstein relation  $E = m \cdot c^2$ . If the difference  $w_I - w_{I,e}$  in entropy units, is related to the entropy  $108.85 \text{ JK}^{-1}\text{mol}^{-1}$  in the gas phase, then  $4168 \cdot (w_I - w_{I,e}) / 108.85 = 23.85 \text{ Jmol}^{-1}$  results [1]. This value is 4 % smaller in comparison to the value  $24.78 \text{ Jmol}^{-1}$  obtained with the state equation  $pV = RT$ . A neutrino mass corresponding to the value from the state equation leads to  $4331 \text{ Jmol}^{-1}$  and  $0.0449$  eV instead of the minimum value  $0.0432$  eV in Equation (23). This case emphasizes the possible decrease of  $a_w$  as function of its source.

Experimentally an upper limit to the neutrino mass from a direct kinematic method by KATRIN is  $1.1$  eV [6]. From combined cosmological observations and particle physics experiments an upper bound of neutrino masses,  $\leq 0.26$  eV and an approximation-independent upper bound for the lightest neutrino mass species,  $< 0.086$  eV [7] are found.

### **The mass of the Higgs- and Z- boson**

Whereas the electron mass  $m_e$  is related to the electromagnetic coupling constant  $a_\infty$ , a mass  $m_I$  can be derived from the connection between the term  $w_I =$

$(1+2\cdot\pi)^{0.5} = 2.6987..$  in the power sequence (2) and the strong force  $a_s$ . Similar to the above use of two limit values  $a_w$ , two limit values are obtained with Equation (20):

$$a_{s,max} = a_{w,max} \cdot 80.4856 = 2.93475 > a_s = e > 2.1485 = a_{s,min}.$$

In conformity with Equation (1), a combination of three particles with relative values  $w_l$  is considered to be representative of three quarks in a particle with a mass  $m_3$ . The interaction of three terms  $w_l$  as  $w_l^3$  can be considered as resulting from the strong force. With a similar relation like Equation (21) the following expression for  $m_3$  is derived:

$$m_3 = \frac{e^{w_1^3} \cdot w_{1,\infty}^3}{M_e} \cdot 3a_s \quad (24)$$

The factor 3 takes into account the interaction of the three particles (quarks). Corresponding to  $Y_\infty(l, w)$  in Equation (21), the term  $Y_\infty(l, w_l^3) = 1$ .

With  $a_{s,max}$  instead of  $a_s$ , the value  $m_H = 125.186$  GeV results from Equation (24). It is the mass for the Higgs boson. The experimental value for this is  $m_H = 125.10 \pm 0.14$  GeV [8,9].

With  $a_{s,min}$  a mass  $m_Z = 91.647$  GeV results with Equation (24). This is similar with the mass of the Z-boson having the experimental value  $m_Z = 91.19$  GeV [5].

### **The mass of the proton**

The term  $w_l = 2.699 < e$  from the interaction function is inside of the application range for algorithm (4) in conformity with Equation (5). That means, only a strong force coupling constant exists. This can be defined as  $a_{s,p} = e^{-e}$  in comparison with  $a_s$  from Equation (18), as only one of 4 coupling constants, resulting with  $e^{-b\infty}$ . That means,  $e^{-e}$  stands for  $e^{-b\infty}$ , as a consequence of the extension into a region with  $q > e$ . With  $a_{s,p}$  and an analogous relation to Equation (21) the following mass of the proton is obtained:

$$m_p = \frac{e^{w^3} \cdot w_{1,\infty}^3}{M_e} \cdot \frac{1}{e^e} = 1836.15073 \cdot m_e \quad (25)$$

The term  $w_{1,\infty}^3 > e$  is treated as a single particle.

The experimental value for the ratio  $m_p/m_e = 1836.1527$  [3].

From this result the similarity with the atomic mass unit derived in Equation (3) is shown.

The value  $1/e^e = 0.066$  demonstrates a decrease of the strong force with decreasing distances between the interacting quark particles [5].

### A hypothetical mass $m_x$

From symmetrical reasons, in addition to the value  $w$  used in algorithm (4), a value  $q = 1/w = w_x$  is defined and introduced in the algorithm (4), producing a limit value  $w_{x,\infty} = 0.0905374$ . Whereas in conformity with Equation (21),

$m_e = \frac{e^w \cdot w_\infty}{Y_\infty(1,w) \cdot M_e}$ , there is a resulting hypothetical particle mass  $m_x$  analogous to Equation (23),

$$m_x = \frac{e^{w_x} \cdot w_{x,\infty}}{M_e} \cdot a_\infty = 3.00447 \cdot 10^{-9} \cdot m_e = 0.00153 \cdot eV. \quad (26)$$

The discussed mass of Axion, supposed as a component of Dark Matter lies within such dimensions [10].

The numerical values obtained above with the simple model based on the interaction function (2), emphasizes the role of this function as a bridge between fundamental constants and emergent properties in liquids.

#### 4. Modelling nuclear binding using the interaction function

The interaction function is not related *a priori* to particles with a special physical structure. Consequently the atomic nucleus represents another system with some similarity in comparison with liquid droplets [11].  $n$  similar particles (the nucleons) are interacting under the influence of strong forces [11,12].

In the following expression for the relative binding energy,  $E_n$ , between the  $n$  nucleons, referred to one nucleon, the term  $(w_{n,e})^3$  plays the main role, where the exponent 3 takes into account the three quarks as separate interacting sub-particles in one nucleon. In addition, a negative term proportional to  $n$  is used and the  $n-1$  interactions related to one nucleon gives:

$$E_n = (w_{n,e}^3 \cdot \pi r^2 - n \cdot \pi(2r)^2) \frac{1}{\pi r^2} \cdot \frac{n-1}{n} = (w_{n,e}^3 - 4n) \cdot \frac{n-1}{n}. \quad (27)$$

The first term  $w_{n,e}^3$  represents the mobility of one nucleon resulting from the interactions, in the form of its cross sectional area. But the movement of the nucleon must overcome the impedance produced by the relative total cross sectional area,  $n\pi(2r)^2/\pi r^2 = 4n$  of the  $n$  nucleons related to the cross section area  $\pi r^2$  of one nucleon.

The significant difference between the application of the interaction function (2) in this case and the examples with liquid properties presented in [1] lies in the involving all  $n$  nucleons in this treatment and not only the particles along a string with a linear arrangement. The place exchange of the  $n$  nucleons must overcome the resistance from the total cross sectional area of the nucleons. Instead of the

electromagnetic interactions in the liquid, strong force interactions are involved in the nuclear “droplets”.

Equation (27) shows a maximum value for  $n = 60$  and approaches zero at  $n = 238$ . That means, no stable elements exist beyond uranium (Figure 3). A correlation between the dimensionless function  $E_n$  to the binding energy  $B/A$  in MeV for the atomic mass  $n = A$ , can be established in the following manner:

The elements Be, Co and Au are selected as references, because each of them has one isotope with 100 % abundance. Their binding energies are  $B/9 = 6.46253$  MeV,  $B/59 = 8.76775$  MeV and  $B/197 = 7.91554$  MeV, respectively, obtained from the atomic masses in [13,section 1-14]. The corresponding  $E_n$  values from Equation (27) are  $E_9 = 139.039$ ,  $E_{59} = 483.025$  and  $E_{197} = 132.89$ .

With  $a1 = (B/59-B/9)/(E_{59}-E_9) = 0.00670222$  and from  $E_{59} \cdot a1 + b1 = B/59$ , with  $b1 = 5.53066$ , the relation  $B1/A$  for  $A \leq 59$  in Equation.(28) results. In the same way the relation  $B2/A$  for  $A \geq 59$  is obtained from  $B/59$ ,  $B/197$ ,  $E_{59}$  and  $E_{197}$ , respectively.

In combination with  $E_n$  from Equation (27) the curves  $B_1/A$  and  $B_2/A$  in Equation (28) represent the calculated  $B/A$  values between  $A = 9$  and  $A = 250$  (Figure 4).

$$\begin{aligned} B_1/A &= 0.00670222 \cdot E_n + 5.53066 \quad (\text{MeV}) && \text{for } n = A \leq 59 \\ B_2/A &= 0.00243466 \cdot E_n + 7.592 \quad (\text{MeV}) && \text{for } n = A \geq 59. \end{aligned} \quad (28)$$

The following Table 1 below contains 26 elements having only one single isotope, or one isotope with  $> 95$  % abundance. The experimental data are obtained from the atomic masses in [13].

Table 1.

Experimental binding energies  $B/A$  from [13, section 1-14] and  $\Delta(B/A)$  between calculated values with Equation (28) and experimental values of 26 elements.

Element	$B/A$ (MeV) a)	$\Delta(B/A)$ <sup>b)</sup>	Element	$B/A$ (MeV) a)	$\Delta(B/A)$ <sup>b)</sup>
Be	6.46253	0.0	Y	8.714	-0.022
C	7.680	-0.821	Nb	8.664	0.0
F	7.779	-0.192	Rh	8.584	0.035
Na	8.111	-0.223	I	8.445	0.021
Al	8.331	-0.207	Cs	8.410	0.014
P	8.481	-0.172	Pr	8.354	0.011
Ar	8.595	-0.013	Tb	8.189	0.039
Sc	8.619	0.053	Ho	8.147	0.033
V	8.742	0.0	Tm	8.114	0.033
Mn	8.765	0.0	Au	7.91554	0.0
Fe	8.790	-0.028	Bi	7.848	-0.035
Co	8.76775	0.0	Th	7.615	0.0
As	8.701	0.044	U	7.570	-0.01

a) [13]; b)  $\Delta(B/A) = (B/A)_{calc} - (B/A)_{exp}$  (MeV)






Figure 3. The interaction function  $E_n$  from Equation (27) versus the number  $n$  of nucleons.




Figure 4. Binding energy/nucleon (MeV) as function of the nucleon number  $A$ . (●) experimental values; (—) calculated values with Equation (28).

It is widely believed that  $^{56}\text{Fe}$  is the most tightly bound atomic nuclide. The higher binding energy of  $^{62}\text{Ni}$  in comparison with  $^{56}\text{Fe}$  has been discussed recently [12]. The scatter of the binding energies for 10 isotopes of Fe, Co and Ni, obtained from their atomic masses published in [13, section 1-14], is compared with the corresponding values calculated with Equation (27) and Equation (28) and shown in Table 2 and Fig.5.

The calculated values with Equation (27) have a maximum for  $A = 60$ . Even if  $^{62}\text{Ni}$ , is used as reference instead of Co, the calculated  $B/A$  values with Equation (28) give a maximum value for  $A = 60$ , as shown in Fig.5.




Figure 5. Binding energies  $B/A$  (●) obtained from atomic masses [13] for the 10 isotopes from Table 10 [13] and calculated (—) with Equation (27). Calculated (- - -) values with  $B/62 = 8.794$  for  $^{62}\text{Ni}$  as reference instead of Co and Equation (27).

Table 2.

Experimental atomic masses  $M$  in mass units  $u$  and binding energies  $B/A$  for 10 isotopes from [13, section 1-14].

Isotope	$M$ (u)	$B/A$ (MeV)	Isotope	$M$ (u)	$B/A$ (MeV)
$^{54}\text{Fe}$	53.93961	8.736	$^{58}\text{Ni}$	57.9353	8.732
$^{56}\text{Fe}$	55.9349	8.790	$^{60}\text{Ni}$	59.9308	8.780
$^{57}\text{Fe}$	56.9354	8.770	$^{61}\text{Ni}$	60.9311	8.765
$^{58}\text{Fe}$	57.9333	8.792	$^{62}\text{Ni}$	61.9283	8.794
$^{59}\text{Co}$	58.9332	8.768	$^{64}\text{Ni}$	63.9280	8.777

An essential difference between Equation (27) for nuclear binding and the vapor pressures in liquid droplets [1] is the inclusion of all  $n$  nucleons in the interaction in Equation (27) and not only  $i = n^{1/3}$  molecules, as in liquid droplets. This

emphasizes the different interaction forces in the two systems, one as strong and the other as electromagnetic, respectively.

## **5. Conclusions**

In addition to interactions between a few elementary particles, powerful principles of organization in the whole physical system must be considered. The properties of natural numbers play a crucial role in this case. The fundamental properties of natural numbers reflect the asymptotical limits reached at equilibrium and these reflect the properties following for the physical systems. This situation is the fundamental position as a bridge between a few particles and macroscopic systems with a very large number of interacting particles. In this connection it is necessary to emphasize the practically unlimited number of small differences between specific systems, because one particle with a different structure can produce different properties in the whole system, but without fundamental changes in the organization.

This bridge, specific for liquid systems, has not been considered until about 1900, when the finite value of  $h$  and a little later the velocity of light  $c$  were introduced. With these two empirical constants of nature, most fundamental interactions between particles in physical systems were predictable with models. Neglecting limited physical values for energy of elementary particles, the Schrödinger Equation is of no sense. The algorithm (4) developed in this manuscript is exact the result of this bridge and demonstrates the impossibility of correct interaction modelling based only of pair interactions, as correct starting points for macroscopic parts of matter, as demonstrated in the fundamental laws of Newton and Einstein.

A physical system of uniform particles with finite energies is correlated with the system of natural numbers. With an algorithm based on the properties of natural numbers, a maximal limit value,  $y(e) = e$  results asymptotically for a relative interacting energy density in the system.

In order to continue the application of the algorithm for initial values  $q > e$ , an invariant relation (6), with  $B = (1/e \cdot A) \cdot e^{e \cdot A}$  for  $B = y_\infty(q)$  has been postulated. In this way a strong correlation between the electromagnetic coupling constant  $\alpha$  and the limit value  $w$  from the interaction function results. From the limit value of the algorithm with  $w$  as an initial value, a limit value  $\pm b_\infty$  is obtained. The number  $\exp(-b_\infty)$  contains the four fundamental coupling constants. The values of the strong and weak coupling constants lie between two extremes, which allow the determination of the masses of the Higgs and Z bosons. In the same way the mass of the neutrino, as related to the Z-boson, is derived.

The limit value  $w$  acts as bridge between the emergent properties of macroscopic systems and the fundamental constants of nature.

Of fundamental importance of the results obtained with this model is the difference of nature between the electromagnetic interaction contained in the coupling constant  $a_\infty$  and the strong coupling constant  $e$ . Related to  $a_\infty$  is the electron and the many interactions between atoms and molecules treated in [1]. Related to  $e$  is the proton and the interactions between nucleons.

A special relation has been derived for the nuclear binding energy of the elements. A maximum value results for  $n = 60$  nucleons and a minimum for  $n = 238$ . Beyond the element U no stable elements can be formed.

Different forces interact in liquid droplets and between nucleons. In liquid droplets the interaction of electromagnetic forces occurs in form of strings, whereas strong forces between nucleons produce a place exchange between all particles in the system.

For all kinds of particles resulted from the different forces, the gravitational interactions between them, based on Equation (12) must be considered.

As a general conclusion, the strong interactions between the different kinds of particles and macroscopic liquid systems, based on the interaction function as bridge, should be emphasized.

## References

- [1] O.G. Piringer, A power sequence interaction function for liquid phase particles; *Fluids* 6 (2021), 354. the strong force is responsible for the interaction between the three particles.
- [2] O.G. Piringer, A predictive power sequence equation for vapour pressures of pure organic fluids and partial pressures in multicomponent systems in equilibrium; *Fluid Phase Equilib.* 506, (2020), 112409.
- [3] CODATA Internationally recommended 2018 values of the fundamental physical constants.
- [4] R.P. Feynman, *The Feynman Lectures on Physics. Vol.1.Mainly Mechanics, Radiation, and Heat.* 1963, California Institute of Technology.
- [5] D.Griffiths, Introduction to elementary particles, Second, revised edition, Wiley-VCH, 2008.
- [6] M. Aker et al., Improved upper limit on the neutrino mass from a direct kinematic method by KATRIN, *Physical Review Letters*, 123, 221802, 2019.
- [7] A. Loureiro et.al., Upper bound of neutrino masses from combined cosmological observations and particle physics experiments, *Physical Review Letters*, 123, 081301, 2019.
- [8] ATLAS collaboration, Observation of a new particle in the search for the standard model Higgs boson with the ATLAS detector at the LHC, *Physics Letters B.* 716, Nr.1, 2012, 1-29.
- [9] CMS collaboration, Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC, *Physics Letters B.* 716, Nr.1, 2012, 30-61.
- [10] ADMX Collaboration and G.C. Hilton, Search for Invisible Axion Dark Matter with the Axion Dark Matter Experiment, *Physical Review Letters* 120, 151301, 2018.

[11] C.F. von Weizsäcker, Zur Theorie der Kernmassen, *Z. Physik*, 96, 431-458, (1935).

[12] M.P. Fewell, The atomic nuclide with the highest mean binding energy, *Am.J.Phys.* 63 (7), 653-658 (1995).

[13] D.R. Lide, (Ed.), *Handbook of Chemistry and Physics*, CRC Press, Taylor & Francis Group 89<sup>th</sup> ed. 2008/2009, Section 6 and 3.