

The Origin of the Two New Baryon Resonances with a Mass of 5935 and 5955 MeV Observed in LHCb Experiment

Sylwester Kornowski

Abstract: Here, within the Scale-Symmetric Everlasting Theory, the origin of the two new baryon resonances with a mass of 5935 and 5955 MeV observed in LHCb experiment is presented.

1. Introduction and motivation

The Scale-Symmetric Everlasting Theory (S-SET) [1] leads to the atom-like structure of baryons. There is the core and the quantum states of pions outside it [1]. In the high-energy collisions the states of pions are destroyed and there dominate the phenomena characteristic for the core of baryons. The mass distance between the charged and neutral core of baryons is $\Delta m^+ = 2.66$ MeV [1]. The cores of baryons can be neutral or charged positively. They produce pairs in such a way that the component charged positively is real whereas the charged negatively is virtual – it leads to the positron fraction in the cosmic rays [2]. On the surface of the core can appear the pions [1]. The mass distance between the charged and neutral pion is $\Delta M^+ = 4.60$ MeV [1]. The mean energy of such condensates is $E_L^+ = (\Delta m^+ + \Delta M^+) / 2 = 3.63$ MeV. The condensates with energy E_L^+ can transform into circular loops [1]. The energy of a wave with the length equal to the radius of the loop is $E_U^+ = 2 \pi E_L = 22.81$ MeV.

In the outer shell of the baryonic plasma can be produced the charged bottom baryons with a mass of $m(\Xi_b^-) = 5791.1 \pm 2.2$ MeV [3].

The calculated mass of the charged pion is $m(\pi^-) = 139.57041$ MeV [1].

Now we can calculate the masses and define charges of the two new baryon resonances

$$m(\Xi_b'^-) = m(\Xi_b^-) + m(\pi^-) + E_L^+ = 5934.30 \pm 2.2 \text{ MeV}. \quad (1)$$

$$m(\Xi_b^{*-}) = m(\Xi_b^-) + m(\pi^-) + E_U^+ = 5953.48 \pm 2.2 \text{ MeV}. \quad (2)$$

The obtained theoretical results are consistent with experimental data obtained in the LHCb experiment presented here [4]. Notice that the natural width is measured to be $\Gamma(\Xi_b^{*-}) = 1.65 \pm 0.41$ MeV [4].

References

- [1] Sylwester Kornowski (14 March 2014). “The Everlasting Theory and Special Number Theory”.
<http://www.rxiv.org/abs/1203.0021> [v3].
- [2] Sylwester Kornowski (3 November 2014). “The Positron Fraction in Primary Cosmic Rays and New Cosmological Problems”.
<http://www.rxiv.org/abs/1411.0018> .
- [3] J. Beringer *et al.* (Particle Data Group).
PR **D86**, 010001 (2012) and 2013 partial update for the 2014 edition.
<http://pdg.lbl.gov> .
- [4] LHCb collaboration (submitted on 18 November 2014). “Observation of two new Ξ_b^- baryon resonances”.
arXiv:1411.4849 [hep-ex].