

How to project onto extended second order cones

O. P. Ferreira* S. Z. Németh †

November 4, 2016

Abstract

The extended second order cones were introduced by G. Zhang and S. Z. Németh for solving mixed complementarity problems and variational inequalities on cylinders. R. Sznajder determined the automorphism groups and the Lyapunov or bilinearity ranks of these cones. G. Zhang and S. Z. Németh found both necessary conditions and sufficient conditions for a linear operator to be a positive operator of an extended second order cone. This note will give formulas for projecting onto the extended second order cones. In the most general case the formula will depend on a piecewise linear equation for one real variable which will be solved by using numerical methods.

Keywords: Semi-smooth equation, extended second order cone, metric projection, piecewise linear Newton method.

2010 AMS Subject Classification: 47H99, 90C33

1 Introduction

The Lorentz cone is an important object in theoretical physics. Recently it has been rebranded as second order cone and used for various application in optimization. Some robust optimization, plant location and investment portfolio management problems were formulated as as a second order cone program [1]. Another good survey paper with a wide range of applications of second order cone programming is [2]. More recent connections of second order cone programming and second order cone complementarity problem with physics, mechanics, economics, game theory, robotics, optimization and neural networks were considered in [3–11]. The importance of the second order cone in optimization is nowadays notorious not only in theoretical physics, but in optimization as well.

The projection mapping plays a crucial role in solving optimization and complementarity problems. Usually there is no closed-form expressions for projecting onto a cone. A nice property of the second order is that it admits an explicit representation of the projection mapping onto it see [12, Proposition 3.3]. The original motivation for extending the second order cone was inspired by using iterative methods for solving complementarity problems and variational inequalities [13, 14]. These iterative methods are based on the property that the projection onto the closed convex set defining the problem is isotone with respect to the order defined out by a cone. Usually this is a very restrictive condition. However, cylinders and in particular cylinders with cone base admit isotone projections onto them with respect to the extended

*IME/UFG, Avenida Esperana, s/n, Campus Samambaia, Goiânia, GO, 74690-900, Brazil (e-mail:orizon@ufg.br). The author was supported in part by FAPEG, CNPq Grant 305158/2014-7 and PRONEX–Optimization(FAPERJ/CNPq).

†School of Mathematics, University of Birmingham, Watson Building, Edgbaston, Birmingham B15 2TT, United Kingdom (e-mail:s.nemeth@bham.ac.uk).

second order cones. Therefore, variational inequalities on cylinders and mixed complementarity problems can be solved by using such iterative techniques based on monotone convergence [13, 15].

Later it turned out that many of these cones could be even more useful because they have nice numerical properties, that is, the bilinearity rank (or Lyapunov rank) [16–20] of these cones is higher than the dimension of the underlying space. More specifically, for $p > 1$ this is true whenever $q^2 - 3q + 2 > 2p$ [21], where p, q are from the definition of the extended second order cone (see Definition 1). Such cones are “numerically good” cones when solving complementarity problems defined on them. The extended second order cones are also irreducible cones [21]. But to be really usable from optimization point of view we need easy ways of projecting onto them. This paper will show that projecting onto an extended second order cone it is “almost possible” by using closed-form expression. We will present a set of formulas for projecting onto an extended second order cone which is only subject to solving a piecewise linear equation with one real variable. The method of finding these expressions is based on the special form of the complementarity set of the extended second order cone and Moreau’s decomposition theorem [22] for projecting onto cones.

The structure of the paper is as follows:

In Section 2 we fix the notation and the terminology used throughout the paper. In Section 3 we present the formulas for projecting onto the extended second order cone. In Section 4 we solve the piecewise linear equation involved in these formulas by using the semi-smooth Newton’s method, a method based on Picard’s iteration and the bisection method. Finally, we make some remarks in the last section.

2 Preliminaries

Let ℓ, m, p, q be positive integers such that $m = p + q$. We identify the the vectors of \mathbb{R}^ℓ with $\ell \times 1$ matrices with real entries. The scalar product in \mathbb{R}^ℓ is defined by the mapping

$$\mathbb{R}^\ell \times \mathbb{R}^\ell \ni (x, y) \mapsto \langle x, y \rangle := x^\top y \in \mathbb{R}$$

and the corresponding norm by

$$\mathbb{R}^\ell \ni x \mapsto \|x\| := \sqrt{\langle x, x \rangle} \in \mathbb{R}.$$

For $x, y \in \mathbb{R}^\ell$ denote $x \perp y$ if $\langle x, y \rangle = 0$. We identify the elements of $\mathbb{R}^p \times \mathbb{R}^q$ with the elements of \mathbb{R}^m through the correspondence

$$\mathbb{R}^p \times \mathbb{R}^q \ni (x, y) \mapsto (x^\top, y^\top)^\top.$$

Through this identification the scalar product in $\mathbb{R}^p \times \mathbb{R}^q$ is defined by

$$\langle (x, y), (u, v) \rangle := \langle (x^\top, y^\top)^\top, (u^\top, v^\top)^\top \rangle = \langle x, u \rangle + \langle y, v \rangle.$$

A closed set $K \subset \mathbb{R}^\ell$ with nonempty interior is called a *proper cone* if $K + K \subset K$, $K \cap (-K) = \{0\}$ and $\lambda K \subset K$, for any λ positive real number. The *dual cone* of a proper cone $K \subset \mathbb{R}^\ell$ is a proper cone defined by

$$K^* = \{x \in \mathbb{R}^\ell : \langle x, y \rangle \geq 0, \forall y \in K\}.$$

A proper cone $K \subset \mathbb{R}^\ell$ is called *subdual* if $K \subset K^*$, *superdual* if $K^* \subset K$ and *self-dual* if $K^* = K$. If $K, D \subset \mathbb{R}^\ell$ are proper cones such that $D = K^*$, then $D^* = K$ and the cones K, D are called *mutually dual*.

For a proper cone $K \in \mathbb{R}^\ell$ denote

$$C(K) = \{(x, y) \in K \times K^* : x \perp y\}$$

the *complementarity set* of K .

Let $C \in \mathbb{R}^\ell$ be a closed convex set. The projection mapping $P_C : \mathbb{R}^\ell \rightarrow \mathbb{R}^\ell$ onto C is the mapping defined by

$$P_C(x) = \operatorname{argmin}\{\|x - y\| : y \in C\}.$$

Recall the following decomposition Theorem of Moreau [22] (stated here for proper cones only):

Moreau's Decomposition Theorem. *Let $K \subset \mathbb{R}^\ell$ be a proper cone, K^* its dual cone and $z \in \mathbb{R}^\ell$. Then, the following two statements are equivalent:*

- (i) $z = x - y$ and $(x, y) \in C(K)$,
- (ii) $x = P_K(z)$ and $y = P_{K^*}(-z)$.

It can be easily seen that Moreau's decomposition theorem implies

$$z = P_K(z) - P_{K^*}(-z),$$

with $P_K(z) \perp P_{K^*}(-z)$.

For a vector $z \in \mathbb{R}^\ell$ denote z_1, \dots, z_ℓ its components, that is, $z = (z_1, \dots, z_\ell)^\top$. Let \geq denote the component-wise order in \mathbb{R}^ℓ , that is, the order defined by $\mathbb{R}^\ell \ni x \geq y \in \mathbb{R}^\ell$ if and only if $x_i \geq y_i$ for $i = 1, \dots, \ell$. Denote by 0 the vector in \mathbb{R}^ℓ with all components zero, by e the vector in \mathbb{R}^ℓ with all components one and by $\mathbb{R}_+^\ell = \{x \in \mathbb{R}^\ell \mid x \geq 0\}$ the nonnegative orthant. The proper cone \mathbb{R}_+^ℓ is self-dual. For a real number $\alpha \in \mathbb{R}$ denote $\alpha^+ = \max(\alpha, 0)$ and $\alpha^- = \max(-\alpha, 0)$. For a vector $z \in \mathbb{R}^\ell$ denote $z^+ = (z_1^+, \dots, z_\ell^+)$, $z^- = (z_1^-, \dots, z_\ell^-)$ and $|z| = (|z_1|, \dots, |z_\ell|)$. It is known that $z^+ = P_{\mathbb{R}^\ell}(z)$ and $z^- = P_{\mathbb{R}^\ell}(-z)$.

Recall the following definition of a pair of mutually dual extended second order cones L, M :

Definition 1.

$$L = \{(x, u) \in \mathbb{R}^p \times \mathbb{R}^q : x \geq \|u\|e\},$$

$$M = \{(x, u) \in \mathbb{R}^p \times \mathbb{R}^q : \langle x, e \rangle \geq \|u\|, x \geq 0\},$$

where \geq denotes the component-wise order and e is the p dimensional vector with all components 1.

It is known that both L and M are proper cones, L is subdual M is superdual and if $p = 1$, then both cones reduce to the second order cone. The cones L and M are polyhedral if and only if $q = 1$. If we allow $q = 0$ as well, then the cones L and M reduce to the nonnegative orthant. More properties of the extended second order cones can be found in [13, 15, 21].

3 Projection formulas for extended second order cones

Let p, q be positive integers.

Proposition 1. *Let $x, y \in \mathbb{R}^p$ and $u, v \in \mathbb{R}^q \setminus \{0\}$. We have that $(x, u, y, v) := ((x, u), (y, v)) \in C(L)$ if and only if there exists a $\lambda > 0$ such that $v = -\lambda u$, $\langle y, e \rangle = \|v\|$ and $(x - \|u\|e, y) \in C(\mathbb{R}_+^p)$.*

Proof. Suppose first that there exists $\lambda > 0$ such that $v = -\lambda u$, $\langle y, e \rangle = \|v\|$ and $(x - \|u\|e, y) \in C(\mathbb{R}_+^p)$. Hence, $(x, u) \in L$ and $(y, v) \in M$. Moreover,

$$\langle (x, u), (y, v) \rangle = \langle x, y \rangle + \langle u, v \rangle = \|u\|\langle e, y \rangle - \lambda\|u\|^2 = \|u\|\|v\| - \lambda\|u\|^2 = 0.$$

Thus, $(x, u, y, v) \in C(L)$.

Conversely, suppose that $(x, u, y, v) \in C(L)$. Then, $(x, u) \in L$, $(y, v) \in M$ and

$$0 = \langle (x, u), (y, v) \rangle = \langle x, y \rangle + \langle u, v \rangle \geq \langle \|u\|e, y \rangle + \langle u, v \rangle \geq \|u\|\|v\| + \langle u, v \rangle \geq 0.$$

Hence, there exists $\lambda > 0$ such that $v = -\lambda u$, $\langle e, y \rangle = \|v\|$ and $\langle x - \|u\|e, y \rangle = 0$. It follows that $(x - \|u\|e, y) \in C(\mathbb{R}_+^p)$. \square

Theorem 1. *Let $(z, w) \in \mathbb{R}^p \times \mathbb{R}^q$. Then, we have*

1. *If $z^+ \geq \|w\|e$, then $P_L(z, w) = (z^+, w)$ and $P_M(-z, -w) = (z^-, 0)$.*
2. *If $\langle z^-, e \rangle \geq \|w\|$, then $P_L(z, w) = (z^+, 0)$ and $P_M(-z, -w) = (z^-, -w)$.*
3. *If $z^+ \not\geq \|w\|e$ and $\langle z^-, e \rangle < \|w\|$, then the piecewise linear equation*

$$\lambda\|w\| = \langle e, [(\lambda + 1)z - \|w\|e]^- \rangle. \quad (1)$$

has a unique positive solution $\lambda > 0$,

$$P_L(z, w) = \left(\left[z - \frac{1}{\lambda + 1}\|w\|e \right]^+ + \frac{1}{\lambda + 1}\|w\|e, \frac{1}{\lambda + 1}w \right) \quad (2)$$

and

$$P_M(-z, -w) = \left(\left[z - \frac{1}{\lambda + 1}\|w\|e \right]^-, -\frac{\lambda}{\lambda + 1}w \right) \quad (3)$$

Proof. We will use Moreau's decomposition theorem for L for proving all three items. In this case this theorem states that, $P_L(z, w) = (x, u)$ and $P_M(-z, -w) = (y, v)$ if and only if $(z, w) = (x, u) - (y, v)$ and $(x, u, y, v) \in C(L)$.

1. This is exactly the case when $v = 0$.

Indeed, $v = 0$ implies $P_L(z, w) = (x, u)$ and $P_M(-z, -w) = (y, 0)$. Hence, $z = x - y$, $w = u$, $x \geq \|u\|e$, $y \geq 0$ and $\langle x, y \rangle = 0$. By using Moreau's decomposition theorem for \mathbb{R}_+^p , we have that $z = x - y$, $x \geq 0$, $y \geq 0$ and $\langle x, y \rangle = 0$ implies $x = z^+$ and $y = z^-$. Since, $w = u$ and $x \geq \|u\|e$, we get $z^+ \geq \|w\|e$.

Conversely, suppose that $z^+ \geq \|w\|e$. Then, it is easy to check that $(z^+, w, z^-, 0) \in C(L)$. Then, by Moreau's decomposition Theorem for L , we get $P_L(z, w) = (z^+, w)$ and $P_M(-z, -w) = (z^-, 0)$. Thus, $v = 0$.

2. This is exactly the case when $u = 0$.

Indeed, $u = 0$ implies $P_L(z, w) = (x, 0)$ and $P_M(-z, -w) = (y, v)$. Hence, $z = x - y$, $w = -v$, $x \geq 0$, $\langle y, e \rangle \geq \|v\|$, $y \geq 0$ and $\langle x, y \rangle = 0$. By using Moreau's decomposition theorem for \mathbb{R}_+^p , we have that $z = x - y$, $x \geq 0$, $y \geq 0$ and $\langle x, y \rangle = 0$ implies $x = z^+$ and $y = z^-$. Since $w = -v$ and $\langle y, e \rangle \geq \|v\|$, we get $\langle z^-, e \rangle \geq \|w\|$.

Conversely, suppose that $\langle z^-, e \rangle \geq \|w\|$. Then, it is easy to check that $(z^+, 0, z^-, -w) \in C(L)$. Then, by Moreau's decomposition Theorem for L , we get $P_L(z, w) = (z^+, 0)$ and $P_M(-z, -w) = (z^-, -w)$. Thus, $u = 0$.

3. This is exactly the case when neither the condition of item 1 nor the condition of item 2 holds, or equivalently when $u \neq 0$ and $v \neq 0$.

From Proposition 1 it follows that $(z, w) = (x, u) - (y, v)$ and $(x, u, y, v) \in C(L)$ is equivalent to $z = x - y$, $w = u - v$ and the existence of a $\lambda > 0$ such that $v = -\lambda u$, $\langle y, e \rangle = \|v\|$ and $(x - \|u\|e, y) \in C(\mathbb{R}_+^p)$. On the other hand, by Moreau's decomposition theorem for \mathbb{R}_+^p , $(x - \|u\|e, y) \in C(\mathbb{R}_+^p)$ is equivalent to $x - \|u\|e = [x - \|u\|e - y]^+$ and $y = [x - \|u\|e - y]^-$. Hence,

$$P_L(z, w) = \left(x, \frac{1}{\lambda + 1} w \right) \quad (4)$$

and

$$P_M(-z, -w) = \left(y, -\frac{\lambda}{\lambda + 1} w \right) \quad (5)$$

if and only if $z = x - y$ and $\lambda > 0$ is such that

$$\langle y, e \rangle = \frac{\lambda}{\lambda + 1} \|w\|, \quad (6)$$

$$x = \left[z - \frac{1}{\lambda + 1} \|w\|e \right]^+ + \frac{1}{1 + \lambda} \|w\|e \quad (7)$$

and

$$y = \left[z - \frac{1}{\lambda + 1} \|w\|e \right]^-. \quad (8)$$

From equations (4) and (7) follows equation (2) and from equations (5) and (8) follows equation (3), where $\lambda > 0$ is given by equation (1), which is a combination of equations (6) and (8). The uniqueness of $\lambda > 0$ which satisfies (1) follows from the uniqueness of $P_L(z, w)$ and $P_M(z, w)$. □

Alternative Proof.

For proving items 1 and 2 it is enough to take the third paragraphs from the original proofs of these items (i.e., the paragraphs which start with the word ‘‘Conversely’’).

3. Denote the right hand sides of equations (2) and (3) by (x, u) and (y, v) , respectively. It is easy to see that $(z, w) = (x, u) - (y, v)$. On the other hand $y = [z - \|u\|e]^-$ and $x - \|u\|e = [z - \|u\|e]^+$. Hence, $\langle x - \|u\|e, y \rangle = 0$. Thus, it follows that $(x - \|u\|e, y) \in C(\mathbb{R}_+^p)$. If equation (1) has a positive solution, then by using some algebra, it can be seen that equation (1) implies that $0 = \langle (x, u), (y, v) \rangle$. It follows that

$$0 = \langle x, y \rangle + \langle u, v \rangle = \langle [z - \|u\|e]^+ + \|u\|e, [z - \|u\|e]^- \rangle + \langle u, v \rangle = \|u\| \langle e, y \rangle + \langle u, v \rangle,$$

which, by using $v = -\lambda u$, implies that $\langle e, y \rangle = \|v\|$. Under the given conditions of item 3, we also have $u \neq 0$ and $v \neq 0$. Hence, Proposition 1 implies $(x, u, y, v) \in C(L)$. Therefore equations (2) and (3) follow from Moreau's decomposition theorem. It remains to show that (1) has a positive solution. Let

$$\psi(\lambda) = \lambda \|w\| - \langle e, [(\lambda + 1)z - \|w\|e]^- \rangle.$$

In this case $z \not\geq \|w\|e$, otherwise it would follow that $z^+ = z \geq \|w\|e$, which contradicts our assumptions. Hence, we get $\psi(0) < 0$. If $\lambda > 0$ is sufficiently large, then we get

$$\psi(\lambda) \geq \lambda \|w\| - \langle e, (\lambda + 1)z^- + \|w\|e \rangle = (\|w\| - \langle z^-, e \rangle)\lambda - \langle e, z^- + \|w\|e \rangle > 0.$$

Since ψ is continuous, there is a $\lambda > 0$ such that $\psi(\lambda) = 0$ and hence equation (1) has a positive solution. The uniqueness of the projection shows that this positive solution is unique.

□

The next remark will recover the well known formulas for projecting onto the second order cone (see for example [12, Proposition 3.3]).

Remark 1. Let $(z, w) \in \mathbb{R} \times \mathbb{R}^q$ and L be the second order cone. Then, Theorem 1 implies that

$$P_L(z, w) = \begin{cases} \frac{1}{2} \left([z - \|w\|]^+ + [z + \|w\|]^+, [[z + \|w\|]^+ - [z - \|w\|]^+] \frac{w}{\|w\|} \right), & w \neq 0, \\ (z^+, 0), & w = 0. \end{cases} \quad (9)$$

For $p = 1$, the conditions in item 3 in Theorem (1) hold if and only if $0 \leq |z| < \|w\|$ and equation (1) becomes $\lambda\|w\| = [(\lambda + 1)z - \|w\|]^-$, which obviously can have only nonnegative solutions, because the right hand side of the equation is nonnegative. Moreover, $\lambda = 0$ cannot be a solution neither because that would imply $|z| - \|w\| \geq z - \|w\| > 0$. Hence, the conditions in item 3 hold if and only if (1) becomes $\lambda\|w\| = (\|w\| - \lambda + 1)z$. This latter equation has the unique positive solution

$$\lambda = \frac{\|w\| - z}{\|w\| + z}. \quad (10)$$

By using equation (2) and (10), it is just a matter of algebraic manipulations to check that (9) holds for this case. The cases described by items 1 and 2 can be similarly checked.

4 Numerical methods for projecting

In this section we will present three well known numerical methods to find the unique zero of the piecewise linear equation (1), in order to project onto the extended second order cones. We note that $(z, w) \in \mathbb{R}^p \times \mathbb{R}^q$ satisfies the two conditions in item 3 of Theorem 1 if and only if

$$\exists i_0 \in \{1, \dots, p\}; \quad 0 \leq z_{i_0}^+ < \|w\|, \quad 0 \leq \sum_{i=1}^p z_i^- < \|w\|. \quad (11)$$

Throughout this section we will assume that $(z, w) \in \mathbb{R}^p \times \mathbb{R}^q$ satisfies (11).

4.1 Semi-smooth Newton method

In order to study (1), we consider the piecewise linear function $\psi : (0, +\infty) \rightarrow \mathbb{R}$ defined by

$$\psi(\lambda) := \lambda\|w\| - \langle e, [(\lambda + 1)z - \|w\|e]^- \rangle. \quad (12)$$

The function ψ is piecewise linear and its unique zero is the solution of (1). The *semi-smooth Newton method* for finding the zero of ψ , with starting point $\lambda_0 \in (0, +\infty)$, it is formally defined by

$$\psi(\lambda_k) + s_k (\lambda_{k+1} - \lambda_k) = 0, \quad s_k \in \partial\psi(\lambda_k), \quad k = 0, 1, \dots, \quad (13)$$

where s_k is any subgradient in $\partial\psi(\lambda_k)$ the Clarke generalized subdifferential of ψ at λ_k . By letting

$$N(\lambda) := \text{diag} \left(-\text{sgn} \left([(\lambda + 1)z - \|w\|e]^- \right) \right), \quad \lambda \in (0, +\infty), \quad (14)$$

where $\text{sgn}(x) := (\text{sgn}(x_1), \dots, \text{sgn}(x_p))$ for each $x \in \mathbb{R}^p$, we can see $\|w\| - \langle e, N(\lambda)z \rangle \in \partial\psi(\lambda)$. Since $N(\lambda)[(\lambda + 1)z - \|w\|e] = [(\lambda + 1)z - \|w\|e]^-$, by setting $s_k = \|w\| - \langle e, N_k z \rangle$ and

$$N_k := N(\lambda_k), \quad (15)$$

equation (13) implies $\lambda_k \|w\| - \langle e, N_k [(\lambda_k + 1)z - \|w\|e] \rangle + [\|w\| - \langle e, N_k z \rangle] [\lambda_{k+1} - \lambda_k] = 0$. After some algebraic manipulations, the last equality becomes

$$[\|w\| - \langle e, N_k z \rangle] \lambda_{k+1} = \langle e, N_k [z - \|w\|e] \rangle, \quad k = 0, 1, \dots, \quad (16)$$

which formally defines the *semi-smooth Newton sequence* $\{\lambda_k\}$ for solving (1).

Remark 2. For $p = 1$, the conditions in (11) hold if and only if $0 \leq |z| < \|w\|$. Thus, if $z \leq 0$, then $N_k \equiv -1$ and $\lambda_{k+1} = [\|w\| - z] / [\|w\| + z]$ for all $k = 0, 1, \dots$. Now, if $z > 0$ then letting $0 < \lambda_0 < [\|w\| - z] / z$, we have $N_0 \equiv -1$ and $\lambda_1 = [\|w\| - z] / [\|w\| + z]$. Therefore, from Remark 1, we conclude that semi-smooth Newton (16) solve equation (1) for $p = 1$ with only one iteration.

Proposition 2. Assume that $2 \langle e, |z| \rangle < \|w\|$. Then, for any $\lambda_0 > 0$, the sequence $\{\lambda_k\}$ defined in (16) is well defined and converges linearly to the unique solution λ_* of (1) as follows

$$|\lambda_* - \lambda_{k+1}| \leq \frac{\langle e, |z| \rangle}{\|w\| - \langle e, N_k z \rangle} |\lambda_* - \lambda_k|, \quad k = 0, 1, \dots \quad (17)$$

Proof. Definition (14) implies that the entries of N_k are equal to 0 or -1 , for all $k = 0, 1, \dots$. Thus, from (11) we have $\|w\| - \langle e, N_k z \rangle > 0$ for all $k = 0, 1, \dots$, which implies that the sequence $\{\lambda_k\}$ is well defined. Since we are under conditions in (11), then item 3 in Theorem 1 implies that (1) has a unique solution $\lambda_* \in (0, +\infty)$. Thus, from the definition of ψ , we conclude that $0 = \psi(\lambda_*)$. Hence, by using (15), (16) and the definition of ψ , after some algebra we obtain that

$$\begin{aligned} \lambda_* - \lambda_{k+1} &= [\|w\| - \langle e, N_k z \rangle]^{-1} \left[-\lambda_* \|w\| + \langle e, [(\lambda_* + 1)z - \|w\|e]^- \rangle + \right. \\ &\quad \left. \lambda_k \|w\| - \langle e, [(\lambda_k + 1)z - \|w\|e]^- \rangle + [\|w\| - \langle e, N_k z \rangle] [\lambda_* - \lambda_k] \right], \quad k = 0, 1, \dots \end{aligned}$$

On the other hand, since $N_k [(\lambda_k + 1)z - \|w\|e] = [(\lambda_k + 1)z - \|w\|e]^-$, after some calculations we get

$$\begin{aligned} -\lambda_* \|w\| + \langle e, [(\lambda_* + 1)z - \|w\|e]^- \rangle + \lambda_k \|w\| - \langle e, [(\lambda_k + 1)z - \|w\|e]^- \rangle + [\|w\| - \langle e, N_k z \rangle] [\lambda_* - \lambda_k] = \\ \langle e, N_* [(\lambda_* + 1)z - \|w\|e] \rangle - \langle e, N_k [(\lambda_k + 1)z - \|w\|e] \rangle - \langle e, N_k z \rangle [\lambda_* - \lambda_k], \end{aligned}$$

for all $k = 0, 1, \dots$, where $N_* := N(\lambda_*)$. By combining the above two equalities, we obtain that

$$\lambda_* - \lambda_{k+1} = [\|w\| - \langle e, N_k z \rangle]^{-1} \left[\langle e, N_* [(\lambda_* + 1)z - \|w\|e] \rangle - \langle e, N_k [(\lambda_k + 1)z - \|w\|e] \rangle - \langle e, N_k z \rangle [\lambda_* - \lambda_k] \right].$$

Define the auxiliary piecewise linear function $\zeta(\lambda) = \langle e, N(\lambda)[(\lambda + 1)z - \|w\|e] \rangle$. Thus, except possibly at p points, ζ is differentiable and there holds

$$\zeta(\lambda_*) = \zeta(\lambda_k) + \int_0^1 \langle e, N(\lambda_k + t(\lambda_* - \lambda_k))z \rangle [\lambda_* - \lambda_k] dt,$$

due to $\langle e, N(\lambda)z \rangle \in \partial\zeta(\lambda)$. Hence, by simple combination of the two latter equalities, we have

$$\lambda_* - \lambda_{k+1} = [\|w\| - \langle e, N_k z \rangle]^{-1} \left[\int_0^1 \langle e, [N(\lambda_k + t(\lambda_* - \lambda_k)) - N_k]z \rangle dt \right] [\lambda_* - \lambda_k], \quad k = 0, 1, \dots$$

Since (14) implies that the entries of the matrix N are equal to 0 or -1 , the last inequality implies (17). Therefore, as we are under the assumption $2 \langle e, |z| \rangle < \|w\|$, we have $[\langle e, |z| \rangle] / [\|w\| - \langle e, N_k z \rangle] < 1$ and (17) implies that the sequence $\{\lambda_k\}$ converges to λ_* , which concludes the proof. \square

4.2 Picard's method

In this section we present a method based on Picard's iteration for solving equation (1). The statement of the result is

Proposition 3. *If $\langle e, |z| \rangle < \|w\|$, then for all $\lambda_0 > 0$ the sequence given by the iteration*

$$\lambda_{k+1} = \frac{1}{\|w\|} \langle e, [(\lambda_k + 1)z - \|w\|e]^- \rangle, \quad k = 1, \dots, \quad (18)$$

converges to the unique solution of the semi-smooth equation (1).

Proof. It is sufficient to prove that $\varphi : [0, +\infty) \rightarrow \mathbb{R}$ defined by $\varphi(\lambda) = \frac{1}{\|w\|} \langle e, [(\lambda + 1)z - \|w\|e]^- \rangle$ is a contraction. Indeed, the definition of φ implies

$$\begin{aligned} |\varphi(\lambda) - \varphi(\mu)| &\leq \frac{1}{\|w\|} \sum_{i=1}^n |[(\lambda + 1)z_i - \|w\|]^- - [(\mu + 1)z_i - \|w\|]^-| \\ &\leq \frac{1}{\|w\|} \sum_{i=1}^n |z_i(\lambda - \mu)| = \frac{\langle e, |z| \rangle}{\|w\|} |\lambda - \mu|, \quad \lambda, \mu \in [0, +\infty). \end{aligned}$$

Since we are under the assumption $\langle e, |z| \rangle < \|w\|$, the last inequality implies that φ is a contraction and the result follows. \square

4.3 Bisection method

In the followings we state the bisection method for finding the unique solution of (1). Let ψ be defined by (12). Since (11) implies $\|w\| > 0$, we have $\psi(0) < 0$. Then, the initializations of the method λ_0 and μ_0 can be obtained by taking

$$\lambda_0 > \max \left\{ \max \left\{ \frac{\|w\|}{z_i} : i \in \{1, \dots, n\}, z_i \neq 0 \right\} - 1, p \right\},$$

and $\mu_0 = 0$. Note that $\psi(\lambda_0)\psi(\mu_0) < 0$. Therefore, the bisection method is formally stated as follows:

0. Let $\lambda_0 > 0$ and $\mu_0 = 0$ satisfying $\psi(\lambda_0)\psi(\mu_0) < 0$. Set $k = 0$;

1. compute

$$\lambda_{k+1} = \frac{\lambda_k + \mu_k}{2}; \quad (19)$$

2. if $\psi(\lambda_{k+1}) = 0$ stop. Otherwise, compute

$$\mu_{k+1} = \begin{cases} \lambda_k, & \text{if } \psi(\lambda_k)\psi(\lambda_{k+1}) < 0, \\ \mu_k, & \text{otherwise;} \end{cases} \quad (20)$$

3. set $k \leftarrow k + 1$ and go to step 1.

It is well known that the bisection method is globally convergent with linear rate; see, for example, [23].

Final remarks

We believe that the extended second order cones will become an important class of cones in the future. First, they are probably the most natural extension of the already important second order cones. Next, the complementarity problems defined on them often have nice computational properties as remarked in the introduction. Finally, we found “almost closed-form” formulas for projecting onto them. The formulas depend only on a piecewise linear equation for a real parameter.

For a given point the projection can be obtained easily in at most 2^p steps, by assigning signs to the components of the second vector in the scalar product on the right hand side of the piecewise linear equation (1), solving for λ , and if there is a solution, then checking that the solution corresponds to the apriori assumed signs. However, this method is computationally unviable for larger p . Therefore, we developed numerical methods for solving (1) based on the semi-smooth Newton method, Picard’s iteration and bisection. The first two of these methods need some restrictions on the point which is projected to prove that they are globally convergent, but the bisection method is always globally linearly convergent.

In the future we plan to solve conic optimization and complementarity problems on the extended second order cone and to find practical examples which can be modelled by such problems.

References

- [1] F. Alizadeh and D. Goldfarb. Second-order cone programming. *Math. Program.*, 95(1, Ser. B):3–51, 2003. ISMP 2000, Part 3 (Atlanta, GA).
- [2] Miguel Sousa Lobo, Lieven Vandenbergh, Stephen Boyd, and Hervé Lebret. Applications of second-order cone programming. *Linear Algebra Appl.*, 284(1-3):193–228, 1998. ILAS Symposium on Fast Algorithms for Control, Signals and Image Processing (Winnipeg, MB, 1997).
- [3] P. Gajardo and A. Seeger. Equilibrium problems involving the Lorentz cone. *J. Global Optim.*, 58(2):321–340, 2014.
- [4] L. Kong, N. Xiu, and J. Han. The solution set structure of monotone linear complementarity problems over second-order cone. *Oper. Res. Lett.*, 36(1):71–76, 2008.
- [5] M. Malik and S. R. Mohan. On \mathbf{Q} and \mathbf{R}_0 properties of a quadratic representation in linear complementarity problems over the second-order cone. *Linear Algebra Appl.*, 397:85–97, 2005.
- [6] L. L. Zhang, J. Y. Li, H. W. Zhang, and S. H. Pan. A second order cone complementarity approach for the numerical solution of elastoplasticity problems. *Comput. Mech.*, 51(1):1–18, 2013.
- [7] K. Yonekura and Y. Kanno. Second-order cone programming with warm start for elastoplastic analysis with von Mises yield criterion. *Optim. Eng.*, 13(2):181–218, 2012.
- [8] G.-M. Luo, X. An, and J.-Y. Xia. Robust optimization with applications to game theory. *Appl. Anal.*, 88(8):1183–1195, 2009.
- [9] R. Nishimura, S. Hayashi, and M. Fukushima. Robust Nash equilibria in N -person non-cooperative games: uniqueness and reformulation. *Pac. J. Optim.*, 5(2):237–259, 2009.
- [10] C.-H. Ko, J.-S. Chen, and C.-Y. Yang. Recurrent neural networks for solving second-order cone programs. *Neurocomputing*, 74:3464–3653, 2011.

- [11] J.-S. Chen and P. Tseng. An unconstrained smooth minimization reformulation of the second-order cone complementarity problem. *Math. Program.*, 104(2-3, Ser. B):293–327, 2005.
- [12] Masao Fukushima, Zhi-Quan Luo, and Paul Tseng. Smoothing functions for second-order-cone complementarity problems. *SIAM J. Optim.*, 12(2):436–460, 2001/02.
- [13] S. Z. Németh and G. Zhang. Extended Lorentz cones and mixed complementarity problems. *J. Global Optim.*, 62(3):443–457, 2015.
- [14] Sándor Zoltán Németh and Guohan Zhang. Extended Lorentz cones and variational inequalities on cylinders. *J. Optim. Theory Appl.*, 168(3):756–768, 2016.
- [15] S.Z. Németh and G. Zhang. Positive operators of Extended Lorentz cones. [arXiv:1608.07455v2](https://arxiv.org/abs/1608.07455v2), 2016.
- [16] G. Rudolf, N. Noyan, D. Papp, and F. Alizadeh. Bilinear optimality constraints for the cone of positive polynomials. *Math. Program.*, 129(1, Ser. B):5–31, 2011.
- [17] M. S. Gowda and J. Tao. On the bilinearity rank of a proper cone and Lyapunov-like transformations. *Math. Program.*, 147(1-2, Ser. A):155–170, 2014.
- [18] M. S. Gowda and D. Trott. On the irreducibility, Lyapunov rank, and automorphisms of special Bishop-Phelps cones. *J. Math. Anal. Appl.*, 419(1):172–184, 2014.
- [19] D. W. Trott. *Topheavy and special Bishop-Phelps cones, Lyapunov rank, and related topics*. ProQuest LLC, Ann Arbor, MI, 2014. Thesis (Ph.D.)—University of Maryland, Baltimore County.
- [20] M. Orlitzky and M. S. Gowda. An improved bound for the Lyapunov rank of a proper cone. *Optim. Lett.*, 10(1):11–17, 2016.
- [21] R. Sznajder. The Lyapunov rank of extended second order cones. *Journal of Global Optimization*, 66(3):585–593, 2016.
- [22] J.-J. Moreau. Décomposition orthogonale d’un espace hilbertien selon deux cônes mutuellement polaires. *C. R. Acad. Sci. Paris*, 255:238–240, 1962.
- [23] David Kincaid and Ward Cheney. *Numerical analysis*. Brooks/Cole Publishing Co., Pacific Grove, CA, 1991. Mathematics of scientific computing.