Showing Fjortof's Theorem does not Applyfor Defining Instability for Early Universe Thermodynamic Potentials. Asking if Nucleated Particles Result at/befoe Electro-Weak Era due to Injection of Matter-Energy at the Big Bang?

Andrew Beckwith Chongqing University Department of Physics; E-mail: abeckwith@uh.edu Chongqing, PRC, 400044

Abstract

This paper uses the "Fjortoft theorem" for defining necessary conditions for instability. The point is that it does not apply in the vicinity of the big bang. We apply this theorem to what is called by T. Padmanabhan a thermodynamic potential which becomes would be unstable if conditions for the applications of "Fjortoft's theorem" hold. In our case, there is no instability, so a different mechanism has to be appealed to. In the case of vacuum nucleation, we argue that conditions exist for the nucleation of particles as of the electroweak regime. Due to injecting material from a node point, in spacetime. This regime of early universe creation, coexits with the failure of applications of "Fjortoft" theorem in such a way as to give necessary and sufficient conditons for matter creation, in a way similar to the Higgs Boson

.Key words: Fjortoft theorem, thermodynamic potential, matter creation, Higgs Boson.

1.Introduction

We first start off with a review of the classicial Fjortoft theorem [1] and from there apply it to an early universe thermodynamic potential described by T. Padamanabhan [2] in Dice 2010. The objective will be to show that one can come up with a first principle creation of nucleated "particles", likely from a semi classical stand point which can be introducing the creation of mass without appealing directly to the Higgs Boson in the first place. That due to the fact that the Fjortoft theorem does not apply.There is an inflection point for the speed up of acceleration of the universe which exists one billion years ago for reasons which we will introduce in this manuscript. But no such inflection point at the origin of the big bang itself, or at the electroweak era either.

2. Describing the Fjortoft theorem

From [1] we have that the theorem to be considered should be written up as follows, namely, look at

Fjortoft theorem:

A necessary condition for instability is that if z_* is a point in spacetime

for which $\frac{d^2U}{dz^2} = 0$ for any given potential U, then there must be some value

 z_0 in the range $z_1 < z_0 < z_2$ such that

$$\left. \frac{d^2 U}{dz^2} \right|_{z_0} \cdot \left[U(z_0) - U(z_*) \right] < 0 \tag{1}$$

For the proof, see [1] and also consider that the main discussion is to find instability in a physical system which will be described by a given potential U. Next, we will construct in the boundary of the EW era, a way to come up with an optimal description for U

2. Constructing an appropriate potential for using Fjortoft theorem in cosmology for the early universe cannot be done. We show why

To do this, we will look at Padamanabhan [2] and his construction of (in Dice 2010) of thermodynamic potentials he used to have another construction of the Einstein GR equations. To start, Padamanabhan [2] wrote

If P_{cd}^{ab} is a so called Lovelock entropy tensor, and T_{ab} a stress energy tensor

$$U(\eta^{a}) = -4 \cdot P_{ab}^{cd} \nabla_{c} \eta^{a} \nabla_{d} \eta^{b} + T_{ab} \eta^{a} \eta^{b} + \lambda(x) g_{ab} \eta^{a} \eta^{b}$$

$$= U_{gravity}(\eta^{a}) + U_{matter}(\eta^{a}) + \lambda(x) g_{ab} \eta^{a} \eta^{b}$$

$$\Leftrightarrow U_{matter}(\eta^{a}) = T_{ab} \eta^{a} \eta^{b}; U_{gravity}(\eta^{a}) = -4 \cdot P_{ab}^{cd} \nabla_{c} \eta^{a} \nabla_{d} \eta^{b}$$

(2)

We now will look at

$$U_{matter}(\eta^a) = T_{ab}\eta^a \eta^b$$
; (3)
 $U_{gravity}(\eta^a) = -4 \cdot P_{ab}^{cd} \nabla_c \eta^a \nabla_d \eta^b$

So happens that in terms of looking at the partial derivative of the top (2) equation, we are looking at

$$\frac{\partial^2 U}{\partial \left(\eta^a\right)^2} = T_{aa} + \lambda(x) g_{aa} \tag{4}$$

Thus, we then will be looking at if there is a specified η_*^a for which the following holds.

$$\begin{bmatrix} \frac{\partial^{2}U}{\partial(\eta^{a})^{2}} = T_{aa} + \lambda(x)g_{aa} \end{bmatrix}_{\eta_{0}^{a}}^{*} < 0$$

$$\begin{bmatrix} -4 \cdot P_{ab}^{cd} \left(\nabla_{c}\eta_{0}^{a}\nabla_{d}\eta_{0}^{b} - \nabla_{c}\eta_{*}^{a}\nabla_{d}\eta_{*}^{b}\right) + \\ T_{ab} \cdot \left[\eta_{0}^{a}\eta_{0}^{b} - \eta_{*}^{a}\eta_{*}^{b}\right] + \lambda(x)g_{ab} \cdot \left[\eta_{0}^{a}\eta_{0}^{b} - \eta_{*}^{a}\eta_{*}^{b}\right] \end{bmatrix}$$

$$(5)$$

What this is saying is that there is no unique point, using this η^a_* for which (5) holds. Therefore, we say there is no official point of **instability of** η_*^a due to (4). The Lagrangian structure of what can be built up by the potentials given in (4) with respect to η^a_* mean that we cannot expect an inflection point with respect to a 2nd derivative of a potential system. Such an inflection point designating a speed up of acceleration due to DE exists a billion years ago [3]. Also note that the reason for the failure for (5) to be congruent to (1) is due to

$$\left[\frac{\partial^2 U}{\partial \left(\eta^a\right)^2} = T_{aa} + \lambda(x)g_{aa}\right] \neq 0, \text{ for } \forall \eta^a_* \text{ choices}$$
(6)

** *

To use Eq (6) properly, we use the material, in our reasoning from [4], [5], [6], [7], and [8]. How and why can we do this ?

We state that (6) tells us is that there is an embedding structure for early universe geometry, some of which may take the form of the following diagram.



Figure 1, from [7,8]

3. Working with a way to achieve energy injection into the universe, without appealing to Fjortoft theorem for alleged instabilities starting from Padmanabhan thermodynamic potential terms

Padmanabhan [2] introduced the following discussion as to entropy, namely starting with energy, we have

$$E = \frac{1}{2} k_B \int dn T_{loc} \tag{7}$$

And the n value as in (7) is given by

$$dn = 32\pi \cdot P_{cd}^{ab} \cdot \varepsilon_{ab} \cdot \varepsilon^{cd} \cdot dA \tag{8}$$

Where P_{cd}^{ab} is a so called Lovelock entropy tensor, and ε_{ab} a bi normal on the co dimension -2 cross section, and then entropy is stated to be

$$S \propto \int_{\partial v} dn \propto \int_{\partial v} 32\pi \cdot P_{cd}^{ab} \cdot \varepsilon_{ab} \cdot \varepsilon^{cd} \sqrt{\sigma} d^{D-2} x$$
(9)

The end result, is that energy is induced via the temperature T_{loc} , while [2]

$$T_{loc} = N \frac{a^{\mu} n_{\mu}}{2\pi} = \text{local acceleration temperature}$$
(10)

Also, the change in n can be given by, if l_p is the Planck's length value[2]

$$\Delta n = \sqrt{\sigma} \, d^2 x / l_P \tag{11}$$

Looking at (9) and (11) we state that the change in number count given in (11) is really a holographic surface pheonmena, with N defined [2]

$$N = E / [(1/2)k_B T]$$

$$\tag{12}$$

The upshot is that we can, as implied by Ng[4] easily reference a change in entropy via[4],[5],[6]

 $S \sim n$

(13)

While having a change in n as due to a change in the spatial surface of spacetime as given in (11), we have to realistically infer that the local acceleration temperature (10) is from another pre universe contruction and that local instability is ruled out by (5) and (6). This leads us to ask as to what would be an acceptable way to form the formation of mass, i.e. say the mass of a graviton, via external factors introduced into our universe prior to the Electroweak era, in cosmology. To do that, look at if there are two branes on the AdS_5 space-time so that with one moving and one stationary, we can look at Figure 1 as background as to introduce such external factors in our present space-time universe during its initial expansion phase

4. Fall out from adopting Figure 1 and that due to no instability in the Padamanabhan supplied potentials. i.e. a way to obtain graviton mass via a root finding method.

Using [7], what we find is that there are two branes on the AdS_5 space-time so that with one moving and one stationary, we can look at figure 1 which is part of the geometry used in the spatial decomposition of the differential operator acting upon the h_{\bullet} Fourier modes of the h_{ij} operator [7]. As given by [7],[8] we have that

$$\left[\partial_t^2 + k^2 - \partial_y^2 + \frac{3}{y} \cdot \partial_y\right] h_{\bullet} = 0$$
(14)

Using [8] (and also [7]) the solution to (14) above takes the form of having

$$h_{\bullet} = H_{ij} = e_{ij} \cdot \exp[i \cdot \omega \cdot t] \cdot (m \cdot y)^2 \cdot A \cdot J_2(m \cdot y)$$
(15)

 e_{ij} is a polarization tensor, and the function $J_2(my)$ is a 2nd order Bessel function [14]. A generalization offered by Durrer et al. [7], [8] leads to

$$h = \left\{ \exp[i \cdot \omega \cdot t] \cdot (m \cdot y)^2 \cdot A \cdot J_2(m \cdot y) \right\} \cdot \left(1 + \frac{\pi}{4} \cdot (m \cdot \ell)^2 \right)$$
(16)

With the factor of $\left(1 + \frac{\pi}{4} \cdot (m \cdot \ell)^2\right)$ coming in due to a boundary condition upon the wall of a brane put in, i.e. looking at [7]. With the right hand side of (16) due to a domain wall tension of a brane.

$$-2 \cdot \partial_{y} H_{ij} = \kappa_{5} \cdot \pi_{ij}^{(T)} \to 0$$
⁽¹⁷⁾

This will be in our example set as not equal to zero, in the right hand side, but equal to an extremely small parameter, namely

$$\partial_{y}H_{ij}\Big|_{y=yb} = \kappa_{5} \cdot \pi_{ij}^{(T)} \sim \xi^{+}$$
(18)

With this turned into

$$\partial_{y}h\Big|_{y=yb} \sim \delta^{+} \tag{19}$$

The right hand side of (19) represents very small brane tension, which is understandable. Then using [7], [8], [9], i.e.

$$\partial_{y}h\Big|_{y=yb} = \partial_{y}\left\{\exp[i\omega t]\cdot(my)^{2}\cdot A\cdot J_{2}(my)\right\}\cdot\left(1+\frac{\pi}{4}\cdot(m\cdot\ell)^{2}\right)\Big|_{y=yb}\sim\delta^{+}$$
(20)

and

$$J_{2}(my) = \frac{(my)^{2}}{2^{2} \cdot 2!} \cdot \left(1 - \frac{(my)^{2}}{2^{2} \cdot 3} + \frac{(my)^{4}}{2^{4} \cdot 2! \cdot 3 \cdot 4} - \frac{(my)^{6}}{2^{6} \cdot 4! \cdot 3 \cdot 4 \cdot 5} + \dots\right)$$
(21)

The upshot is, that afterwards,

$$\frac{(my)^{4}}{2^{2} \cdot 2!} \cdot \frac{1}{y} \begin{bmatrix} \left(1 - \frac{(my)^{2}}{2^{2} \cdot 3} + \frac{(my)^{4}}{2^{4} \cdot 2! \cdot 3 \cdot 4} - \frac{(my)^{6}}{2^{6} \cdot 4! \cdot 3 \cdot 4 \cdot 5} + ...\right) \\ - \left(\frac{2 \cdot (my)^{2}}{2^{2} \cdot 3} + \frac{4 \cdot (my)^{4}}{2^{4} \cdot 2! \cdot 3 \cdot 4} - \frac{6 \cdot (my)^{6}}{2^{6} \cdot 4! \cdot 3 \cdot 4 \cdot 5} + ...\right) \end{bmatrix}$$
(22)
$$= \frac{\delta^{+} \cdot \exp[\mp i\omega t]}{A} \cdot \left[1 - \frac{\pi}{4} \cdot (m \cdot \ell)^{2}\right]$$

Should the term

$$\frac{\delta^{+} \cdot \exp[\mp i\omega t]}{A} \cdot \left[1 - \frac{\pi}{4} \cdot \left(m \cdot \ell\right)^{2}\right] \xrightarrow{\delta^{+} \to 0} 0$$
(23)

Then, (22) is acting much as in [7], and [8], whereas, one is recovering a simple numerical exercise as to obtain a suitable solution as given by (18), and (19) due to [7,8] where the domain tension of the brane vanishes. The novelty as to this approach given in (22) is to obtain a time dependent behavior of the mass of the graviton,

$$(my) = f(t) \Leftrightarrow m \equiv \frac{f(t)}{y}$$
(24)

In doing all of this, keep in mind the mathematical information given in [9] which is repeatedly used in [7] and [8];

Needless to say, (22) can only be solved for, numerically, i.e. fourth order polynomial solutions for quartic equations still give over simplified dynamics, especially if (24) holds, and makes things more complicated. This is all being done to keep fidelity with respect to [10], and [11] as a possible feature of brane world dynamics as reflected in [10],[11], as well as certain issues brought up in [12], as to what is a semi classical argument can obtain a usually quantum result. It also would allow for eventually understanding if entropy can also be stated in terms of gravitons alone in early universe models as was proposed by Kiefer & Starobinsky, et. al. [13]. Finally, it would address if QM is embedded in a larger deterministic theory as advocated by t' Hooft [14], as well as degrees of freedom in early universe cosmology as brought up by Beckwith in Dice 2010 [15]. We argue that making this step is consistant with keeping the value of Planck's constant uniform in spite of Avessans theories suggesting it vary in time[16] . To do this, we make extensive use of [17] and [18].

It is now then time to do a re cap and to organize how such speculation can be vetted using experimental proceedures. To do this we re cap what can be said about traces massive gravitons can be detected, prior to our conclusion which MAY LEAD TO AN EXPLANTION OF THE FOLLOWING ENTROPY FORMULA[19]

$$S_{gw} = V \cdot \int_{v}^{v} \int_{0}^{1} r(v) \cdot v^2 dv \cong (10^{29})^3 \cdot (H_1/M_P)^{3/2} \approx 10^{87} - 10^{88}$$
(25)

This is a bridge to future projects which should be kept in mind. I.e, could our formulation of graviton physics lead to identification of gravitons, in the early universe as the main driver of graviton physics being the primary entropy generator as suggested by [19]?

. 5. CONCLUSION Semiclassical method of obtaining graviton mass procedure cannot be ruled out, and it impacts relic GW searches

First of all, review the details of a massive graviton imprint upon h_{ij} , and then we will review the linkage between that and certain limits upon h_{\bullet} . As read from Hinterbichler [20], if $r = \sqrt{x_i x_i}$, and we look at a mass induced h_{ij} suppression factor put in of $\exp(-m \cdot r)$, then if

$$h_{00}(x) = \frac{2M}{3M_{Planck}} \cdot \frac{\exp(-m \cdot r)}{4\pi \cdot r}$$
(26)

$$h_{0i}(x) = 0$$
 (27)

$$h_{ij}(x) = \left[\frac{M}{3M_{Planck}} \cdot \frac{\exp(-m \cdot r)}{4\pi \cdot r}\right] \cdot \left(\frac{1 + m \cdot r + m^2 \cdot r^2}{m^2 \cdot r^2} \cdot \delta_{ij} - \left[\frac{3 + 3m \cdot r + m^2 \cdot r^2}{m^2 \cdot r^4}\right] \cdot x_i \cdot x_j\right)$$
(28)

Here, we have that these h_{ij} values are solutions to the following equation, as given by [21], with D a dimensions value put in.

$$\left(\partial^{2} - m^{2}\right)h_{\mu\nu} = -\kappa \cdot \left[T_{\mu\nu} - \frac{1}{D-1} \cdot \left(\eta_{\mu\nu} - \frac{\partial_{\mu}\partial_{\nu}}{m^{2}}\right) \cdot T\right]$$
(29)

To understand the import of the above equations, set

$$M = 10^{50} \cdot 10^{-27} g \equiv 10^{23} g \propto 10^{61} - 10^{62} eV$$

$$M_{Plank} = 1.22 \times 10^{28} eV$$
(30)

We should use the $m_{massive-graviton} \sim 10^{-26} eV$ value in (29) above.

In reviewing what was said about (27),(28) we should keep in mind the overall Fourier decomposition linkage between h_{\bullet} , h_{ii} which is written up as

$$h_{ij}(t,x;k) = \frac{1}{(2\pi)^{3/2}} \int d^3k \sum_{\bullet=+,\otimes} e^{ik \cdot x} e^{\bullet}_{ij} h_{\bullet}(t,y;k)$$
(31)

The bottom line is that the simple de composition with a basis in two polarization states, of $+, \otimes$ will have to be amended and adjusted, if one is looking at massive graviton states.

In addition further developments as to (31) could influence giving a semi classical interpretation as to entrophic origins of gravity, along the lines brought up by both t'Hooft, indirectly [14], and Lee [22] directly.

The experimental gravity considerations are covered in [23], [24], [25], and [26], and the idea should be especially to in our work to examine if [23] and [25] in terms of gravity are adhered to. As these are LIGO projects, we should be looking to see if what we are doing contravenes or backs the post Newtonian approximations of physics, so brought up.

Reference [25] is a must to review. In it, Corda reviews GR tests and our document must not contravene these basics. Can we obtain through our representation of gravitons, confirmation, or refutation of if the data sets are in adherence, or partially refute General Relativity.

As far as [26], in terms of quantum cosmology, it is another similar parallel development to the ideas raised here. I urge readers to investigate it.

Finally, in lieu of [27] the author urges readers to look at Appendix A, which is a summary of what the author views as to what would be foundational investigation of gravity, and to see if it can be made in adherence to GR, and possibly quantized.

The author urges readers to look at [27], as well as [28] by Kieffer, and that when seeing **Appendix A**, that this is a schematic the author believes would be appropriate for an investigation to confirm if Gravity could be derived as having quantum roots. Which in turn would affect the viability of presumed quantum gravity

Acknowledgements

This work is supported in part by National Nature Science Foundation of China grant No110752

References

[1] J. Pringle and A. King; Astrophysical Flows, Cambridge University Press, New York, 2007

[2] Thanu Padmanabhan 2011 J. Phys.: Conf. Ser. 306 01200;"Lessons from Classical Gravity about the Quantum structure of space-time", DICE 2010, editor T. Elze, <u>http://iopscience.iop.org/1742-6596/306/1/012001</u>

[3] A. Beckwith, "Identifying a Kaluza Klein Treatment of a Graviton Permitting a deceleration parameter Q (Z) as an Alternative to Standard DE", Journal of Cosmology, 2011, Vol. 13 http://journalofcosmology.com/BeckwithGraviton.pdf

[4] Y.J. Ng, Entropy 10(4), pp441-461 (2008); Y.J. Ng and H. van Dam, FoundPhys30, pp795–805 (2000);

[5] Y.J. Ng and H. van Dam, Phys. Lett. B477, pp429–435 (2000);

[6]Y.J. Ng, https://cgc.physics.miami.edu/Miami2012/Ng.pdf

[7] R. Durrer et.al., "Dynamical Casmir effect for gravitons in Bouncing Braneworlds", 2009,

http://theory.physics.unige.ch/~durrer/papers/casimir.pdf; Earlier version http://www.scribd.com/doc/79001898/Ruth-Durrer-and-Marcus-Ruser-

Thedynamical-Casimir-effect-in-braneworlds ; Version put into PRD 10 Marcus Ruser, Ruth Durrer, "Dynamical Casimir effect for gravitons in bouncing braneworlds", http://arxiv.org/abs/0704.0790; Phys.Rev.D76:104014,2007

[8] R. Durrer, "Branewords", http://arxiv.org/pdf/hep-th/0507006v1.pdf

[9] Bessel functions and Hankel Functions: NIST: <u>http://dlmf.nist.gov/10.2</u>
[10] J. Koury, P.J. Steinhardt, N. Turkok, PRL92,

031302(2004),hepth/0307132

[11] J. Koury, P.J. Steinhardt, N. Turkok, PRL91,

161301(2003),astroph/0302012

[12] C.Cartier, Ru. Durrer, M. Ruser, PRD 72, 104018(2005),hep-th/0510155
[13] C. Kiefer, D. Polarski, A.A Starobinsky, "Entropy of gravitons produced in the early Universe", PRD 62, 043518 (2000) [6 pages]; arxiv grqc/9910065

[14] G. t' Hooft, "Quantum Mechanics as a dissipative deterministic system" http://arxiv.org/PS_cache/gr-qc/pdf/9903/9903084v3.pdf

[15] A. Beckwith, "How to use the cosmological Schwinger principle for energy flux, entropy, and "atoms of space–time" to create a thermodynamic space–time and multiverse", <u>http://iopscience.iop.org/1742-</u>

<u>6596/306/1/012064</u>, in "5th International Workshop DICE2010: Space-Time-Matter – Current Issues in Quantum Mechanics and Beyond", T. Elze ed. <u>http://iopscience.iop.org/1742-</u>

<u>6596/306/1;jsessionid=A05372A78C18D970BF35F40A9A863B51.c2</u>

[16] A.K. Avessian, "Planck's constant evolution as a cosmological

evolution test for the early universe", .Gravitation and Cosmology, 2009, Vol. 15, number 1, pp 10-12

[17] U. Bruchholz; "Derivation of Planck's constant from Maxwell's Electrodynamics", Progress in Physics, V4, October 2009, page 67; http://www.ptep-online.com/index_files/2009/PP-19-07.PDF

[18]U. Bruzchholz, "Key notes on a geometric theory of fields", Progress in Physics, 2009, v.2, 107–113.

[19] Giovannini, Massimo, " A Primer on the Physics of the Cosmic Microwave Background", World Press Scientific, 2008, Singapore, Republic of Singapore, (2008)

[20] K.Hinterbichler, "Theoretical Aspects of Massive Gravity", http://arxiv.org/abs/1105.3735.pdf

[21] http://web.mit.edu/redingtn/www/netadv/Xgravitati.html

[22] Jae-Weon Lee, "On the Origin of Entropic Gravity and Inertia", http://arxiv.org/abs/1003.4464

[23] B. P. Abbott et al., "Observation of Gravitational Waves from a Binary Black Hole Merger", PRL 116, 061102 (2016);

https://physics.aps.org/featured-article-pdf/10.1103/PhysRevLett.116.061102 [24] Corda, C. (2009) Interferometric Detection of Gravitational Waves: The Definitive Test for General Relativity. International Journal of Modern Physics D, 18, 2275-2282. ArXiv: 0905.2502 [gr-gc]

[25] B. P. Abbot et.al. "GW151226: Observation of Gravitational Waves from a 22-Solar-Mass Binary Black Hole Coalescence" Published in **Phys. Rev. Lett. 116**, 241103 (**2016**)

[26] Halliwell, J., "Quantum Cosmology" pp 515 -599, in "The ICTP Serioes in Theoretical Physics, Volume 7, 1990 Summer School in High energy Physics and Cosmology, Trieste, Italy 18 June to 28 July, 1990, Editors Pati, J. Randjbar-Daemi, Sezgin, E. Shafi, Q, Word Scientific, Singapore, Republic of Singapore, 1991 [27] Andrew Beckwith, "History Lessons from the 5th Solvay Meeting, 1927", accepted as talk in a September 15 conference, in Kiev, Ukranine, 2017, <u>http://vixra.org/abs/1708.0399</u>

[28] Kieffer, Claus; "Quantum Gravity, 3rd edition", Oxford Science Publications, Oxford University Press, Oxford, United Kingdom, 2012
[29] Dirac, Paul, 'Lectures on Quantum Mechanics' Dover Publications, Inc, Mineola, New York, USA 2001

[30] Ryden, Barbara, 'Introduction to Cosmology, 2nd Edition', Cambridge University Press, Cambridge, United Kingdom, 2017

[31] G. Bacciagaluppi and A. Valentini Quantum Theory at the Crossroads: Reconsidering the 1927 Solvay Conference (Cambridge University Press, 2009), Cambridge, Massachusetts, USA

[32] Roland Omnes , "Understanding Quantum Mechanics", Princeton University Press, Princeton, New Jersey, USA, 1999

[33] Roland Omnes , "The Interpretation of Quantum Mechanics", Princeton University Press, Princeton, New Jersey, USA,1994

Appendix A, from [27], a quote of would be first principles of how to quantize the Einstein field equations elucidated by first principles:

From [27] where we quote large parts of our text in [27]

XIX. A generalized problem to making quantization of the Einstein field equations elucidated by first principles.(This reflects our evaluation of [28])

i.e. the lectures on quantization of a classical Hamiltonian given by Dirac, in [29] [108], pages 25 – 43 is ironically made more fraught by the requirement of extending the Hamiltonian i.e. if we have say $\phi_{a'}$ as so called first class secondary constraints, page 25 of [29][108] we find that there is an inability to do the

following, if we wish to transfer to quantum systems, we need to do the following, i.e. add to the initial classical Hamiltonian, H_{τ}

$$\begin{aligned} H_E &= H_T + v_{a'} \cdot \phi_{a'} \\ \dot{g} &\approx \left\{ g, H_E \right\} \end{aligned} \tag{A1}$$

Eq. (A1), in a Poisson bracket formulation, was used by Dirac to transform to a set of quantization conditions, in pages 25 to 43 of. The problem is, that it is difficult to come up with constraint equations, as given in the top level of Eq.(A1)

The following is easy to do, if you ignore constraints

$$\frac{d\langle P \rangle}{dt} = -\frac{i}{\hbar} \langle \Psi \| [P, H] | \Psi \rangle = \frac{1}{i\hbar} \langle \Psi \| [P, V] | \Psi \rangle$$

$$\xrightarrow{3\dim \to 1\dim} -\int \Psi^* \cdot \frac{dV(x)}{dx} \cdot \Psi dx$$

$$\xrightarrow{1\dim \to Any \ \dim} \langle -\frac{dV(x)}{dx} \rangle \sim \langle -\vec{\nabla}V \rangle \equiv F(force)$$
(A2)

Try doing this, to have equivalence with Eq. (A1) and match that with Eq.(A2) the equations below given as (A3) to Eq. (A4). i.e. what is so difficult is to put in a Hamiltonian system, for gravity, which is comme Worse than that, we do not have a quantum mechanical equivalent, and this due to the difficulties in terms of finding a quantum mechanical equivalent to the Poisson brackets $\{p_N, H(Hamiltonian)\} \approx 0$ which is readily transferrable to the Friedman equation , i.e. so far a quantum bridge between quantized versions of Eq. (A3) and Eq. (A4)does not exist, right now. As we wrote in [27]

So as given in [27], as stated by Kieffer, [28] there is a relationship between a Hamiltonian form,, H(Hamiltonian), and a constraint equation, for momentum p_N , along the lines of

$$p_{N} \\ \& \qquad (A3) \\ \left\{ p_{N}, H(Hamiltonian) \right\} \approx 0$$

This is , according to Kieffer, the Poisson brackets, equivalent to the following

What we are looking at is, if we set the Lapse function, N, as = 1

$$\dot{a}^{2} = -1 + a^{2} \cdot \left(\dot{\phi}^{2} + \frac{\Lambda}{3} + m^{2} \cdot \phi^{2}\right)$$

$$\Leftrightarrow \ddot{\phi}^{2} + 3\frac{\dot{a}}{a} \cdot \dot{\phi} + m^{2}\phi = 0$$
(A4)

Here, the ϕ is a scalar field (here, called a 'homogeneous field'), m is amass term, and a the scale factor, and Λ the cosmological constant. If m is set equal to zero, this has a simple m= 0 solution with

$$p_{\phi} = a^{3} \cdot \dot{\phi} = \kappa = const$$

$$\& \qquad (A5)$$

$$\phi = \pm \frac{1}{2} \cdot ar \cosh \frac{\kappa}{a^{2}}$$

It cannot be solved analytically, if m is not equal to zero. Now as to a general problem between the Solvay 1927 conference methods and the application to GR will be alluded to, next

Dirac claims the bridge from Poisson brackets to the situation represented by Eq. (A5) always involves a carefully set extended Hamiltonian situation. I,.e. see his discussion in 33 to page 35 of [29]. The challenge would be to make those extensions somehow commensurate with Eq. (A4) and Eq. (A5)

Having said, this, we will next go to the problem of Quantum Geometrodynamics. Before going to it, a notice as to the problems of bridging to general relativity using conventional Quantum mechanics, will be raised as a bridge to the use of $H_{ADM} \Psi = 0$ which makes a plausible bridge to the Fluid equation of general relativity, [30] but also a summary as to how and why the connection to the rest of general relativity is extremely difficult, i.e. the Friedman equation as seen in [30]has a classical analogue which cannot be linked to its general relativistic form, but the fluid equation of General relativity in [30] does have a Newtonian derivation yielding the exact same result in both Newtonian and GR physics. Hence, the quantum-classical bridge as exemplified by Eq.(A5) works for the fluid equation, but would not work for the GR Friedman equation, since the Friedman equation classical would be the only bridge to the quantum result, using the Eq. (A5) bridge.

And of course, both the GR Friedman bridge plus the fluid cosmology bridge are both needed in the acceleration equation, i.e. from [30] the following cannot be linked to quantum mechanics, via Eq. (A5), namely the acceleration equation of GR has

$$\left(\frac{\ddot{a}}{a}\right) = -\frac{4\pi G}{3c^2} \cdot \left(\varepsilon + 3P\right) \tag{A6}$$

This requires two equations, namely,

$$\left(\frac{\ddot{a}}{a}\right) = -\frac{4\pi G}{3c^2} \cdot (\varepsilon + 3P)$$

$$Due - to$$

$$\dot{\varepsilon} + 3\left(\frac{\dot{a}}{a}\right) \cdot (\varepsilon + 3P) = 0 \qquad (A7)$$

$$And$$

$$\dot{a}^2 = \frac{8\pi G}{3c^2} \cdot \varepsilon a^2 - \frac{\kappa c^2}{R_0^2} (GR - Friedman)$$

$$as - opposed - to, if \quad U = const.$$

$$\dot{a}^2 = \frac{8\pi G}{3c^2} \cdot \rho a^2 + \frac{2U}{r_c^2} (Newtonian - Friedman)$$

The derivation of the acceleration equation for GR, using the two equations cited is in[30], page 60

In addition we will derive the Fluid equation also used, which is the same form used in Eq.(A5) making a linkage to relativity and quantum mechanics, possible, if one uses the following steps, as given on page 59 of [30] I.e. If exists a commoving radius r_s

We then will get a clean derivation of the so called fluid equation, used in Cosmology. This fluid equation, which has the same form used in both GR and Newtonian physics may be in principle linkable to the quantization program outlined in Eq. (A5). So with that, we go to the interactions given in Eq.(A8) below.

$$V(t) = Volume(universe) = \frac{4\pi r_s^3 a^3}{3}$$

$$\dot{w}$$

$$\dot{V} = V \cdot \left(\frac{3\dot{a}}{a}\right)$$

$$E = V(t) \cdot \varepsilon(t)$$

$$\dot{E} = V \cdot \left(\dot{\varepsilon} + \frac{3\dot{a}}{a}\varepsilon\right)$$

$$\dot{w}$$
First - law - thermo(universe)
$$\dot{E} + P\dot{V} = 0$$

$$\Rightarrow V \cdot \left(\dot{\varepsilon} + \frac{3\dot{a}}{a}\varepsilon + \frac{\dot{a}}{a}P\right) = 0$$

$$\Rightarrow \dot{\varepsilon} + \frac{3\dot{a}}{a}\varepsilon + \frac{\dot{a}}{a}P = 0$$
(A8)

The GR and classical physics forms of the fluid equation, so derived, in Eq.(A8) and the results at the bottom of Eq. (A7) would allow us to make connection, with a lot of work to the sort of reasoning used in Eq. (A5) above, but due to the difference in the Friedman equation, in classical and GR form, as noted in Eq.(A6), it would be using the Solvay methods, extremely difficult to make connection between an acceleration equation, using scale factors, as given in Eq. (A6) and Eq. (A7) with the Eq. (A5)(58) connection between classical and quantum mechanics with respect to an acceleration of the universe acceptable in both GR and quantum form.

We can state though that a bridge to the Fluid equation, as given i[27]n Eq.(A8) and Eq.(A5) would at least in principle very doable. Having said that, let us now go to the ideas of Quantum Geometrodynamics, as far as their use and future prospects to the study of Solvay 1927 methods [31], and quantum gravity issues[28]

End of our quote from [27]

As a close to this, all this, in terms of quantum gravity should also keep in mind issues brought up in [31], and [32], in particular, quantum entanglement and how information is transferred in cosmology. i.e. the Geometrodynamics part of [28]