

Chen's Formulas of the Fine-structure Constant (viXra:2002.0203)

Gang Chen[†], Tianman Chen and Tianyi Chen

Guangzhou Huifu Research Institute Co., Ltd., Guangzhou, P. R. China

7-20-4, Greenwich Village, Wangjianglu 1, Chengdu, P. R. China

[†]Correspondence to: gang137.chen@connect.polyu.hk

Dedicated to Prof. Albert Sun-Chi Chan on the occasion of his 70th birthday

Abstract

This paper gives two series of formulas of the fine-structure constant α which are reasonable, precise, smart and elegant. It also demonstrates there are two values of α , i.e., $\alpha_1=1/137.035999037435$ and $\alpha_2=1/137.035999111818$, which are consistent with but much more accurate than those experiment measured values. The formulas consist of 2π -e formulas and some factors related to nucleon numbers of nuclides. A brief explanation of the fine-structure constant shows $1/\alpha \approx 137.036$ is the equal ratio factor between 112 and 168 (more precisely 168-1/3). Based on these, all 119th to 170th ideal extended elements were predicted, the speed of light in atomic units was mathematically calculated by $c_{au}=1/(\alpha_1\alpha_2)^{1/2}=137.035999074627$, Schrödinger equation of hydrogen atom was simplified and correlated with α_1/α_2 , classical electron radius was calculated to be 2.81794032658(43) fm and proton charge radius was hypothetically calculated to be 0.833027202999(13) fm. In the end, it was found that the approximate rational numbers of 2π marvelously related to nuclides, a mathematic shell model of nuclides was established and a picture of elements and ideal extended elements was depicted.

Keywords: formulas; the fine-structure constant; the ideal extended elements; the speed of light; Schrödinger equation of hydrogen atom; the proton charge radius; 2π .

1. Introduction

The fine-structure constant (Sommerfeld constant) is a critical dimensionless constant in physics, it is a century mystery of physics, it has been one of the biggest enigmas in physics since it was introduced by Arnold Sommerfeld in 1916. Its definition, some interpretations and the latest measured values are as follows^{1,2}:

$$\alpha = \frac{\lambda_e}{2\pi a_0}, \quad \alpha = \frac{2\pi r_e}{\lambda_e}, \quad \frac{a_0}{r_e} = \frac{1}{\alpha^2}; \quad \alpha = \frac{e^2}{4\pi\epsilon_0\hbar c} = \frac{v_e}{c}, \quad \frac{c}{v_e} = \frac{1}{\alpha}$$

in atomic units, the speed of light $c_{au} = \frac{1}{\alpha}$

the 2014 CODATA recommended value: $\alpha = 1/137.035999139(31)$

the 2018 CODATA recommended value: $\alpha = 1/137.035999084(21)$

Science 13 April 2018 reported value: $\alpha = 1/137.035999046(27)$

The ratio of Bohr radius of hydrogen atom a_0 to the classical electron radius r_e is $1/\alpha^2$. The ratio of the speed of light c to the line velocity of ground state electron in hydrogen atom v_e is $1/\alpha$, this means in atomic units $c=1/\alpha$ and $E=mc^2=m/\alpha^2$ or $\alpha^2=m/E$. In quantum electrodynamics it substantially characterizes the strength of electromagnetic interaction between elementary charged particles such as electron and proton, so it is the coupling constant of electric charges. It is one of the 25 fundamental constants (could not be calculated theoretically, could only be determined by experiments) in Standard Model of physics and should be the most important one. As it is dimensionless, it could be called the proportional ruler of the nature or the bridge of mathematics and physics. However, to our knowledge, up to now (except this work), no one knows how it comes from, no one could give reasonable explanations to it or formulas of it since it was introduced.

In 2016 Paul Davis gave the following comment³: “Physicists have long wondered where this number, $1/137.035999$, comes from. Is there a deep reason why α has to be precisely this number for the world to function as it does? There is a long history of attempts to derive α from physical theory or to concoct a mathematical formula that has this value. For a brief time in the 1920s, when it looked as if α might be exactly $1/137$, astronomer Arthur Eddington searched for a theory that would throw up the numbers naturally, but his ideas ultimately led nowhere. Then in 1969 a young Swiss mathematician, Armand Wyler, pointed out that $(9/16\pi^3)(\pi/5!)^{1/4}$ comes close to $1/137.036$, which matched the value of α to the precision known at the time. However, his formula was not accompanied by any credible theory and was regarded as little more than a numerical curiosity. Several other attempts at α numerology have been made since, none of which have gained traction in the physics community.”

As for the fascination of the fine-structure constant, in the middle of 1980s, Richard Feynman stated⁴: “It has been a mystery ever since it was discovered more than fifty years ago, and all good theoretical physicists put this number up on their wall and worry about it. Immediately you would like to know where this number for a coupling comes from: is it related to pi or perhaps to the base of natural logarithms? Nobody knows. It's one of the greatest damn mysteries of physics: a magic number that comes to us with no understanding by man. You might say the hand of God wrote that number, and we don't know how He pushed his pencil.”

This paper shows how God pushed his pencil to write the fine-structure constant and how God used it to coordinate elements.

2. 2π -e formula(s)

2π -e formula, its related formulas and their preliminary applications were deduced independently by us from April to December of 2013.

Fig. 1. Diagram of $y=1/x$.





Fig. 2. Diagram of $y=\log(x)$.



$$\text{Euler-Mascheroni constant } \gamma : \sum_{n=1}^{\infty} \frac{1}{n} = 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{\infty} = \ln \infty + \gamma$$

$$\text{As for } y = 1/x \text{ (Fig. 1), } \gamma = 0.577215\dots = 0.5 + 0.077215\dots = \sum_{n=1}^{\infty} \delta_{2n} + \sum_{n=1}^{\infty} \delta_{1n} = \frac{1}{2} + \gamma_1$$

$$\gamma_1 = \sum_{n=1}^{\infty} \delta_{1n} = \lim_{N \rightarrow \infty} \left(\sum_{n=1}^{N-1} \frac{1}{n} - \int_1^N \frac{1}{x} dx \right) - \frac{1}{2}, \text{ Generally } \gamma_s = \lim_{N \rightarrow \infty} \left(\sum_{n=1}^{N-1} \frac{1}{n^s} - \int_1^N \frac{1}{x^s} dx \right) - \frac{1}{2}, \quad s \in \mathbb{N}$$

$$\begin{aligned} \text{As for } y = \log(x) \text{ (Fig. 2), } \delta_{1,n} &= \int_n^{n+1} \log x dx - \frac{1}{2} \ln \frac{n+1}{n} - \ln n = (x \ln x - x) \Big|_n^{n+1} - \frac{1}{2} \ln(n+1)n \\ &= (n+1) \ln(n+1) - n \ln n - 1 - \frac{1}{2} \ln(n+1) - \frac{1}{2} \ln n = \left(n + \frac{1}{2}\right) \ln\left(1 + \frac{1}{n}\right) - 1 \end{aligned}$$

$$\gamma_{c,N} = \sum_{n=1}^N \delta_{1,n} = \sum_{n=1}^N \left[\left(n + \frac{1}{2}\right) \ln\left(1 + \frac{1}{n}\right) - 1 \right] = \sum_{n=1}^N \ln \frac{\left(1 + \frac{1}{n}\right)^{\left(n + \frac{1}{2}\right)}}{e} = \ln \prod_{n=1}^N \frac{\left(1 + \frac{1}{n}\right)^{\left(n + \frac{1}{2}\right)}}{e}$$

$$\gamma_c = \gamma_{c,\infty} = \sum_{n=1}^{\infty} \delta_{1,n} = \lim_{N \rightarrow \infty} \left(\int_1^{N+1} \log(x) dx - \sum_{n=1}^N \log(n) - \frac{\log(N+1)}{2} \right)$$

$$= \sum_{n=1}^{\infty} \left[\left(n + \frac{1}{2}\right) \ln\left(1 + \frac{1}{n}\right) - 1 \right] = \sum_{n=1}^{\infty} \ln \frac{\left(1 + \frac{1}{n}\right)^{\left(n + \frac{1}{2}\right)}}{e} = \ln \prod_{n=1}^{\infty} \frac{\left(1 + \frac{1}{n}\right)^{\left(n + \frac{1}{2}\right)}}{e}$$

$$\ln N! = \sum_{n=1}^N \ln n = \int_1^{N+1} \ln x dx - \sum_{n=1}^N \delta_{1,n} - \sum_{n=1}^N \delta_{2,n} = (x \ln x - x) \Big|_1^{N+1} - \ln e^{\gamma_{c,N}} - \sum_{n=1}^N \frac{\ln(n+1) - \ln n}{2}$$

$$= (N+1) \ln \frac{(N+1)}{e} + \ln \frac{e}{e^{\gamma_{c,N}}} - \frac{1}{2} \ln(N+1) = \ln \left[\frac{e^{1-\gamma_{c,N}}}{\sqrt{N+1}} \left(\frac{N+1}{e} \right)^{(N+1)} \right]$$

$$N! = \frac{e^{1-\gamma_{c,N}}}{\sqrt{N+1}} \left(\frac{N+1}{e} \right)^{(N+1)}, \text{ compared to Stirling formula : } N! \sim \sqrt{2\pi N} \left(\frac{N}{e} \right)^N$$

$$(N+1)! = (N+1)N! \sim \sqrt{2\pi(N+1)} \left(\frac{N+1}{e} \right)^{N+1}, \quad N! \sim \frac{\sqrt{2\pi}}{\sqrt{N+1}} \left(\frac{N+1}{e} \right)^{N+1}$$

$$\text{Compared to previous formula, gives } \sqrt{2\pi} \sim e^{1-\gamma_{c,N}} \text{ or } 2\pi = \left(\frac{e}{e^{\gamma_c}} \right)^2$$

$$2\pi - e \text{ formula(s): } 2\pi = \left(\frac{e}{e^{\gamma_c}} \right)^2 = e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \frac{e^2}{\left(\frac{4}{3}\right)^7} \dots, \quad (2\pi)_k = \left(\frac{e}{e^{\gamma_{c,k}}} \right)^2 = e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{k+1}{k}\right)^{2k+1}}$$

$$\gamma_c = 0.0810614668, \quad e^{\gamma_c} = 1.0844375$$

2π -e formula is an expanding form of Stirling formula. To our knowledge, it was first deduced by us. If it was new, it could be named Chen's 2π -e formula.

3. Some Formulas Related to 2π -e Formula

The following formulas which correlate each other and has similar form could be called Chen's natural group formulas, and the form is called natural group.

$$\begin{aligned}
1 &= 4\gamma_c + \frac{4\gamma_1}{1(1+1)} + \frac{4\gamma_2}{2(2+1)} + \frac{4\gamma_3}{3(3+1)} + \dots \\
&= |B| \frac{\pi}{2} + \sum_{n=1}^{\infty} \frac{|B_{2n}|(\pi/2)^{2n}}{(2n)!} = \sum_{n=1}^{\infty} \frac{|B_{2n}|\pi^{2n}}{(2n)!} = -|B| \frac{3\pi}{2} + \sum_{n=1}^{\infty} \frac{|B_{2n}|(3\pi/2)^{2n}}{(2n)!} \\
N &\sim -\frac{3}{2}|B| + \sum_{n=1}^N \frac{|B_{2n}|(2\pi)^{2n}}{2(2n)!} \\
e &= 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \dots \\
2\pi &= \left(\frac{e}{e^{\gamma_c}}\right)^2 = e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \frac{e^2}{\left(\frac{4}{3}\right)^7} \dots
\end{aligned}$$

B, B_{2n} : the Bernoulli numbers such as $-\frac{1}{2}, -\frac{1}{6}, -\frac{1}{30}, \frac{1}{42}, -\frac{1}{30}, \dots$

$$\gamma_c = \lim_{N \rightarrow \infty} \left(\int_1^{N+1} \log(x) dx - \sum_{n=1}^N \log(n) - \frac{\log(N+1)}{2} \right) = 0.0810614668$$

$$\gamma_s = \lim_{N \rightarrow \infty} \left(\sum_{n=1}^{N-1} \frac{1}{n^s} - \int_1^N \frac{1}{x^s} dx \right) - \frac{1}{2}, \quad s \in \mathbb{N}$$

$\gamma_1 = 0.077215, \gamma_2 = 0.144934, \gamma_4 = 0.24899, \gamma_8 = 0.36122, \gamma_{16} = 0.433349, \dots, \gamma_{\infty} = 0.5$

$\gamma_c, \gamma_1, \gamma_2, \gamma_3, \dots$ are called Chen's natural group constants (analogue to Bernoulli numbers).

The following are some other formulas related to 2π -e Formula.

$$\begin{aligned}
\sqrt{2\pi} &= e^{1-\gamma_c}, \quad e = \sqrt{2\pi} e^{\gamma_c} = \sqrt{2\pi} \left(1 + \sum_{n=1}^{\infty} \frac{\gamma_c^n}{n!}\right) \\
\gamma_c &= \sum_{n=1}^{\infty} \left[\left(n + \frac{1}{2}\right) \ln\left(1 + \frac{1}{n}\right) - 1 \right] = \sum_{n=1}^{\infty} \frac{(2^{2n}-1)|B_{2n}|\pi^{2n} - 2(2n)!}{2(2n+1)!} = \frac{1}{4} - \sum_{s=1}^{\infty} \frac{\gamma_s}{s(s+1)} \\
\gamma_g &= \sum_{n=1}^{\infty} \left(n + \frac{1}{2}\right) \ln\left(1 + \frac{1}{n}\right) - \int_1^{\infty} \left(x + \frac{1}{2}\right) \ln\left(1 + \frac{1}{x}\right) dx \\
\gamma_{cg} &= \frac{1}{2} \lim_{N \rightarrow \infty} \left[\sum_{n=1}^N \frac{(2^{2n}-1)|B_{2n}|\pi^{2n}}{(2n+1)!} - \ln N \right] \\
\frac{\pi}{2} &= \left(\frac{e}{e^{\gamma_g}}\right)^2, \quad e = \sqrt{\frac{\pi}{2}} e^{\gamma_g} = \sqrt{\frac{\pi}{2}} \left(1 + \sum_{n=1}^{\infty} \frac{\gamma_g^n}{n!}\right); \quad \frac{\pi}{2} = \left(\frac{e^{\gamma/2}}{e^{\gamma_{cg}}}\right)^2, \quad \gamma = \ln \frac{\pi}{2} + 2\gamma_{cg} \\
\gamma_c &= \gamma_g - \ln 2 = 1 - \frac{\gamma}{2} + \gamma_{cg} - \ln 2, \quad \gamma_{cg} = \frac{1}{2} + \sum_{s=2}^{\infty} \frac{\gamma_s}{s(s+1)} - \ln 2 \\
\gamma_c &= 0.0810614668, \quad \gamma_g = 0.7742086474, \quad \gamma_{cg} = 0.0628164798 \\
\frac{\pi}{2} &= \sum_{n=1}^{\infty} \frac{|B_{2n}|\pi^{2n}}{2n(2n)!}; \quad \sum_{n=1}^{\infty} [\zeta(2n) - 1] = \frac{3}{4}, \quad \zeta(2n) = \sum_{k=1}^{\infty} \frac{1}{k^{2n}} \\
\sum_{n=1}^{\infty} \frac{1}{n} &= \sum_{n=1}^{\infty} \frac{|B_{2n}|(2\pi)^{2n}}{2n(2n)!} = \sum_{n=1}^{\infty} \frac{|B_{2n}|(2n2^{2n}+1)\pi^{2n}}{2n(2n+1)!}
\end{aligned}$$

4. Some Applications of 2π -e Formula and its Related Formulas

(1). 2π -e formula is basically an algebraic expanding of Stirling formula, but it is more meaningful, it exhibits the relationship between 2π and e. In 2π -e formula, γ_c is a real constant with geometric definition like Euler-Mascheroni constant γ . With 2π -e formula and its related formulas, 2π can be calculated from e and vice versa. So it is the real 2π -e relationship formula.

$$2\pi = \left(\frac{e}{e^{\gamma_c}}\right)^2 = e^2 \prod_{n=1}^{\infty} \frac{e^2}{\left(1 + \frac{1}{n}\right)^{2n+1}} = e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \frac{e^2}{\left(\frac{4}{3}\right)^7} \dots$$

$$e = \sqrt{2\pi} e^{\gamma_c} = \sqrt{2\pi} \left(1 + \sum_{n=1}^{\infty} \frac{\gamma_c^n}{n!}\right), \quad \gamma_c = \sum_{n=1}^{\infty} \frac{(2^{2n}-1)|B_{2n}|\pi^{2n} - 2(2n)!}{2(2n+1)!}$$

(2). 2π -e formula demonstrates 2π is a natural constant rather than π . $\pi/2$ is somewhat fundamental but not as complete as 2π . π is neither fundamental nor complete. In 2001 mathematician Bob Palais said “ π is wrong”⁵. 2π -e formula and the Taylor expansion of e have similar form (natural group form), this should give a conclusive proof that 2π is a real natural constant and π is not.

$$2\pi = \left(\frac{e}{e^{\gamma_c}}\right)^2 = e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \frac{e^2}{\left(\frac{4}{3}\right)^7} \dots \Rightarrow 2\pi \text{ or } \sqrt{2\pi} \text{ is a natural constant}$$

$$\frac{\pi}{2} = \left(\frac{e}{e^{\gamma_g}}\right)^2 = \left(\frac{e^{\gamma/2}}{e^{\gamma_{cg}}}\right)^2 \Rightarrow \frac{\pi}{2} \text{ or } \sqrt{\frac{\pi}{2}} \text{ is almost a natural constant}$$

$$\pi = \left(\frac{e}{e^{\gamma_c} \sqrt{2}}\right)^2 = \left(\frac{e\sqrt{2}}{e^{\gamma_g}}\right)^2 \Rightarrow \pi \text{ or } \sqrt{\pi} \text{ is not a natural constant}$$

Table 1 lists some points of view of Piist who support π is a natural constant, Tauist who support 2π is a natural constant and this work which supports the later.

Table 1. Comparison of points of view of Piist, Tauist and this work.

	Piist	Tauist	This work
Circumference of a circle	πd	$2\pi R$	$2\pi R$
Area of a circle	πR^2		$(1/2)(2\pi R)R$
Volume of sphere	$(4/3)\pi R^3$	$(2/3)(2\pi)R^3$	$(2\pi R^2/3)2R$
Volume of n-dimension sphere	$\frac{\pi^{n/2}}{\Gamma(n/2+1)}R^n$	$\frac{(2\pi)^{n/2}}{2^{n/2}\Gamma(n/2+1)}R^n$	$\frac{2\pi R^2}{n}V_{n-2}$
Euler's identity	$e^{i\pi}+1=0$	$e^{2\pi i}=1$	$e^{2\pi i}=1$
Gauss integral	$\int_{-\infty}^{+\infty} e^{-x^2} dx = \sqrt{\pi}$		$\int_{-\infty}^{+\infty} e^{-x^2} dx = \frac{e}{e^{\gamma_c}} \frac{1}{\sqrt{2}}$

(3). As 2π is a square number, the frequent appearing of its square root in some

important equations such as Gaussian distribution (normal distribution) and Maxwell-Boltzmann distribution becomes reasonable and understandable. And the distributions can be transformed as follows.

$$\text{Standard Normal Distribution: } f(x, 0, 1) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} = e^{-\frac{x^2+2(1-\gamma_e)}{2}}$$

$$\text{Maxwell-Boltzmann Distribution: } f(v) = \frac{2}{\sqrt{2\pi}} v^2 \left(\frac{m}{kT}\right)^{\frac{3}{2}} e^{-\frac{mv^2}{2kT}} = 2\left(\frac{m}{kT}\right)^{\frac{3}{2}} v^2 e^{-\left(\frac{m}{kT} \frac{v^2}{2} + 1 - \gamma_e\right)}$$

(4). Euler's identity (Euler's equation) $e^{i\pi} + 1 = 0$ is called God formula and the most beautiful formula in mathematics. However, as 2π is the real natural constant and π is not, $e^{2\pi i} = 1$ should be more beautiful.

(5). $\gamma = \ln(2\pi) + \gamma_{cg}$ may help to prove γ is an irrational number or even a transcendental number.

(6). The natural group formulas help us to establish "Chen's Periodic Table of Elements and Natural Group Theory"⁶ (2014-2017).

(7). The mathematic expression of chirality is $\pm 2\pi$. This concept is helpful for us to establish "Chirality and Poetry Model of Atomic Nuclei"⁷ (2017/12-2018/3).

(8). Based on the above theories, Chen's theory of the fine-structure constant was deduced (2018/4-6)⁸ and has been revised, modified and improved (2018/7-2020/1).

5. Original Inspiration for Formulas of the Fine-structure Constant

1. According to $\alpha = \frac{e^2}{4\pi\epsilon_0\hbar c} = \frac{\lambda_e}{2\pi a_0} = \frac{2\pi r_e}{\lambda_e} \approx \frac{1}{137.036}$, the formulas of α should relate to 2π .

$$2. \frac{137.036}{2\pi} = \frac{137.036}{6.28318} = 21.81, \quad 137.036 = 21.81 \times 2\pi$$

$$3. \text{According to } 2\pi\text{-e formula: } 2\pi = \left(\frac{e}{e^{\gamma_e}}\right)^2 = e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \frac{e^2}{\left(\frac{4}{3}\right)^7} \dots$$

2π is a square number, suppose $21.81 = x^2$, $x = 4.670 \approx 14/3$

$$\text{so: } \frac{1}{\alpha} \approx \left(\frac{14}{3}\right)^2 2\pi \text{ or } \alpha \approx \left(\frac{3}{14}\right)^2 \frac{1}{2\pi} \quad (\text{Discover: about 2 am on 2018/4/12})$$

4. Apply with $2\pi\text{-e}$ formula (in the afternoon of 2018/4/12, a meeting in the morning)

$$\alpha = \left(\frac{3}{14}\right)^2 \frac{1}{(2\pi)_{112}} = \left(\frac{3}{14}\right)^2 \frac{1}{e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{113}{112}\right)^{225}}} = 137.035781520, \text{ closest to the real value.}$$

As 112 is one of the most important stable numbers and the 112th element ${}_{112}^{285}\text{Cn}_{173}^*$ is the natural end of elements according to our Chen's Chirality and Poetry Model of Atomic Nuclei⁶.

$$\text{So: } \textit{Eureka!} \quad \text{Subsequently transformed to: } \alpha = \frac{6^2}{7(2\pi)_{112}} \frac{1}{112} = 137.035781520,$$

$$\text{Finally modified to: } \alpha = \frac{6^2}{7(2\pi)_{112}} \frac{1}{112 + \frac{1}{75^2}} = 137.035999037435$$

6. Logical Deduction of Chen's Formulas of the Fine-structure Constant

Physicist Richard Feynman noticed a hydrogen-like atom with Z protons and only one electron, according to Bohr model, the line velocity of the nth rank electron $v_{e/z/n}$ satisfies:

$$\frac{v_{e/z/n}}{c} = \frac{Ze^2}{n4\pi\epsilon_0\hbar c} = \frac{Z}{n}\alpha, \text{ as } v_{e/z/n} \leq c, \alpha = \frac{v_{e/z/n}}{c} \frac{n}{Z} \approx \frac{1}{Z_{\max\text{-ideal}}} = \frac{1}{Fy} = \frac{1}{137}$$

The 137th hydrogen-like element Fy (Feynmanium) is an ideal (imaginative) element,

in reality, the above formula should be modified to: $\alpha = f(Z_{\text{real}}) \frac{1}{Z_{\max\text{-real}}}$

According to Chen's Chirality and Peotry Model of Atomic Nuclei⁶,

$$Z_{\max\text{-real}} = 112 = 2 \cdot 56, \text{ so } \alpha = f(Z_{\text{real}}) \frac{1}{Z_{\max\text{-real}}} = f(Z_{\text{real}}) \frac{1}{112}$$

Compared to $\alpha = \frac{\lambda_e}{2\pi a_0}$, the formula should have a 2π factor:

$$\alpha = f(Z_{\text{real}}) \frac{1}{Z_{\max\text{-real}}} = \frac{n}{m(2\pi)_k} \frac{1}{Z_{\max\text{-real}}} = \frac{6^2}{7 \cdot (2\pi) 112} = 1/136.8$$

Apply with 2π -e formula: $2\pi = e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \frac{e^2}{\left(\frac{4}{3}\right)^7} \dots$

the formula is transformed to:

$$\alpha = \frac{n}{m(2\pi)_k} \frac{1}{Z_{\max\text{-real}}} = \frac{6^2}{7 \cdot (2\pi)_{112}} \frac{1}{112} = \frac{6^2}{7 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{113}{112}\right)^{225}}} \frac{1}{112} = 1/137.035782$$

Above deduction on 2018/4/12, only $(2\pi)_{112}$ gives the closest value to α , this coincidence of one part per infinity proves the formula itself is correct.

Added an calibration factor ($\delta=1/75^2$) on 2018/4/20, the accurate formula is:

$$\alpha_1 = \frac{\lambda_e}{2\pi a_0} = \frac{6^2}{7 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{113}{112}\right)^{225}}} \frac{1}{112 + \frac{1}{75^2}}$$

Discover: 2018/4/12; Revise: 2018/4/20 (add $1/75^2$ factor)

By the same procedure but compared to $\alpha = \frac{2\pi r_e}{\lambda_e}$, the other formula is:

$$\alpha_2 = \frac{2\pi r_e}{\lambda_e} = \frac{m(2\pi)_k}{n} \frac{1}{Z_{\max\text{-real}}} = \frac{13 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{279}{278}\right)^{557}}}{10^2} \frac{1}{112 - \frac{1}{3 \cdot 29 \cdot 64}} = 1/137.035999111818$$

Discover: 2018/4/24; Revise: 2018/9/18-20 ($280 \rightarrow 278$, $-\frac{1}{39^2} + \frac{1}{780^2} \rightarrow -\frac{1}{3 \cdot 29 \cdot 64}$)

Another amazing coincidence is 6^2 and 10^2 are square numbers in accordance with $2\pi = \left(\frac{e}{e^{\gamma c}}\right)^2$

This also demonstrates that α has two values with two kinds of formulas.

As $f(Z_{\text{real}}) = \frac{n}{m(2\pi)_k}$ or $f(Z_{\text{real}}) = \frac{m(2\pi)_k}{n}$, m n k δ should relate to nucleon numbers of nuclides.

7. The Two Most Important Formulas

The above two formulas for α_1 and α_2 were our first gained formulas and are the most important formulas among their serial formulas which will be given followed in this paper. Calculation to give the values of α_1 and α_2 is shown in **Fig. 3** and **Table 2**.

Fig. 3. Calculation diagram of α_1 and α_2 (2018/4-6).




Table 2. Calculation of α_1 and α_2 (2018/4-6).

k	$(2\pi)_k$	$1/\alpha_1$	k	$(2\pi)_k$	$1/\alpha_2$
	7.389056099	160.917477134		7.389056099	116.596364743
1	6.824768754	148.628533230	1	6.824768754	126.236816375
2	6.640803185	144.622165589	2	6.640803185	129.733867427
3	6.549956514	142.643723845	3	6.549956514	131.533251879
4	6.49586908	141.465817857	4	6.49586908	132.628454999
5	6.46000004	140.684668634	5	6.46000004	133.364872233
6	6.434476503	140.128821836	6	6.434476503	133.893888578
7	6.415388754	139.713132398	7	6.415388754	134.292263980
8	6.400576029	139.390543654	8	6.400576029	134.603053878
9	6.388747203	139.132937708	9	6.388747203	134.852272701
10	6.379083388	138.922480953	10	6.379083388	135.056563407
14	6.353377324	138.362659116	14	6.353377324	135.603008624
28	6.319398093	137.622665802	28	6.319398093	136.332142298
56	6.301583891	137.234711452	56	6.301583891	136.717545138
110	6.29262658	137.039640822	112	6.292459356	136.915795771
111	6.292542221	137.037803660	224	6.28784124	137.016353814
112	6.292459356	137.035999037435	276	6.286966940	137.035408057
113	6.292377945	137.034226098	277	6.286953333	137.035704647
114	6.292297952	137.032484014	278	6.286939823	137.035999111818
			279	6.286926410	137.036291474
			280	6.286913093	137.036581756


In these two formulas (deduced from the modification of Z_{\max}), there are some factors which are essentially related to nucleon numbers of some nuclides especially some important stable numbers (stipulated by Chen's Chirality and Poetry Model of Atomic Nuclei⁷) such as 28, 42, 56, 83, 84, 112, 126, 166, 167, 168 *et al.* And these numbers correlate with each others. This kind of relationship is shown in the follows.

A brief illustration of the relationships between the fine-structure constant and nuclides:




Above nuclides indicate that 136–138, which can be called the fine-structure constant numbers, definitely relate to 112 and 166–168 (double of 56 and 83–84, the most stable numbers in nuclides).

$$\alpha_1 = \frac{6^2}{7 \cdot (2\pi)_{112}} \frac{1}{112 + \frac{1}{75^2}} = \frac{6^2}{7 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{113}{112}\right)^{225}}} \frac{1}{112 + \frac{1}{75^2}} = 1/137.035999037435$$




$$\alpha_2 = \frac{13 \cdot (2\pi)_{278}}{10^2} \frac{1}{112 - \frac{1}{3 \cdot 29 \cdot 64}} = \frac{13 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{9 \cdot 31}{278}\right)^{557}}}{10^2} \frac{1}{112 - \frac{1}{3 \cdot 29 \cdot 64}} = 1/137.035999111818$$



The value of the front part of each above formula is almost equal to $1/(3/2)^{1/2}$ (because 112 is the element natural proton end and 168 is the element natural neutron end as shown in $^{112}\text{Cn}_{168+5}$), so the formulas can be transformed to the follows.


$$\alpha_1 = \alpha_{1-(3/2)} = \frac{1}{\left(\frac{3}{2} - \frac{1}{3 \cdot 112 + 1} + \frac{1}{2^2 \cdot 3 \cdot 5^3 \cdot 13 \cdot 23 - \frac{30}{64}}\right)^{1/2}} \frac{1}{112 + \frac{1}{75^2}} = 1/137.035999037435$$

2019/4/25 Relations to nuclides :



$$\alpha_2 = \alpha_{2-(3/2)} = \frac{1}{\left(\frac{3}{2} - \frac{1}{3 \cdot 112 + 1} + \frac{1}{2 \cdot 7 \cdot 11 \cdot 19 \cdot 29 + \frac{36}{75^2}}\right)^{1/2}} \frac{1}{112 - \frac{1}{3 \cdot 29 \cdot 64}} = 1/137.035999111818$$

2019/4/25 Relations to nuclides:



8. The Integrated Fine-structure Constant

Multiplication of α_1 and α_2 should almost divide out the 2π factors and give $3/2$ and 112×112 factors, this means $\alpha_1 \alpha_2$ is almost equal to 112×168 , so we define $\alpha_c = (\alpha_1 \alpha_2)^{1/2}$ as the integrated fine-structure constant or Chen's fine-structure constant.

$$\begin{aligned} \frac{1}{\alpha_c^2} &= \frac{1}{\alpha_1 \alpha_2} = \frac{2\pi a_0}{\lambda_e} \frac{\lambda_e}{2\pi r_e} = \frac{a_0}{r_e} = \left(\frac{c}{v_e}\right)^2 \\ &= 112 \times \left(168 - \frac{1}{3} + \frac{1}{12 \cdot 47} - \frac{1}{6 \cdot 29 \cdot 53 \cdot 59 - 79/47}\right) \quad 2018/6/8-9, 9/18-19, 2019/4/19 \\ &= 136 \left(138 + \frac{1}{2} - \frac{1}{10 \cdot 29} + \frac{1}{12 \cdot 53 \cdot (6 \cdot 53 - 1) - 27/47}\right) \quad 2019/4/17-19 \\ &= 137 \left(137 + \frac{1}{13} - \frac{1}{7 \cdot 29} + \frac{1}{32 \cdot 33 \cdot 89 + 16/49}\right) \quad 2019/4/17-19 \\ &= 112 \cdot 167.668437878408 = 18778.865042381 \\ & \begin{matrix} 27 & 29 & 47,49 & 53 & 54,56,58 & 59 & 58,60,61 & 63,65 & 79 & 87 \\ 13 & Al_{14} & 14 & Si_{15} & 22 & Ti_{25,27} & 24 & Cr_{29} & 26 & Fe_{28,30,32} & 27 & Co_{32} & 28 & Ni_{30,32,33} & 29 & Cu_{34,36} & 35 & Br_{44} & 38 & Sr_{49} \end{matrix} \\ & \begin{matrix} 100,102 & 112 & 113 & 135-138 & 136,138 & 3-47 & 158,160 & 159 & 166,168 \\ 44 & Ru_{56,58} & 48 & Cd_{64} & 49 & In_{64} & 56 & Ba_{79-82} & 58 & Ce_{78,80} & 82 & Pr_{82} & 64 & Gd_{94,96} & 65 & Tb_{94} & 68 & Er_{98,100} \end{matrix} \\ & \begin{matrix} 174 & 188 & 197 & 203 & 223 & 226 & 227 & 262 & 285 & 293 \\ 70 & Yb_{104} & 76 & Os_{112} & 79 & Au_{118} & 81 & Tl_{122} & 87 & Fr_{136}^* & 88 & Ra_{138}^* & 89 & Ar_{138}^* & 103 & Lr_{159}^* & 112 & Cn_{173}^* & 116 & Lv_{177}^{ie} \end{matrix} \\ \alpha_c^2 &= \alpha_1 \alpha_2 = \left[\frac{6^2}{7 \cdot (2\pi)_{112}} \frac{1}{112 + \frac{1}{75^2}} \right] \left[\frac{13 \cdot (2\pi)_{278}}{10^2} \frac{1}{112 - \frac{1}{3 \cdot 29 \cdot 64}} \right] \\ &= \frac{13 \cdot 3^2}{7 \cdot 5^2} \frac{e^2}{\left(\frac{2 \cdot 3 \cdot 19}{113}\right)^{227}} \frac{e^2}{\left(\frac{115}{114}\right)^{229}} \dots \frac{e^2}{\left(\frac{9 \cdot 31}{2 \cdot 139}\right)^{557}} \frac{1}{112^2 - \frac{1}{30^2 \cdot 5} + \frac{1}{60^2 \cdot 15} - \frac{1}{120^2 \cdot 15 \cdot 29}} \\ &= 1/18778.865042381 \quad 2019/12/14 \\ & \begin{matrix} 27 & 31 & 39 & 55 & 54,56,57,58 & 63,65 & 69,71 & 79,81 & 87 \\ 13 & Al_{14} & 15 & P_{16} & 19 & K_{20} & 25 & Mn_{30} & 26 & Fe_{28,30,31,32} & 29 & Cu_{34,36} & 31 & Ga_{38,40} & 35 & Br_{44,46} & 38 & Sm_{49} \end{matrix} \\ & \begin{matrix} 89 & 93 & 112-120-124 & 135-138 & 139 & 136,138 & 144,145 & 157 \\ 39 & Y_{50} & 41 & Nb_{52} & 50 & Sn_{62-70-74} & 56 & Ba_{79-82} & 57 & La_{82} & 58 & Ce_{78,80} & 60 & Nd_{83,84} & 64 & Gd_{93} \end{matrix} \\ & \begin{matrix} 200 & 209 & 223 & 237 & 278+7 & 284 \\ 80 & Hg_{120} & 83 & Bi_{126}^* & 87 & Fr_{136}^* & 93 & Np_{144}^* & 112 & Cn_{166+7}^* & 113 & Nh_{9,19}^{ie} \end{matrix} \\ \alpha_c^2 &= \alpha_1 \alpha_2 = \frac{1}{\left(\frac{3}{2} - \frac{1}{3 \cdot 112 + 1} + \frac{1}{7 \cdot 19 \cdot 29 \cdot 37 - \frac{25}{44}}\right)} \frac{1}{112 + \frac{1}{75^2}} \frac{1}{112 - \frac{1}{3 \cdot 29 \cdot 64}} \\ &= 1/18778.865042381 \quad 2019/12/14 \\ & \begin{matrix} 39 & 47,50 & 55 & 63,65 & 85,87 & 87,88 & 99,100,102,104 & 112 \\ 19 & K_{20} & 22 & Ti_{25,28} & 25 & Mn_{30} & 29 & Cu_{34,36} & 37 & Rb_{48,50} & 38 & Sr_{49,50} & 44 & Ru_{55,56,58,60} & 48 & Cd_{64} \end{matrix} \\ & \begin{matrix} 112,114,115,116,120,124 & 5-37,11-17 & 223 & 226 \\ 50 & Sn_{62,64,65,66,70,74} & 75 & Re_{110,112} & 87 & Fr_{136}^* & 88 & Ra_{138}^* \end{matrix} \end{aligned}$$

9. A Brief Explanation of the Fine-structure Constant

According to Chen's Chirality and Poetry Model of Atomic Nuclei⁷, the ratio of neutron number N to proton number Z in nuclides increases from 1/1 to 3/2 (eventually slightly above 3/2) along with the increasing of atomic number, for example, from ${}_{14}\text{Si}_{14}$, ${}_{26}\text{Fe}_{30}$, ${}_{29}\text{Cu}_{34,36}$, ${}_{56}\text{Ba}_{82}$, ${}_{84}\text{Po}_{125}$ to ${}_{112}\text{Cn}_{168+5}^*$. In this process, $(3/2)^{1/2}$ will act as a transition foothold. As for nuclide ${}_{112}\text{Cn}_{168+5}$ with Z=112, N=168+5 and $168/112=3/2$, 137 is just right their $(3/2)^{1/2}$ times intermediate stage. This should be why 137 exists and what's the real meaning of 137.

electromagnetic wave or light, it should be reasonable to suppose the speed of light to be the integrated fine-structure constant, i.e., $c_{au}=1/\alpha_c=1/(\alpha_1\alpha_2)^{1/2}=137.035999074627$. It means we've theoretically/mathematically calculated the speed of light, the formula is intrinsically consistent with Maxwell's formula, and the value is much accurate.

In atomic units ($e = m_e = \hbar = 1$ and $\varepsilon_0 = \frac{1}{4\pi}$), $v_{e/au} = \alpha c_{au} = \frac{e^2}{4\pi\varepsilon_0\hbar} = 1$, so $c_{au} = \frac{1}{\alpha}$

There are two α (α_1 and α_2), but there shouldn't be two c or c_{au} ,

so it should be: $c_{au} = \frac{1}{\alpha_c} = \frac{1}{\sqrt{\alpha_1\alpha_2}}$ (au : atomic units)

Compared to Maxwell Formula $c = \frac{1}{\sqrt{\mu_0\varepsilon_0}}$, $c_{au} = \frac{1}{\alpha_c} = \frac{1}{\sqrt{\alpha_1\alpha_2}}$ should be reasonable.

$$c_{au} = \frac{1}{\sqrt{\mu_{0/au}\varepsilon_{0/au}}}, \mu_{0/au}\varepsilon_{0/au} = \alpha_1\alpha_2, \mu_{0/au} = 4\pi\alpha_1\alpha_2 \quad (2019/11/30)$$

So the theoretical formula of the speed of light in atomic units is as follows:

$$\begin{aligned} c_{au} &= \frac{1}{\alpha_c} = \frac{1}{\sqrt{\alpha_1\alpha_2}} = \frac{1}{\sqrt{\left(\frac{6^2}{7 \cdot (2\pi)_{112}} \frac{1}{112 + \frac{1}{75^2}}\right) \left(\frac{13 \cdot (2\pi)_{278}}{10^2} \frac{1}{112 - \frac{1}{3 \cdot 29 \cdot 64}}\right)}} \\ &= \frac{5}{3} \sqrt{\frac{7 \cdot (2\pi)_{112}}{13 \cdot (2\pi)_{278}} \left(112^2 - \frac{1}{30^2 \cdot 5} + \frac{1}{60^2 \cdot 15} - \frac{1}{120^2 \cdot 15 \cdot 29}\right)} \\ &= \sqrt{\frac{5 \cdot 17 - \frac{10}{11 \cdot 11 \cdot 23}}{7 - \frac{1}{7 \cdot 19}} \frac{(2\pi)_{12389}}{(2\pi)_{28186}} \left(112^2 - \frac{2 \cdot 7^2 \cdot 43 \cdot 67 + \frac{5}{7}}{5^2 \times 10^{10}}\right)} \\ &= \frac{5}{3} \sqrt{\left(\frac{2^3 \cdot 17}{11 \cdot 23} + \frac{2 \cdot 17}{11 \cdot 23 \cdot 97}\right) \frac{(2\pi)_{34450}}{(2\pi)_{28186}} \left(112^2 - \frac{2^5 \cdot 3 \cdot 7 \cdot 13}{10^{10}}\right)} \\ &= \sqrt{\left(\frac{3}{2} - \frac{1}{3 \cdot 112 + 1} + \frac{1}{7 \cdot 19 \cdot 29 \cdot 37 - \frac{25}{44}}\right) \left(112^2 - \frac{1}{30^2 \cdot 5} + \frac{1}{60^2 \cdot 15} - \frac{1}{120^2 \cdot 15 \cdot 29}\right)} \\ &= \sqrt{\frac{3}{2} \left(112 - \frac{1}{3^2} + \frac{1}{12^2 \cdot 13 - \frac{30 \cdot 19}{100} - \frac{1}{125 \cdot 100}}\right)} = \sqrt{\frac{3}{2} - \frac{1}{3 \cdot 112 + 1} + \frac{1}{14 \cdot 53 \cdot 193 - \frac{33}{2 \cdot 47}}} \times 112 \\ &= \sqrt{112 \times \left(168 - \frac{1}{3} + \frac{1}{12 \cdot 47} - \frac{1}{6 \cdot 29 \cdot 53 \cdot 59 - 79/47}\right)} \\ &= \sqrt{137.035999037435 \times 137.035999111818} = 137.035999074627 \end{aligned}$$

Note: $112/278 \approx 27/67$, $12389/28186 \approx 11/25$, $34450/28186 \approx 11/9 \approx 66/29$

Discover: 2019/12/16; Revise and Supplement: 2020/1/5-8, 2/24

12. The Special 29 and 75 Factors

In the above formulas some factors especially 29 and 75 appear several times. This feature should be analyzed and explained. Accompanying N/Z ratio from 1/1 to slightly above 3/2 along with the increasing of atomic number, ${}_{29}\text{Cu}_{34,36}$ is the critical point of N/Z ratio approaching $(3/2)^{1/2}$ and ${}_{75}\text{Re}_{110,112}$ is the critical point of N/Z ratio approaching 3/2 (**Table 3**, **Fig. 4** and **Fig. 5**), so 29 and 75 are important factors and hence frequently appear in the formulas.

Table 3. N/Z ratios of the Elements (2019/4/23).

Z	N	N/Z	Z	N	N/Z	Z	N	N/Z	Z	N	N/Z				
H	1	0	0	Ga	31	38.80	1.25	Pm	61	84	1.38	Pa*	91	140	1.54
He	2	2.00	1.00	Ge	32	40.71	1.27	Sm	62	88.45	1.43	U*	92	146	1.59
Li	3	3.92	1.31	As	33	42	1.27	Eu	63	89.04	1.41	Np*	93	144	1.55
Be	4	5	1.25	Se	34	45.05	1.33	Gd	64	93.33	1.46	Pu*	94	150	1.60
B	5	5.80	1.16	Br	35	44.98	1.29	Tb	65	94	1.45	Am*	95	148	1.56
C	6	6.01	1.00	Kr	36	47.89	1.33	Dy	66	96.57	1.46	Cm*	96	151	1.57
N	7	7.00	1.00	Rb	37	48.56	1.31	Ho	67	98	1.46	Bk*	97	150	1.55
O	8	8.00	1.00	Sr	38	49.71	1.31	Er	68	99.33	1.46	Cf*	98	153	1.56
F	9	10	1.11	Y	39	50	1.28	Tm	69	100	1.45	Es*	99	153	1.55
Ne	10	10.19	1.02	Zr	40	51.32	1.28	Yb	70	103.11	1.47	Fm*	100	157	1.57
Na	11	12	1.09	Nb	41	52	1.27	Lu	71	104.03	1.47	Md*	101	157	1.55
Mg	12	12.32	1.03	Mo	42	54.04	1.29	Hf	72	106.54	1.48	No*	102	157	1.54
Al	13	14	1.08	Td	43	55	1.28	Ta	73	108	1.48	Lr*	103	159	1.54
Si	14	14.11	1.01	Ru	44	57.16	1.30	W	74	109.89	1.49	Rf*	104	161	1.55
P	15	16	1.07	Rh	45	58	1.29	Re	75	111.25	1.48	Db*	105	163	1.55
S	16	16.09	1.01	Pd	46	60.51	1.32	Os	76	114.27	1.50	Sg*	106	165	1.56
Cl	17	18.48	1.09	Ag	47	60.96	1.30	Ir	77	115.25	1.50	Bh*	107	163	1.52
Ar	18	21.99	1.22	Cd	48	64.52	1.34	Pt	78	117.12	1.50	Hs*	108	169	1.56
K	19	20.13	1.06	In	49	65.91	1.35	Au	79	118	1.49	Mt*	109	167	1.53
Ca	20	20.12	1.01	Sn	50	68.81	1.38	Hg	80	120.62	1.51	Ds*	110	171	1.55
Sc	21	24	1.14	Sb	51	70.86	1.39	Tl	81	123.41	1.52	Rg*	111	169	1.52
Ti	22	25.92	1.18	Te	52	75.70	1.46	Pb	82	125.24	1.53	Cn*	112	173	1.54
V	23	28	1.22	I	53	74	1.40	Bi*	83	126	1.52	Nh*	113	171	1.51
Cr	24	28.06	1.17	Xe	54	77.39	1.43	Po*	84	125	1.49	Fl*	114	175	1.54
Mn	25	30	1.20	Cs	55	78	1.42	At*	85	125	1.47	Mc*	115	173	1.50
Fe	26	29.91	1.15	Ba	56	81.42	1.45	Rn*	86	136	1.58	Lv*	116	177	1.53
Co	27	32	1.19	La	57	82	1.44	Fr*	87	136	1.56	Ts*	117	177	1.51
Ni	28	30.76	1.10	Ce	58	82.21	1.42	Ra*	88	138	1.57	Og*	118	176	1.49
Cu	29	34.62	1.19	Pr	59	82	1.39	Ac*	89	138	1.55				
Zn	30	35.45	1.18	Nd	60	84.41	1.41	Th*	90	142	1.58				

Z: atomic number, N: average neutron number or neutron number of the most stable isotope.

1. N/Z from 1/1 (${}_6\text{C}$) to slightly above 3/2 (such as ${}_{112}\text{Cn}$ which is the natural end of elements demonstrated by Chen's Chirality and Poetry Model of Atomic Nuclei⁷).
2. For ${}_{29}\text{Cu}$, N/Z ratio 1.19 is near to $(3/2)^{1/2}=1.22$, slightly less is because of stability effect.
3. For ${}_{75}\text{Re}$, N/Z ratio 1.48 is near to $3/2=1.50$, slightly less is because of stability effect.
4. From ${}_6\text{C}$ to ${}_{112}\text{Cn}$, the middle of N/Z 1.5 range is at $(76.5-5)/(112-5)=0.668 \approx 2/3$ position.

Fig. 4 and **Fig. 5** shows that stability effect of nucleon number 64 makes the neutron numbers of ${}_{29}\text{Cu}$'s isotopes are relatively less (34 and 36) than normal so that its N/Z ratio is a little less than $(3/2)^{1/2}$ which is otherwise it should be. Also the

stability effect of nucleon numbers 110 and 112 make the neutron numbers of ^{75}Re 's nuclides are relatively less (110 and 112) than normal so that its N/Z ratio is a little less than $3/2$ which otherwise it should be.

Fig. 4. Complete Graph of N/Z Ratios of Elements (2019/4/23-24).





Fig. 5. Partially Amplified Graph of N/Z Ratios of Elements (2019/4/24).




The general trend of N/Z ratio of elements is from 1/1 (${}^6\text{C}_6$) to slightly above 3/2 (${}^{112}\text{Cn}_{173}$) definitely. However, the increasing process is not smooth, the N/Z ratio rising fluctuates consecutively. According to Chen's Chirality and Poetry Model of Atomic Nuclei⁷, there are some stable numbers (magic numbers) which can bring about this kind of fluctuation (**Table 4** and **Fig. 6**).

Table 4. Effect of Stable Numbers on N/Z ratio's fluctuation (2019/4/22).

Element	Z	N(Average)	Z(3/2) ^{1/2}	N-Z(3/2) ^{1/2}	Stable Number
K	19	20.13	23.27	-3.17	20
Ca	20	20.12	24.49	-4.41	20+20
Sc	21	24	25.72	-1.74	
Ti	22	25.92	26.94	-1.07	22+26=48
V	23	28.00	28.17	-0.23	28
Cr	24	28.06	29.39	-1.39	28
Mn	25	30	30.62	-0.68	
Fe	26	29.91	31.84	-1.99	26+30=56
Co	27	32.00	33.07	-1.14	
Ni	28	30.76	34.29	-3.60	28+30=58、28+32=60
Cu	29	34.62	35.52	-0.97	64
Zn	30	35.45	36.74	-1.36	30+34=64、30+36=66
Ga	31	38.80	37.97	0.75	
Ge	32	40.71	39.19	1.44	32+40=72
As	33	42.00	40.42	1.50	
Se	34	45.05	41.64	3.30	34+46=80
Br	35	44.98	42.87	2.03	
Kr	36	47.89	44.09	3.71	36+48=84

Fig. 6. Effect of Stable Numbers on N/Z ratio's fluctuation (2019/4/22-23)



13. α_1/α_2 in Schrödinger Equation of Hydrogen Atom

Stationary Schrodinger Equation $-\frac{\hbar^2}{2m}\nabla^2\psi + U\psi = E\psi$, applied to hydron atom:

$$\nabla^2\psi + \frac{2m_e}{\hbar^2}(E + \frac{e^2}{4\pi\epsilon_0 r})\psi = 0, \quad E = -\frac{m_e e^4}{2n^2(4\pi\epsilon_0)^2\hbar^2}, \text{ do substitution and simplification:}$$

$$\frac{2m_e}{\hbar^2}(\frac{m_e e^4}{2n^2(4\pi\epsilon_0)^2\hbar^2} - \frac{e^2}{4\pi\epsilon_0 r})\psi = \nabla^2\psi, \quad [\frac{1}{n^2}(\frac{m_e e^2}{4\pi\epsilon_0\hbar^2})^2 - \frac{2}{r} \frac{m_e e^2}{4\pi\epsilon_0\hbar^2}]\psi = \nabla^2\psi,$$

$$[\frac{1}{n^2}(\frac{e^2}{4\pi\epsilon_0\hbar c} \frac{m_e c}{\hbar})^2 - \frac{2}{r} \frac{e^2}{4\pi\epsilon_0\hbar c} \frac{m_e c}{\hbar}]\psi = \nabla^2\psi,$$

$$\text{As } \sqrt{\alpha_1\alpha_2} = \frac{v_e}{c} = \frac{e^2}{4\pi\epsilon_0\hbar c}, \lambda_e = \frac{h}{m_e c} \text{ and } \alpha_1 = \frac{\lambda_e}{2\pi a_0}:$$

$$[\frac{1}{n^2}(\sqrt{\alpha_1\alpha_2} \frac{2\pi}{\lambda_e})^2 - \frac{2}{r} \sqrt{\alpha_1\alpha_2} \frac{2\pi}{\lambda_e}]\psi = \nabla^2\psi,$$

$$[\frac{1}{n^2(\lambda_e/2\pi/\sqrt{\alpha_1\alpha_2})^2} - \frac{2}{(\lambda_e/2\pi/\sqrt{\alpha_1\alpha_2})r}]\psi = \nabla^2\psi,$$

$$[\frac{1}{n^2 a_0^2 (\alpha_1/\alpha_2)} - \frac{2}{a_0 r \sqrt{\alpha_1/\alpha_2}}]\psi = \nabla^2\psi$$

$$\text{As } \alpha_1/\alpha_2 \approx 1, \text{ simplified to: } [\frac{1}{n^2 a_0^2} - \frac{2}{a_0 r}]\psi = \nabla^2\psi \text{ (factor 2 seems not beautiful)}$$

In atomic units (*au*: $e = m_e = \hbar = 1$ and $\epsilon_0 = \frac{1}{4\pi}$),

$$a_{0/au} = \frac{4\pi\epsilon_0\hbar^2}{m_e e^2} = 1, \quad v_{e/au} = \frac{e^2}{4\pi\epsilon_0\hbar} = 1, \quad c_{au} = \frac{v_{e/au}}{\alpha_c} = \frac{1}{\alpha_c} = \frac{1}{\sqrt{\alpha_1\alpha_2}}$$

$$[\frac{1}{n^2(\alpha_1/\alpha_2)} - \frac{2}{r_{au}\sqrt{\alpha_1/\alpha_2}}]\psi = \nabla_{au}^2\psi, \text{ or } (\frac{c_{au}^2}{\alpha_1^2 n^2} - \frac{2c_{au}}{\alpha_1 r_{au}})\psi = \nabla_{au}^2\psi$$

the above equation could be called Schrodinger-Chen equation of hydrogen atom, the later form of the equation shows factor 2 is still reasonable and beautiful.

$$\text{As } \alpha_1/\alpha_2 \approx 1, \text{ simplified to: } [\frac{1}{n^2} - \frac{2}{r_{au}}]\psi = \nabla_{au}^2\psi$$

Discover: 2018/4-6; Revise: 2019/12/13 (add *au* form)

$$\alpha_1/\alpha_2 = \frac{137.035999111818}{137.035999037435} = 1.0000000005428 = 1 + \frac{23 \cdot 59}{25 \cdot 10^{11}} = (1 + \frac{23 \cdot 59}{50 \cdot 10^{11}})^2$$

$$\sqrt{\alpha_1/\alpha_2} = 1 + \frac{23 \cdot 59}{50 \cdot 10^{11}} = 1.0000000002714$$

Relations to nuclides: ${}_{11}^{23}\text{Na}_{12}$ ${}_{23}^{50,51}\text{V}_{27,28}$ ${}_{25}^{55}\text{Mn}_{30}$ ${}_{44}^{99,100}\text{Ru}_{55,56}$ ${}_{46}^{105}\text{Pd}_{59}$ ${}_{56}^{137}\text{Ba}_{81}$
 ${}_{50}^{118+1}\text{Sn}_{69}$ ${}_{59}^{141}\text{Pr}_{82}$ ${}_{69}^{169}\text{Tm}_{100}$ ${}_{75}^{185,187}\text{Re}_{110,112}$ ${}_{88}^{169}\text{Ra}^*_{137}$

2019/8/28-29

Solution of Schrödinger equation of hydrogen atom gives some quantum numbers such as n , l and m_l which determine the electron shell structure and the chemical

properties of atoms. That means Schrödinger equation of hydrogen atom is the base of chemical periodicity of elements. On the other hand, from above analysis, we have already demonstrated the formulas of the fine-structure constant α are derived from Chen's Chirality and Poetry Model of Atomic Nuclei⁷ and hence mainly connected to the stability of atomic nuclei. So, a question is whether and how α is connected to Schrödinger Equation of hydrogen atom. This question should reveal the connection of the theory of electron shell of atoms and the theory of nuclei of elements. The above deduction provides the answer. The fine-structure constant α relates to Schrödinger Equation of hydrogen atom in α_1/α_2 way which is subtle and negligible but could show the equation is really reasonable and beautiful.

14. The Two Kinds of General Formulas of the Fine-structure Constant

Based on the above two formulas of α_1 and α_2 , it should be reasonable to assume there are two kinds of serial formulas of α_1 and α_2 which are listed in follows. Among these formulas, the above two first discovered formulas are the most fundamental and important. Some formulas both with a big m and an extra large k should be more important referring to the trend of the approximate values of α .

Approximate formulas:

$$\alpha_{1-m'} = \frac{n}{m \cdot (2\pi)_k} \frac{1}{112} = \frac{n}{m \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{k+1}{k}\right)^{2k+1}}} \frac{1}{112} \approx 1/137.036$$

$$\alpha_{2-m'} = \frac{m \cdot (2\pi)_k}{n} \frac{1}{112} = \frac{m \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{k+1}{k}\right)^{2k+1}}}{n} \frac{1}{112} \approx 1/137.036$$

Accurate Formulas:

$$\alpha_{1-m} = \frac{n}{m \cdot (2\pi)_k} \frac{1}{112 + \delta_1} = \frac{n}{m \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{k+1}{k}\right)^{2k+1}}} \frac{1}{112}$$

$$= 1/137.035999037435$$

$$\alpha_{2-m} = \frac{m \cdot (2\pi)_k}{n} \frac{1}{112 - \delta_2} = \frac{m \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{k+1}{k}\right)^{2k+1}}}{n} \frac{1}{112 - \delta_2}$$


$$= 1/137.035999111818$$

Discover: 2019/6/27; Revise: 2019/7/2-3


Table 5. Parameters and Results of Approximate Formulas of α_1 (2019/7/2).

m	n	k	$\alpha_{1-m'}$	m	n	k	$\alpha_{1-m'}$
1	6	1	122.265854937	24	124	27	137.016359405
2	11	2	135.230901223	25	129	34	137.030171763
3	16	4	136.413250690	26	134	46	137.027100696
4	21	7	136.861626741	27	139	66	137.033636049
5	26	13	136.950569252	28	144	112	137.035781520
6	31	27	137.016359405	29	149	321	137.035917078
7	36	112	137.035781520	30	155	27	137.016359405
8	42	7	136.861626741	31	160	32	137.035453560
9	47	9	137.018237882	32	165	40	137.034309209
10	52	13	136.950569252	33	170	52	137.034083409
11	57	18	137.001388822	34	175	72	137.034617877
12	62	27	137.016359405	35	180	112	137.035781520
13	67	46	137.027100696	36	185	236	137.035810961
14	72	112	137.035781520	43	221	200	137.035845637
15	78	13	136.950569252	50	257	181	137.035307038
16	83	16	136.992590996	59	303	2645	137.035986189
17	88	20	137.028423583	81	416	1605	137.035992406
18	93	27	137.016359405	96	493	5806	137.035998789
19	98	37	137.034579883	103	529	1310	137.035994308
20	103	58	137.030572071	133	683	12389	137.035999034
21	108	112	137.035781520	140	719	1923	137.035994882
22	113	782	137.035967638	155	796	3988	137.035997989
23	119	22	137.001596764	170	873	34450	137.035999031


Fig. 7. Results of Approximate Formulas of α_1 (2019/7/2).




$$\alpha_{1-1} = \frac{6}{1 \cdot e^2 \left(\frac{2}{1}\right)^2} \frac{1}{112 + \frac{17}{2} - \frac{1}{40} + \frac{1}{6 \cdot 23 \cdot 25 - \frac{36}{55}}} = 1/137.035999037434$$




$$\alpha_{1-2} = \frac{11}{2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5}} \frac{1}{112 + \frac{3}{2} - \frac{1}{200} + \frac{1}{5 \cdot (3 \cdot 42 + 1) \cdot (6 \cdot 37 - 1) + \frac{2}{7}}} = 1/137.035999037435$$




$$\alpha_{1-3} = \frac{4^2}{3 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \frac{e^2}{\left(\frac{4}{3}\right)^7} \frac{e^2}{\left(\frac{5}{4}\right)^9}} \frac{1}{112 + 1 - \frac{1}{2} + \frac{1}{88} - \frac{1}{13 \cdot (2 \cdot 9 \cdot 5 \cdot 13 + 1) - \frac{2}{73}}} = 1/137.035999037435$$




$$\alpha_{1-4} = \frac{21}{2^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{8}{7}\right)^{15}}} \frac{1}{112 + \frac{1}{7} - \frac{1}{8 \cdot 19 \cdot 41 - \frac{75}{98}}} = 1/137.035999037435$$




$$\alpha_{1-5} = \frac{26}{5 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{14}{13}\right)^{27}}} \frac{1}{112 + \frac{1}{14} - \frac{1}{9 \cdot 71} + \frac{1}{67 \cdot (75 \cdot 100 - 1) + \frac{1}{10}}} = 1/137.035999037435$$




$$\alpha_{1-6} = \frac{31}{6 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{28}{27}\right)^{55}}} \frac{1}{112 + \frac{1}{2 \cdot 31} - \frac{1}{3 \cdot 11 \cdot 13 \cdot 31 - \frac{43}{4 \cdot 27}}} = 1/137.035999037435$$




$$\alpha_1 = \alpha_{1-7} = \frac{6^2}{7 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{113}{112}\right)^{225}}} \frac{1}{112 + \frac{1}{75^2}} = 1/137.035999037435$$




$$\alpha_{1-9} = \frac{47}{3^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{10}{9}\right)^{19}} 112 + \frac{1}{4 \cdot 17} - \frac{1}{2(8 \cdot 9 \cdot 37 - 1) + \frac{3 \cdot 17}{157}}} = 1/137.035999037436$$




$$\alpha_{1-11} = \frac{3 \cdot 19}{11 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{19}{18}\right)^{37}} 112 + \frac{1}{35} - \frac{1}{88 \cdot 41 - \frac{5 \cdot 53}{22 \cdot 13}}} = 1/137.035999037435$$




$$\alpha_{1-13} = \frac{67}{13 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{47}{2 \cdot 23}\right)^{3 \cdot 31}} 112 + \frac{1}{137} - \frac{1}{6(2 \cdot 27 \cdot 59 + 1) + \frac{9}{50}}} = 1/137.035999037435$$




$$\alpha_{1-16} = \frac{83}{4^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{17}{16}\right)^{33}} 112 + \frac{1}{28} - \frac{1}{6 \cdot (18 \cdot 41 + 1) + \frac{173}{2 \cdot (2 \cdot 75 - 1)}}} = 1/137.035999037435$$




$$\alpha_{1-17} = \frac{2^2 \cdot 22}{17 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{21}{20}\right)^{41}} 112 + \frac{1}{137} - \frac{1}{2 \cdot 19 \cdot 23 \cdot 59 - \frac{30}{100}}} = 1/137.035999037435$$



$$\alpha_{1-19} = \frac{2 \cdot 7^2}{19 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{38}{37}\right)^{75}} 112 + \frac{1}{2 \cdot (8 \cdot 54 - 1) + \frac{54}{19^2}}} = 1/137.035999037440$$




$$\alpha_{1-20} = \frac{103}{2^2 \cdot 5 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{59}{2 \cdot 29}\right)^{9 \cdot 13}} 112 + \frac{1}{32 \cdot 45 \cdot 79 + \frac{22}{3 \cdot 17}}} = 1/137.035999037435$$




$$\alpha_{1-22} = \frac{113}{22 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{27 \cdot 29}{2 \cdot 17 \cdot 23}\right)^{5 \cdot (2 \cdot 157 - 1)}}} \frac{1}{112 + \frac{1}{2 \cdot [2 \cdot 3 \cdot 17 \cdot (10 \cdot 19 + 1) + 1] + \frac{29}{49}}}$$


$$= 1/137.035999037435$$




$$\alpha_{1-23} = \frac{7 \cdot 17}{23 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{23}{22}\right)^{45}}} \frac{1}{112 + \frac{1}{35} - \frac{1}{4 \cdot 13 \cdot 43 - \frac{2 \cdot 29}{16 \cdot 17 - 1}}} = 1/137.035999037435$$




$$\alpha_{1-25} = \frac{3 \cdot 43}{5^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{35}{34}\right)^{3 \cdot 23}}} \frac{1}{11 \cdot 19 - \frac{1}{13^2(16 \cdot 17 - 1) + \frac{11}{25}}} = 1/137.035999037435$$




$$\alpha_{1-27} = \frac{139}{27 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{67}{66}\right)^{7 \cdot 19}}} \frac{1}{11 \cdot 47 + \frac{18}{23} + \frac{1}{138 \cdot 137}} = 1/137.035999037435$$




$$\alpha_{1-29} = \frac{149}{29 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{14 \cdot 23}{3 \cdot 107}\right)^{643}}} \frac{1}{6 \cdot 8 \cdot (12 \cdot 26 - 1) + \frac{11}{18}} = 1/137.035999037434$$




$$\alpha_{1-31} = \frac{4^2 \cdot 10}{31 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{33}{32}\right)^{5 \cdot 13}}} \frac{1}{12 \cdot 11 \cdot 17 - \frac{4 \cdot 49}{5 \cdot 41}} = 1/137.035999037434$$




$$\alpha_{1-32} = \frac{15 \cdot 11}{2 \cdot 4^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{41}{40}\right)^{81}}} \frac{1}{25 \cdot 29 - \frac{5 \cdot 83}{19 \cdot 23}} = 1/137.035999037435$$




$$\alpha_{1-33} = \frac{170}{33 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{53}{4 \cdot 13}\right)^{105}}} \frac{1}{112 + \frac{1}{22 \cdot 29 + \frac{4 \cdot 73}{5 \cdot 83}}} = 1/137.035999037436$$




$$\alpha_{1-34} = \frac{7 \cdot 5^2}{34 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{73}{72}\right)^{5 \cdot 29}}} \frac{1}{112 + \frac{1}{15 \cdot 59 + \frac{13}{15} + \frac{1}{3 \cdot (2 \cdot 15 \cdot 17 - 1)}}}} = 1/137.035999037435$$




$$\alpha_{1-36} = \frac{5 \cdot 37}{6^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{3 \cdot 79}{4 \cdot 59}\right)^{11 \cdot 43}}} \frac{1}{112 + \frac{1}{5 \cdot (31 \cdot 42 - 1) + \frac{3 \cdot 31}{14 \cdot 13}}} = 1/137.035999037436$$




$$\alpha_{1-43} = \frac{13 \cdot 17}{43 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{3 \cdot 67}{200}\right)^{401}}} \frac{1}{112 + \frac{1}{8 \cdot (12 \cdot 83 + 1) + \frac{4}{3 \cdot 13}}} = 1/137.035999037436$$




$$\alpha_{1-50} = \frac{2 \cdot 257}{100 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{14 \cdot 13}{181}\right)^{311^2}}} \frac{1}{112 + \frac{1}{29 \cdot 61 + \frac{157}{16 \cdot 11}}} = 1/137.035999037436$$



$$\alpha_{1-59} = \frac{3 \cdot 101}{59 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{2 \cdot 27 \cdot 49}{5 \cdot 23^2}\right)^{11 \cdot 13 \cdot 37}}} \frac{1}{112 + \frac{1}{48 \cdot 64 \cdot 31 - \frac{17}{81}}} = 1/137.035999037435$$




$$\alpha_{1-81} = \frac{4^2 \cdot 26}{9^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{22 \cdot 73}{15 \cdot 107}\right)^{13^2 \cdot 19}}} \frac{1}{112 + \frac{1}{2 \cdot 81 \cdot 17 \cdot 67 + \frac{35}{88}}} = 1/137.035999037435$$




$$\alpha_{1-96} = \frac{17 \cdot 29}{4^2 \cdot 6 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{16 \cdot 3 \cdot 11^2 - 1}{27 \cdot 5 \cdot 43 + 1}\right)^{79 \cdot 147}}} \frac{1}{112 + \frac{163 \cdot (8 \cdot 21 \cdot 37 + 1)}{50 \cdot 10^{11}}}$$

$$= 1/137.035999037435$$




$$\alpha_{1-103} = \frac{23^2}{103 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{3 \cdot 19 \cdot 23}{7 \cdot 11 \cdot 17 + 1}\right)^{2621}}} \frac{1}{112 + \frac{1}{6 \cdot (12 \cdot (8 \cdot (64 \cdot 7 + 1) + 1) + 1) + \frac{3}{4}}}$$

$$= 1/137.035999037435$$




$$\alpha_{1-133} = \frac{683}{133 \cdot (2\pi)_{12389}} \frac{1}{112 + \frac{14651}{50 \cdot 10^{11}}} = \frac{6^2 \cdot 19 - 1}{7 \cdot 19 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{59 \cdot 210}{13 \cdot (17 \cdot 56 + 1)}\right)^{71 \cdot (12 \cdot 29 + 1)}}} \frac{1}{112 + \frac{7^2 \cdot 13 \cdot 23}{50 \cdot 10^{11}}}$$

$$= 1/137.035999037435$$




$$\alpha_{1-140} = \frac{6^2 \cdot 20 - 1}{140 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{4 \cdot 13 \cdot 37}{3 \cdot (64 \cdot 10 + 1)}\right)^{3847}}} \frac{1}{112 + \frac{1}{4 \cdot 9 \cdot (2 \cdot 3 \cdot 29 \cdot 47 + 1) + \frac{29}{54}}}$$

$$= 1/137.035999037435$$



$$\alpha_{1-155} = \frac{2^2 \cdot 199}{5 \cdot 31 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left[\frac{19 \cdot 210 - 1}{3^2 \cdot (2 \cdot 13 \cdot 17 + 1) + 1}\right]^{7977}}} \frac{1}{112 + \frac{1}{5 \cdot 17 \cdot 31 \cdot (2 \cdot 13 \cdot 17 + 1) - \frac{15}{43}}}$$

$$= 1/137.035999037435$$



$$\alpha_{1-170} = \frac{873}{170 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{34451}{34450}\right)^{68901}}} \frac{1}{112 + \frac{4171}{8 \times 10^{11}}}$$

$$= \frac{3^2 \cdot 97}{170 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left[\frac{47(12 \cdot 61 + 1)}{2 \cdot 25 \cdot 13 \cdot 53}\right]^{3 \cdot 7 \cdot 17 \cdot 193}}} \frac{1}{112 + \frac{43 \cdot 97}{8 \cdot 10^{11}}} = 1/137.035999037435$$






Table 6. Parameters and Results of Approximate Formulas of α_2 (2019/7/3).

m	n	k	$\alpha_{2-m'}$	m	n	k	$\alpha_{2-m'}$
1	8	4	137.933814383	22	170	32	137.047480404
2	16	4	137.933814383	23	177	161	137.036664793
3	24	4	137.933814383	24	185	62	137.040748949
4	31	20	137.054511358	25	193	39	137.039552569
5	39	11	137.120466691	26	200	278	137.036218856
6	47	8	137.070996332	27	208	80	137.038980680
7	54	48	137.042951195	28	216	48	137.042951195
8	62	20	137.054511358	29	223	655	137.036002235
9	70	14	137.109928583	30	231	104	137.037530964
10	77	104	137.037530964	31	239	58	137.040063944
11	85	32	137.047480404	32	247	41	137.044550585
12	93	20	137.054511358	33	254	138	137.036795730
13	100	278	137.036218856	34	262	70	137.038016730
14	108	48	137.042951195	35	270	48	137.042951195
15	116	28	137.059466839	36	277	190	137.036562950
16	124	20	137.054511358	37	285	85	137.037566566
17	131	70	137.038016730	38	293	56	137.041569603
18	139	37	137.048943854	39	300	278	137.036218856
19	147	26	137.063298933	125	961	4293	137.035999678
20	154	104	137.037530964	253	1945	28186	137.035999128
21	162	48	137.042951195	269	2068	41654	137.035999118


Fig. 8. Results of Approximate Formulas of α_2 (2019/7/3).




$$\alpha_{2-1} = \frac{e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \frac{e^2}{\left(\frac{4}{3}\right)^7} \frac{e^2}{\left(\frac{5}{4}\right)^9}}{2 \cdot 2^2} \frac{1}{112 - 1 + \frac{1}{3} - \frac{1}{16} + \frac{1}{41 \cdot (12 \cdot 13 + 1) + \frac{13}{41}}} = 1/137.035999111816$$




$$\alpha_{2-4} = \frac{2^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{21}{20}\right)^{41}}}{31} \frac{1}{112 - \frac{1}{66} + \frac{1}{71 \cdot (14 \cdot 43 - 1) - \frac{56}{95}}} = 1/137.035999111818$$




$$\alpha_{2-5} = \frac{5 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{12}{11}\right)^{23}}}{39} \frac{1}{112 - \frac{1}{14} + \frac{1}{10 \cdot 41} - \frac{1}{23 \cdot (14 \cdot 11 \cdot 79 + 1) + \frac{11}{16}}} = 1/137.035999111818$$




$$\alpha_{2-6} = \frac{6 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{9}{8}\right)^{17}}}{47} \frac{1}{112 - \frac{1}{2 \cdot 17} + \frac{1}{2 \cdot (36 \cdot 17 + 1) - \frac{4}{47}}} = 1/137.035999111818$$



$$\alpha_{2-7} = \frac{7 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{48}{47}\right)^{95}}}{6 \cdot 3^2} \frac{1}{112 + \frac{1}{2 \cdot 13 \cdot 17} - \frac{1}{2 \cdot 29 \cdot (24 \cdot 89 + 1) + \frac{11}{2 \cdot 17}}} = 1/137.035999111818$$



$$\alpha_{2-9} = \frac{3^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{15}{14}\right)^{29}}}{70} \frac{1}{112 - \frac{1}{16} + \frac{1}{11 \cdot 43} - \frac{1}{70 \cdot 17 \cdot (3 \cdot 64 - 1) - \frac{41}{70}}} = 1/137.035999111818$$



$$\alpha_{2-10} = \frac{10 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{5 \cdot 21}{8 \cdot 13}\right)^{11 \cdot 9}}}{77} \frac{1}{112 - \frac{1}{3 \cdot 14 \cdot 19} + \frac{1}{14 \cdot (4 \cdot 27 \cdot (2 \cdot 15 \cdot 19 + 1) - 1)}} = 1/137.035999111818$$

¹⁴N₇ ²⁷Al₁₄ ²⁸Si₁₄ ^{39,40,41}K_{20,21,22} ⁵⁶Fe₃₀ ^{97,99}Mo_{54,56} ^{131,136}Xe_{77,82} ^{138,139}La_{81,82} ¹⁹¹Ir_{2,57} ²⁰⁹Bi₁₂₆* ²⁰⁹Po₁₂₅* ²⁸⁵Cn₁₇₃*
^{344,346,348}Fy_{208,209,210}^{ie} ^{418,420}Ch_{250,252}^{ie}

$$\alpha_{2-11} = \frac{11 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{33}{32}\right)^{65}}}{85} \frac{1}{112 - \frac{1}{106} + \frac{1}{30 \cdot (4 \cdot 163 + 1) - \frac{35}{52}}} = 1/137.035999111818$$

⁷⁹Br₄₄ ⁷⁹Br₄₄ ^{96,99,100,104}Ru₄₄ ¹²⁵Te_{52,55,56,60} ¹²⁷I₇₄ ¹⁴⁵Nd₈₅ ²¹⁰At₁₂₅* ²⁶⁸Db₁₆₃* ²⁶⁹Sg₁₆₃* ²⁷⁰Bh₁₆₃*
^{2,163}Ch_{128,129}^{ie} ^{198=1811,197} ⁴¹⁰Ch₁₆₃^{ie} ⁴²⁰Ch₁₆₅₌₅₋₃₃^{ie} ⁴²⁰Ch₁₇₀^{ie} ⁴²⁰Ch₂₅₀^{ie}

$$\alpha_2 = \alpha_{2-13} = \frac{13 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{9 \cdot 31}{2 \cdot 139}\right)^{557}}}{10^2} \frac{1}{112 - \frac{1}{3 \cdot 29 \cdot 64}} = 1/137.035999111818$$

¹⁹F₁₀ ²⁷Al₁₃ ²⁹Si₁₄ ^{54,56,57,58}Fe₂₆ ^{28,30,31,32}Ni₂₈ ^{63,65}Cu₂₉ ^{34,36} ^{69,71}Ga₃₁ ^{38,40} ⁸⁹Y₃₉ ⁹³Nb₄₁ ¹⁰⁰Ru₄₄ ^{112,116}Cd₄₈ ¹¹⁶Sn₅₀ ⁶⁶
¹²⁸Te₅₂ ^{136,137,138}Ba₅₆ ^{80,81,82} ¹³⁹La₅₇ ^{136,138,140,142}Ce₅₈ ^{78,80,82,84} ¹⁴⁹Sm₆₂ ^{156,157,160}Gd₆₄ ¹⁶⁶⁻¹⁶⁸Er₆₈ ¹⁶⁹Tm₆₉ ¹⁷⁵Lu₇₁ ^{8,13}
¹⁹²Os₇₆ ²⁰⁸Pb₈₂ ²⁰⁹Bi₈₃* ²²³Fr₈₇* ²³⁷Np₉₃* ²⁵⁷Fm₁₀₀* ¹⁵¹⁹Cn₁₁₂* ^{310,24-13}Ch_{124,125}^{ie} ^{186,187} ^{330,332}Ch_{130,131}^{ie} ^{200,201} ¹²⁻²⁹Ch₁₃₉^{ie} ⁶⁻⁷¹Ch₁₆₉^{ie} ²⁵⁷

$$\alpha_{2-15} = \frac{15 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{29}{28}\right)^{57}}}{2^2 \cdot 29} \frac{1}{112 - \frac{1}{4 \cdot 13} + \frac{1}{12 \cdot (36 \cdot 43 + 1) - \frac{1}{16}}} = 1/137.035999111818$$

^{24,25,26}Mg₁₂ ^{12,13,14} ³¹P₁₅ ⁵⁶Fe₂₆ ^{63,65}Cu₂₉ ^{34,36} ^{84,86}Kr₃₆ ^{48,50} ⁹⁹Tc₄₃* ²²²Rn₈₆* ¹³⁶ ²⁹³Ch₁₁₆₌₄₋₂₉^{ie} ^{3,59}

$$\alpha_{2-17} = \frac{17 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{71}{70}\right)^{3 \cdot 47}}}{131} \frac{1}{112 - \frac{1}{6 \cdot 101} + \frac{1}{23 \cdot (30 \cdot 35^2 - 1) + \frac{6}{23}}} = 1/137.035999111818$$

³⁵Cl₁₇ ⁵¹V₂₃ ^{64,66,68}Zn₃₀ ^{34,36,38} ^{80,82}Se₃₄ ^{46,48} ⁸¹Br₃₅ ⁸³Kr₃₆ ¹⁰⁶Pd₄₆ ¹⁰⁷Ag₄₇ ^{121,122}Sb₅₁ ¹⁴⁴Nd₆₀ ^{171,172}Yb₇₀ ^{101,102}
¹⁷⁵Lu₇₁ ²⁶²Lr₁₀₃* ^{312,314}Ch_{115,126}^{ie} ^{11-17,4-47} ^{330,4-83}Ch_{130,131}^{ie} ^{200,201} ^{4-89,358/360}Ch_{141=3-47,142=2-71}^{ie} ^{5-43,6-36/218}

$$\alpha_{2-18} = \frac{18 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{38}{37}\right)^{75}}}{139} \frac{1}{112 - \frac{1}{2 \cdot 47} + \frac{1}{2 \cdot 31 \cdot (2 \cdot 136 - 1) + \frac{83}{137}}} = 1/137.035999111818$$

³¹P₁₅ ^{32,33,34,36}S₁₆ ^{35,37}Cl₁₇ ^{36,38}Ar₁₈ ⁶⁹Ga₃₁ ^{70,72,74,76}Ge₃₂ ^{38,40,42,44} ^{74,76}Se₃₄ ^{83,84}Kr₃₆ ^{47,48}
¹⁰⁹Ag₄₇ ^{148,150,152}Sm₆₂ ^{86,88,90} ^{130,132,136,137,138}Ba₅₆ ^{74,76,80,81,82} ^{185,187}Re₇₅ ²⁰⁹Bi₈₃* ¹²⁶
^{310,312,314}Ch_{124,125,126}^{ie} ^{186,187,188} ^{344,346,348}Fy_{208,209,210}^{ie} ³⁴⁸Ch₁₃₉^{ie} ⁴¹⁸Ch₁₆₆^{ie} ²⁵²

$$\alpha_{2-19} = \frac{19 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{27}{26}\right)^{53}}}{3 \cdot 49} \frac{1}{112 - \frac{1}{44} + \frac{1}{16 \cdot (4 \cdot 37 + 1) - \frac{23}{6 \cdot 47 + 1}}} = 1/137.035999111818$$

$$\alpha_{2-23} = \frac{23 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{2 \cdot 81}{7 \cdot 23}\right)^{17 \cdot 19}}}{3 \cdot 59} \frac{1}{112 - \frac{1}{2 \cdot (40 \cdot 23 - 1) + \frac{9}{32 \cdot 10}}} = 1/137.035999111818$$

$$\alpha_{2-24} = \frac{2^2 \cdot 6 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{63}{2 \cdot 31}\right)^{125}}}{5 \cdot 37} \frac{1}{112 - \frac{1}{257} + \frac{1}{10 \cdot (12 \cdot 13 \cdot 83 + 1) + \frac{23}{81}}} = 1/137.035999111818$$


$$\alpha_{2-25} = \frac{5^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{40}{3 \cdot 13}\right)^{79}}}{193} \frac{1}{112 - \frac{1}{8 \cdot 43} + \frac{1}{18 \cdot 23 \cdot (32 \cdot 27 - 1) - \frac{3}{7}}} = 1/137.035999111818$$

$$\alpha_{2-27} = \frac{3 \cdot 3^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{81}{80}\right)^{7 \cdot 23}}}{4^2 \cdot 13} \frac{1}{112 - \frac{1}{10 \cdot 41} + \frac{1}{2 \cdot 27 \cdot 43 \cdot (3 \cdot 64 + 1) - \frac{19}{26}}} = 1/137.035999111818$$

$$\alpha_{2-29} = \frac{29 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{16 \cdot 41}{5 \cdot 131}\right)^{3 \cdot 19 \cdot 23}}}{223} \frac{1}{112 - \frac{1}{29 \cdot 59 \cdot (12 \cdot 19 + 1) + \frac{19}{29}}} = 1/137.035999111818$$


$$\alpha_{2-31} = \frac{31 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{16 \cdot 31}{7 \cdot 131}\right)^{3 \cdot 19 \cdot 23}}}{223} \frac{1}{112 - \frac{1}{29 \cdot 59 \cdot (12 \cdot 19 + 1) + \frac{19}{29}}} = 1/137.035999111818$$

$$\alpha_{2-31} = \frac{31 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{59}{58}\right)^{9 \cdot 13}}}{7 \cdot 34 + 1} \frac{1}{112 - \frac{1}{7 \cdot 43 + \frac{9}{5 \cdot 113}}} = 1/137.035999111819$$




$$\alpha_{2-32} = \frac{2 \cdot 4^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{42}{41}\right)^{83}}}{13 \cdot 19} \frac{1}{112 - \frac{1}{11 \cdot 13} + \frac{1}{6 \cdot 37 \cdot (5 \cdot 210 - 1) + \frac{10}{11}}} = 1/137.035999111818$$


$$= 1/137.035999111818$$




$$\alpha_{2-33} = \frac{33 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{139}{138}\right)^{277}}}{2 \cdot (2 \cdot 8^2 - 1)} \frac{1}{112 - \frac{1}{32 \cdot 48 - \frac{36}{35 \cdot 13}}} = 1/137.035999111818$$




$$\alpha_{2-36} = \frac{6^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{191}{190}\right)^{3 \cdot 127}}}{2 \cdot 138 + 1} \frac{1}{112 - \frac{1}{10 \cdot 7 \cdot 31 + \frac{13}{25 \cdot 23}}} = 1/137.035999111818$$



$$\alpha_{2-37} = \frac{37 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{2 \cdot 43}{5 \cdot 17}\right)^{9 \cdot 19}}}{3 \cdot 5 \cdot 19} \frac{1}{112 - \frac{1}{4 \cdot 3 \cdot 5 \cdot 13} + \frac{1}{5 \cdot 37^2 \cdot 149}}} = 1/137.035999111818$$




$$\alpha_{2-38} = \frac{2 \cdot 19 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{3 \cdot 19}{56}\right)^{113}}}{6 \cdot 7^2 - 1} \frac{1}{112 - \frac{1}{3 \cdot 73} + \frac{1}{30(8 \cdot 27 \cdot 17 + 1) - \frac{12}{13}}} = 1/137.035999111816$$



$$\alpha_{2-125} = \frac{5 \cdot 5^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{4294}{4293}\right)^{8587}}}{31^2} \frac{1}{112 - \frac{1}{2159481}}$$


$$= \frac{5 \cdot 5^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{2 \cdot 19 \cdot 113}{81 \cdot 53}\right)^{31 \cdot (12 \cdot 23 + 1)}}}{31^2} \frac{1}{112 - \frac{1}{3 \cdot 101 \cdot (8 \cdot 81 \cdot 11 - 1)}}} = 1/137.035999111818$$



$$\alpha_{2-253} = \frac{253 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{28187}{28186}\right)^{56373}}}{1945} \frac{1}{112 - \frac{10411}{8 \times 10^{11}}}$$

$$11 \cdot 23 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left[\frac{71(36 \cdot 11 + 1)}{2 \cdot 17(36 \cdot 23 + 1)}\right]^{3 \cdot 19 \cdot 23 \cdot 43}}$$

$$= \frac{1}{5(4 \cdot 97 + 1)} \frac{1}{112 - \frac{29(360 - 1)}{8 \times 10^{11}}} = 1/137.035999111818$$




$$\alpha_{2-269} = \frac{269 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{41655}{41654}\right)^{83309}}}{2068} \frac{1}{112 - 5.317 \times 10^{-9}}$$

$$(4 \cdot 67 + 1) \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{15 \cdot (6 \cdot (16 \cdot 29 - 1) - 1)}{2 \cdot 59 \cdot (6 \cdot 59 - 1)}\right)^{227 \cdot (6 \cdot 61 + 1)}}$$

$$= \frac{1}{4 \cdot 11 \cdot 47} \frac{1}{112 - \frac{13 \cdot (24 \cdot 17 + 1)}{10^{12}}}$$

$$= 1/137.035999111818$$



In above formulas, there are many amazing coincidences. As $136=8 \times 17$ and $138=6 \times 23$, 17 and 23 both appear in α_{1-1} , α_{1-17} , α_{1-22} , α_{1-23} , α_{1-25} , α_{1-59} , α_{1-103} , α_{1-133} , α_{2-17} and α_{2-23} , 17 frequently appears in α_1 and 23 frequently appears in α_2 . 157 and 257 in α_{1-50} should relate to ${}_{100}\text{Fm}_{157}^*$, 173 in α_{1-16} should relate to ${}_{112}\text{Cn}_{173}^*$, and so on. As the factors in formulas of α are reasonably assumed to relate to nuclides, some ideal extended elements such as ${}_{136,137,138}\text{Fy}_{208,209,210}$ and ${}_{169}\text{Ch}_{257}$ are predicted.

15. Radius of Electron and Proton

The classical electron radius r_e has been calculated very accurately. However, the proton charge radius r_p hasn't yet been determined precisely. Recent two experiments

measured r_p and had given the best results up to now which was $r_p=0.833(19) \text{ fm}^9$ and $r_p=0.831(19) \text{ fm}^{10}$, and hence CODATA revised its recommended data of r_p to $0.8414(19) \text{ fm}$. Here we give our calculation results of r_e and r_p . And it seems there is α_p similar to α . α_p could be called “the second fine-structure constant”.

Ratio of Bohr radius of hydrogen atom to classical electron radius:

$$\frac{a_0}{r_e} = \frac{1}{\alpha_c^2} = \frac{1}{\alpha_1 \alpha_2} = 112 \times \left(168 - \frac{1}{3} + \frac{1}{2^2 \cdot 3 \cdot 47} - \frac{1}{2 \cdot 3 \cdot 29 \cdot 53 \cdot 59 - 79 / 47} \right) = 18788.865042381$$

$$r_e = \alpha_c^2 a_0 = \alpha_1 \alpha_2 a_0 = \frac{5.29177210903(80) \times 10^{-11} \text{ m}}{18788.865042381} = 2.81794032658(43) \text{ fm}$$

Comparable to CODATA recommended value $r_e = 2.8179403262(13) \text{ fm}$ but more precise.

Ratio of Bohr radius of hydrogen atom to the proton charge radius should have the similar form, and is assumed to have the following hypothetical formulas:

$$\frac{a_0}{r_p} = \frac{1}{\alpha_{p/c}^2} = \frac{1}{\alpha_{p/1} \alpha_{p/2}} = 225 \cdot \left(282 + \frac{1}{3} - \frac{1}{12 \cdot 47} + \frac{1}{6 \cdot 29 \cdot 53 \cdot 59 - 79 / 47} \right) = 63524.60147736$$


$$= 247 \cdot \left(257 + \frac{1}{5} - \frac{1}{5 \cdot 13} + \frac{1}{30 \cdot (28 \cdot (2 \cdot 100 - 1) + 1) + \frac{8}{45}} \right)$$

$$= \left(252 + \frac{1}{24} - \frac{1}{2 \cdot 17 \cdot 37} + \frac{1}{11 \cdot 13 \cdot 19 \cdot (2 \cdot 11 \cdot 19 + 1) + \frac{11}{20}} \right)^2 = 252.040872632515^2$$


$$r_p = \alpha_{p/c}^2 a_0 = \alpha_{p/1} \alpha_{p/2} a_0 = \frac{5.29177210903(80) \times 10^{-11} \text{ m}}{63524.60147736} = 0.833027202999(13) \text{ fm}$$

$\alpha_{p/c} \approx \alpha_{p/1} \approx \alpha_{p/2} \approx 252.04$, α_p could be called the second fine-structure constant.

2019/12/19-23




$$\alpha_{p/1} = \frac{5 \cdot 3^2}{8 \cdot (2\pi)_{58}} \frac{1}{225 + \frac{1}{4 \cdot 112} - \frac{1}{5 \cdot 19 \cdot 83 \cdot 103 + \frac{19}{20}}} = 1 / 252.040872632515$$



$$\alpha_{p/2} = \frac{23 \cdot (2\pi)_{227}}{2 \cdot 9^2} \frac{1}{225 - \frac{1}{3 \cdot 16 \cdot 29 - \frac{71}{2 \cdot 67}}} = 1 / 252.040872632514$$

2020/1/2



$$\alpha_{p/2} = \frac{22 \cdot (2\pi)_{164}}{5 \cdot 31} \frac{1}{225 - \frac{1}{7 \cdot 137 - \frac{7 \cdot 13}{197}}} = 1/252.040872632512 \quad 2020/1/3$$

47,48,49,50 ²²Ti_{25,26,27,28} 56,57 ²⁶Fe_{30,31} 69,71 ³¹Ga_{38,40} 89 ³⁹Y₅₀ 99,100 ⁴⁴Ru_{55,56} 134,136,137,138 ⁵⁶Ba_{78,80,81,82} 5-31 ⁶⁴Gd₇₋₁₃ 164 ⁶⁶Dy₉₈ 196,198 ⁷⁸Pt_{118,120}

$$\alpha_{p/2} = \frac{21 \cdot (2\pi)_{126}}{2^2 \cdot 37} \frac{1}{225 - \frac{1}{16 \cdot 29} + \frac{1}{20 \cdot 13^2 \cdot 179 + \frac{8}{17}}} = 1/252.040872632515 \quad 2020/1/3$$

197 ⁷⁹Au₁₁₈ 206,207,208 ⁸²Pb_{124,125,126} 223 ⁸⁷Ra₁₃₆ 226 ⁸⁸Ra₁₃₈ 310,312 ⁸⁸Ch_{186,187}^{ie} 326,326/328 ⁸⁸Ch_{186,187}^{ie} 128,129 ¹²⁶Ch_{198,197/199}^{ie} 12-31 ¹⁴⁷Ch₂₂₅^{ie} 155=5-31,156=12-13 8-49,2-197 ¹⁶⁴Ch_{237,238}^{ie} 32-13 ¹⁶⁴Ch₂₅₂^{ie}

45 ²¹Sc₂₄ 63,65 ²⁹Cu_{34,36} 85,87 ³⁷Rb_{48,50} 126 ⁵²Te₇₄ 148 ⁶⁰Nd₈₈ 169 ⁶⁹Tm₁₀₀ 179 ⁷²Hf₁₀₇ 16-13 ⁸²Pb₁₂₆ 298,300 ^{119,120}Ch_{179,180}^{ie} 312,314 ^{125,126}Ch_{187,188}^{ie} 426 ¹⁶⁹Ch₂₅₇^{ie}

16. Direct Relationships of 2π with Nuclides

In Chen's formulas of the fine-structure constant, there are 2π-e formulas, in which k gets certain numbers and relate to nucleon numbers of some nuclides. So in the end of this paper we feel curious about whether 2π directly relate to nuclides.

$$2\pi = 6.2831853 \dots \approx \frac{4 \cdot 157}{100} = 6.28 \approx \frac{3 \cdot 7 \cdot 44 \cdot 68}{100^2} = 6.2832 \quad \begin{matrix} 7 \\ 3 \end{matrix} \text{Li}_4 \quad \begin{matrix} 100 \\ 44 \end{matrix} \text{Ru}_{56} \quad \begin{matrix} 157 \\ 64 \end{matrix} \text{Gd}_{93} \quad \begin{matrix} 168 \\ 68 \end{matrix} \text{Er}_{100} \quad \begin{matrix} 257 \\ 100 \end{matrix} \text{Fm}_{157}^* \quad \begin{matrix} 400 \\ 157 \end{matrix} \text{Ch}_{243}^{\text{ie}}$$

$$2\pi \approx \frac{4 \cdot 157}{100} = \frac{157}{25} = 6.28 \quad \begin{matrix} 55 \\ 25 \end{matrix} \text{Mn}_{30} \quad \begin{matrix} 100 \\ 44 \end{matrix} \text{Ru}_{56} \quad \begin{matrix} 157 \\ 64 \end{matrix} \text{Gd}_{93} \quad \begin{matrix} 118,119,120 \\ 50 \end{matrix} \text{Sn}_{68,69,70} \quad \begin{matrix} 168 \\ 68 \end{matrix} \text{Er}_{100} \quad \begin{matrix} 169 \\ 69 \end{matrix} \text{Tm}_{100} \quad \begin{matrix} 185,187 \\ 75 \end{matrix} \text{Re}_{110,112}$$

$$2\pi \approx \frac{16 \cdot 3 \cdot 7 \cdot 11 \cdot 17}{100^2} = \frac{48 \cdot 7 \cdot 11 \cdot 17}{100^2} = \frac{3 \cdot 7 \cdot 44 \cdot 68}{100^2} = \frac{3 \cdot 112 \cdot 11 \cdot 17}{100^2} = \frac{2 \cdot 168 \cdot 11 \cdot 17}{100^2} = \frac{2 \cdot 3 \cdot 7 \cdot 11 \cdot 136}{100^2} = \dots = 6.2832$$

⁷Li₄ ²⁰⁻²²Ne₁₀₋₁₂ ²³Na₁₂ ⁴⁵Sc₂₄ 46,47,49,50 ²²Ti_{24,25,27,28} ⁶¹Ni₃₃ ⁵⁵Mn₃₀ 54,56 ²⁶Fe_{28,30} 78,80 ³⁴Se_{44,46} 98,100 ⁴²Mo_{56,58} ¹⁰⁰Ru₅₆ ¹¹²Cd₄₈ ^{136,137,138}Ba_{80,81,82} ¹⁶⁸Er₁₀₀ ^{185,11-17}Re_{110,112} ²⁰⁹Po₁₂₅ ²²²Rn₁₃₆ ²²³Fa₁₃₆ ²²⁶Ra₁₃₈ ²²⁷Ac₁₃₈ ²⁷⁸⁺⁷Cn₁₇₃ ^{344,346,348}Fy_{208,209,210}^{ie}

$$2\pi \approx \frac{44}{7} = \frac{2 \cdot 22}{7} = 6.2857 \dots \quad \begin{matrix} 50 \\ 22 \end{matrix} \text{Ti}_{28} \quad \begin{matrix} 61 \\ 28 \end{matrix} \text{Ni}_{33} \quad \begin{matrix} 100 \\ 44 \end{matrix} \text{Ru}_{56} \quad \begin{matrix} 136,137,138 \\ 56 \end{matrix} \text{Ba}_{80,81,82} \quad \begin{matrix} 226 \\ 88 \end{matrix} \text{Ra}_{138} \quad \begin{matrix} 294 \\ 118 \end{matrix} \text{Og}_{176} \quad \begin{matrix} 8-44 \\ 140 \end{matrix} \text{Ch}_{212}^{\text{ie}} \quad \begin{matrix} 22-17 \\ 148=4 \cdot 37 \end{matrix} \text{Ch}_{226}^{\text{ie}}$$

$$2\pi \approx \frac{201}{32} = \frac{3 \cdot 67}{32} = 6.2812 \dots \quad \begin{matrix} 32 \\ 16 \end{matrix} \text{S}_{16} \quad \begin{matrix} 59 \\ 27 \end{matrix} \text{Co}_{30} \quad \begin{matrix} 67 \\ 30 \end{matrix} \text{Zn}_{37} \quad \begin{matrix} 112 \\ 48 \end{matrix} \text{Cd}_{64} \quad \begin{matrix} 117 \\ 50 \end{matrix} \text{Sn}_{67} \quad \begin{matrix} 128,134 \\ 54 \end{matrix} \text{Xe}_{74,80} \quad \begin{matrix} 134 \\ 56 \end{matrix} \text{Ba}_{78} \quad \begin{matrix} 165 \\ 67 \end{matrix} \text{Ho}_{98} \quad \begin{matrix} 201 \\ 80 \end{matrix} \text{Hg}_{121} \quad \begin{matrix} 332 \\ 131 \end{matrix} \text{Ch}_{201}^{\text{ie}} \quad \begin{matrix} 402 \\ 158 \end{matrix} \text{Ch}_{244}^{\text{ie}}$$

$$2\pi \approx \frac{245}{39} = \frac{5 \cdot 7^2}{3 \cdot 13} = 6.2820 \dots \quad \begin{matrix} 7 \\ 3 \end{matrix} \text{Li}_4 \quad \begin{matrix} 27 \\ 13 \end{matrix} \text{Al}_{14} \quad \begin{matrix} 54,56 \\ 26 \end{matrix} \text{Fe}_{28,30} \quad \begin{matrix} 89 \\ 39 \end{matrix} \text{Y}_{50} \quad \begin{matrix} 79,81 \\ 35 \end{matrix} \text{Br}_{44,46} \quad \begin{matrix} 113,115 \\ 49 \end{matrix} \text{In}_{64,66} \quad \begin{matrix} 24-13,314 \\ 125,126 \end{matrix} \text{Ch}_{187,188}^{\text{ie}}$$

$$2\pi \approx \frac{289}{46} = \frac{17^2}{2 \cdot 23} = 6.2826 \dots \quad \begin{matrix} 3-17 \\ 23 \end{matrix} \text{V}_{28} \quad \begin{matrix} 78,80 \\ 34 \end{matrix} \text{Se}_{44,46} \quad \begin{matrix} 6-17 \\ 46 \end{matrix} \text{Pd}_{56} \quad \begin{matrix} 168 \\ 68 \end{matrix} \text{Er}_{100} \quad \begin{matrix} 169 \\ 69 \end{matrix} \text{Tm}_{100} \quad \begin{matrix} 136,137,138 \\ 56 \end{matrix} \text{Ba}_{80,81,82} \quad \begin{matrix} 11-17 \\ 75 \end{matrix} \text{Re}_{112} \quad \begin{matrix} 222 \\ 86 \end{matrix} \text{Rn}_{136}^*$$

$$2\pi \approx \frac{333}{53} = \frac{9 \cdot 37}{53} = 6.2830 \dots \quad \begin{matrix} 223 \\ 87 \end{matrix} \text{Fa}_{136}^* \quad \begin{matrix} 226 \\ 88 \end{matrix} \text{Ra}_{138}^* \quad \begin{matrix} 227 \\ 89 \end{matrix} \text{Ac}_{138}^* \quad \begin{matrix} 238 \\ 92 \end{matrix} \text{U}_{146}^* \quad \begin{matrix} 17-17 \\ 114 \end{matrix} \text{Fl}_{175}^{\text{ie}} \quad \begin{matrix} 344,346,348 \\ 136,137,138 \end{matrix} \text{Fy}_{208,209,210}^{\text{ie}} \quad \begin{matrix} 22-17 \\ 148=4 \cdot 37 \end{matrix} \text{Ch}_{226}^{\text{ie}}$$

$$2\pi \approx \frac{377}{60} = \frac{13 \cdot 29}{4 \cdot 3 \cdot 5} = 6.2833 \dots \quad \begin{matrix} 85,87 \\ 37 \end{matrix} \text{Rb}_{48,50} \quad \begin{matrix} 3 \cdot 37=111 \\ 48 \end{matrix} \text{Cd}_{7-9} \quad \begin{matrix} 127 \\ 53 \end{matrix} \text{I}_{74} \quad \begin{matrix} 180,184,189 \\ 74 \end{matrix} \text{W}_{106,110,112} \quad \begin{matrix} 222 \\ 86 \end{matrix} \text{Rn}_{136}^* \quad \begin{matrix} 269 \\ 106 \end{matrix} \text{Sg}_{163} \quad \begin{matrix} 280 \\ 111 \end{matrix} \text{Rg}_{169} \quad \begin{matrix} 22-17 \\ 148 \end{matrix} \text{Ch}_{226}^{\text{ie}}$$

$$2\pi \approx \frac{465}{74} = \frac{30 \cdot 31}{4 \cdot 37} = 6.2837 \dots \quad \begin{matrix} 24,25,26 \\ 12 \end{matrix} \text{Mg}_{12,13,14} \quad \begin{matrix} 28,29,30 \\ 14 \end{matrix} \text{Si}_{14,15,16} \quad \begin{matrix} 31 \\ 15 \end{matrix} \text{P}_{16} \quad \begin{matrix} 54,56 \\ 26 \end{matrix} \text{Fe}_{28,30} \quad \begin{matrix} 63,65 \\ 29 \end{matrix} \text{Cu}_{34,36} \quad \begin{matrix} 116,120 \\ 50 \end{matrix} \text{Sn}_{66,70}$$

$$2\pi \approx \frac{509}{81} = \frac{2 \cdot 3 \cdot 5 \cdot 17 - 1}{9^2} = 6.2839 \dots \quad \begin{matrix} 140,142 \\ 58 \end{matrix} \text{Ce}_{82,84} \quad \begin{matrix} 144,145,146,148,150 \\ 60 \end{matrix} \text{Nd}_{84,85,86,88,90} \quad \begin{matrix} 200 \\ 80 \end{matrix} \text{Hg}_{120} \quad \begin{matrix} 223 \\ 87 \end{matrix} \text{Fa}_{136}^* \quad \begin{matrix} 24-13,314 \\ 125,126 \end{matrix} \text{Ch}_{187,188}^{\text{ie}}$$

$$2\pi \approx \frac{622}{99} = \frac{4 \cdot (24 \cdot 13 - 1)}{9 \cdot 22} = 6.2828 \dots \quad \begin{matrix} 31 \\ 15 \end{matrix} \text{P}_{16} \quad \begin{matrix} 67 \\ 30 \end{matrix} \text{Zn}_{37} \quad \begin{matrix} 69,71 \\ 31 \end{matrix} \text{Ga}_{38,40} \quad \begin{matrix} 6-31 \\ 74 \end{matrix} \text{W}_{112} \quad \begin{matrix} 85,67 \\ 37 \end{matrix} \text{Rb}_{48,50} \quad \begin{matrix} 4-37 \\ 60 \end{matrix} \text{Nd}_{88} \quad \begin{matrix} 157 \\ 64 \end{matrix} \text{Gd}_{93} \quad \begin{matrix} 243 \\ 95 \end{matrix} \text{Am}_{4-37}^* \quad \begin{matrix} 22-17 \\ 148=4 \cdot 37 \end{matrix} \text{Ch}_{226}^{\text{ie}}$$

$$2\pi \approx \frac{509}{81} = \frac{2 \cdot 3 \cdot 5 \cdot 17 - 1}{9^2} = 6.2839 \dots \quad \begin{matrix} 19 \\ 9 \end{matrix} \text{F}_{10} \quad \begin{matrix} 35,37 \\ 17 \end{matrix} \text{Cl}_{18,20} \quad \begin{matrix} 64,70 \\ 30 \end{matrix} \text{Zn}_{34,40} \quad \begin{matrix} 80,82 \\ 34 \end{matrix} \text{Se}_{46,48} \quad \begin{matrix} 136,137,138 \\ 56 \end{matrix} \text{Ba}_{80,81,82} \quad \begin{matrix} 203,205 \\ 81 \end{matrix} \text{Tl}_{61,62} \quad \begin{matrix} 210 \\ 85 \end{matrix} \text{At}_{125}^*$$

$$2\pi \approx \frac{622}{99} = \frac{4 \cdot (24 \cdot 13 - 1)}{9 \cdot 22} = 6.2828 \dots \quad \begin{matrix} 3-81 \\ 95 \end{matrix} \text{Am}_{148}^* \quad \begin{matrix} 344,346,348 \\ 136,137,138 \end{matrix} \text{Fy}_{208,209,210}^{\text{ie}} \quad \begin{matrix} 22-17 \\ 148=4 \cdot 37 \end{matrix} \text{Ch}_{226}^{\text{ie}} \quad \begin{matrix} 400 \\ 157 \end{matrix} \text{Ch}_{243}^{\text{ie}}$$

$$2\pi \approx \frac{622}{99} = \frac{4 \cdot (24 \cdot 13 - 1)}{9 \cdot 22} = 6.2828 \dots \quad \begin{matrix} 23 \\ 11 \end{matrix} \text{Na}_{12} \quad \begin{matrix} 27 \\ 12 \end{matrix} \text{Al}_{13} \quad \begin{matrix} 46,48,49 \\ 22 \end{matrix} \text{Ti}_{24,26,27} \quad \begin{matrix} 50,52,54 \\ 24 \end{matrix} \text{Cr}_{26,28,30} \quad \begin{matrix} 54,56,58 \\ 26 \end{matrix} \text{Fe}_{28,30,32} \quad \begin{matrix} 99 \\ 44 \end{matrix} \text{Ru}_{55} \quad \begin{matrix} 167 \\ 68 \end{matrix} \text{Er}_{99} \quad \begin{matrix} 252 \\ 99 \end{matrix} \text{Es}_{153}^*$$

$$2\pi \approx \frac{2 \cdot 355}{113} = \frac{4 \cdot 5 \cdot 71}{2 \cdot 113} = 6.2831858 \dots \quad \begin{matrix} 71 \\ 31 \end{matrix} \text{Ga}_{40} \quad \begin{matrix} 112,113 \\ 48 \end{matrix} \text{Cd}_{64,65} \quad \begin{matrix} 113,115 \\ 49 \end{matrix} \text{In}_{64,66} \quad \begin{matrix} 120,122 \\ 50 \end{matrix} \text{Sn}_{70,72} \quad \begin{matrix} 2-71 \\ 60 \end{matrix} \text{Nd}_{82} \quad \begin{matrix} 171 \\ 70 \end{matrix} \text{Yb}_{101} \quad \begin{matrix} 175 \\ 71 \end{matrix} \text{Lu}_{74} \quad \begin{matrix} 186 \\ 74 \end{matrix} \text{W}_{112}$$

187 ⁷⁵Re₁₁₂ 188,189 ⁷⁶Os_{112,113} 226 ⁸⁸Ra₁₃₈ 232 ⁹⁰Th₁₃₈ 4-71 ¹¹³Nh₁₇₁^{ie} 358/360 ¹⁴²⁼²⁻⁷¹Ch_{6-36/218}^{ie} 22-17 ^{148=4 \cdot 37}Ch₂₂₆^{ie} 6-71 ¹⁶⁹Ch₂₅₇^{ie} 2020/1/8-10

The approximate rational numbers of 2π (could be called 2π formulas) relate to nuclides marvelously. This means 2π (along with 2π -e formula) plays important roles in atomic nuclei, and acts as a rational number rather than an irrational number in the world of atomic nuclei.

17. Correlations among α , 2π and nuclides

Some Chen's formulas of the fine-structure constant and 2π formulas correlate with each others with the same factors and all together relate to the same nuclides. For example, α_{1-9} and $2\pi \approx 4 \times 157/100$ have the same 157 and 100 factors, α_{1-50} and $2\pi \approx 3 \times 7 \times 44 \times 68/100^2$ have the same 100, 7, 11 and 16 factors, and they relate to the same corresponding nuclides. They also have common factors with α_{1-7} and α_{2-13} which should relate to $2\pi \approx 5 \times 7^2/3/13$ and $2\pi \approx 13 \times 29/4/3/5$.

$$\alpha_{1-9} = \frac{47}{3^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{10}{9}\right)^{19}} 112 + \frac{1}{4 \cdot 17}} - \frac{1}{2(8 \cdot 9 \cdot 37 - 1) + \frac{3 \cdot 17}{157}}$$

$$\alpha_{1-50} = \frac{2 \cdot 257}{100 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{14 \cdot 13}{181}\right)^{3 \cdot 11^2}} 112 + \frac{1}{29 \cdot 61 + \frac{157}{16 \cdot 11}}}$$

$$2\pi \approx \frac{4 \cdot 157}{100} = \frac{157}{25} = 6.28 \quad 2\pi \approx \frac{3 \cdot 7 \cdot 44 \cdot 68}{100^2} = \frac{3 \cdot 7 \cdot 11 \cdot 17}{25^2} = 6.2832$$

$$\alpha_{1-7} = \frac{6^2}{7 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{113}{112}\right)^{225}} 112 + \frac{1}{75^2}} \quad \alpha_{2-13} = \frac{13 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{279}{278}\right)^{557}}}{10^2} - \frac{1}{112 - \frac{1}{3 \cdot 29 \cdot 64}}$$

$$2\pi \approx \frac{245}{39} = \frac{5 \cdot 7^2}{3 \cdot 13} = 6.2820 \dots \quad 2\pi \approx \frac{377}{60} = \frac{13 \cdot 29}{4 \cdot 3 \cdot 5} = 6.2833 \dots$$

α_{1-22} relates to $2\pi \approx 2 \times 22/7$, $2\pi \approx 17^2/7/23$ and $2\pi \approx 2 \times 355/113$ as follows. And $2\pi \approx 17^2/7/23$ also relates to α_{1-1} , α_{1-17} , α_{1-22} , α_{1-23} , α_{1-25} , α_{1-59} , α_{1-103} , α_{1-133} , α_{2-17} and α_{2-23} , in which both 17 and 23 factors appear.

$$\alpha_{1-22} = \frac{113}{22 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{27 \cdot 29}{2 \cdot 17 \cdot 23}\right)^{5 \cdot (2 \cdot 157 - 1)}} 112 + \frac{1}{2 \cdot [2 \cdot 3 \cdot 17 \cdot (10 \cdot 19 + 1) + 1] + \frac{29}{49}}}$$

$$2\pi \approx \frac{2 \cdot 22}{7} = 6.2857 \dots, \quad 2\pi \approx \frac{17^2}{2 \cdot 23} = 6.2826 \dots, \quad 2\pi \approx \frac{2 \cdot 355}{113} = \frac{4 \cdot 5 \cdot 71}{2 \cdot 113} = 6.2831858 \dots$$

α_{1-13} and α_{1-43} relate to $2\pi \approx 3 \times 67/32$, $2\pi \approx 5 \times 7^2/39$, $2\pi \approx 17^2/46$ and others as follows.

$$\alpha_{1-13} = \frac{67}{13 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{47}{2 \cdot 23}\right)^{3 \cdot 31}}} \frac{1}{112 + \frac{1}{137} - \frac{1}{6(2 \cdot 27 \cdot 59 + 1) + \frac{9}{50}}}$$

$$\alpha_{1-43} = \frac{13 \cdot 17}{43 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{3 \cdot 67}{200}\right)^{401}}} \frac{1}{112 + \frac{1}{8 \cdot (12 \cdot 83 + 1) + \frac{4}{3 \cdot 13}}}$$

$$2\pi \approx \frac{3 \cdot 67}{32}, 2\pi \approx \frac{5 \cdot 7^2}{3 \cdot 13}, 2\pi \approx \frac{289}{46} = \frac{17^2}{2 \cdot 23}, 2\pi \approx \frac{13 \cdot 29}{60}, 2\pi \approx \frac{30 \cdot 31}{4 \cdot 37}$$

α_{1-11} , α_{1-36} , α_{2-24} , α_{2-23} , α_{2-37} and α_{2-125} relate to $2\pi \approx 9 \times 37/53$, $2\pi \approx 15 \times 31/2/37$ and $2\pi \approx (30 \times 17 - 1)/81$ as follows.

$$\alpha_{1-11} = \frac{57}{11 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{19}{18}\right)^{37}}} \frac{1}{112 + \frac{1}{35} - \frac{1}{88 \cdot 41 - \frac{5 \cdot 53}{22 \cdot 13}}}$$

$$\alpha_{1-36} = \frac{5 \cdot 37}{6^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{3 \cdot 79}{4 \cdot 59}\right)^{11 \cdot 43}}} \frac{1}{112 + \frac{1}{5 \cdot (31 \cdot 42 - 1) + \frac{3 \cdot 31}{14 \cdot 13}}}$$

$$\alpha_{2-24} = \frac{2^2 \cdot 6 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{63}{2 \cdot 31}\right)^{125}}}{5 \cdot 37} \frac{1}{112 - \frac{1}{257} + \frac{1}{10 \cdot (12 \cdot 13 \cdot 83 + 1) + \frac{23}{81}}}$$

$$\alpha_{2-23} = \frac{23 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{2 \cdot 81}{7 \cdot 23}\right)^{17 \cdot 19}}}{3 \cdot 59} \frac{1}{112 - \frac{1}{2 \cdot (40 \cdot 23 - 1) + \frac{9}{32 \cdot 10}}}$$


$$\alpha_{2-37} = \frac{37 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{2 \cdot 43}{5 \cdot 17}\right)^{9 \cdot 19}}}{3 \cdot 5 \cdot 19} \frac{1}{112 - \frac{1}{4 \cdot 3 \cdot 5 \cdot 13} + \frac{1}{5 \cdot 37^2 \cdot 149}}$$

$$\alpha_{2-125} = \frac{5 \cdot 5^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{2 \cdot 19 \cdot 113}{81 \cdot 53}\right)^{31 \cdot (12 \cdot 23 + 1)}}}{31^2} \frac{1}{112 - \frac{1}{101 \cdot (20 \cdot (12 \cdot 89 + 1) + 1)}}$$

$$2\pi \approx \frac{9 \cdot 37}{53} = 6.2830 \dots \quad 2\pi \approx \frac{15 \cdot 31}{2 \cdot 37} = 6.2837 \dots \quad 2\pi \approx \frac{30 \cdot 17 - 1}{81} = 6.2839 \dots$$

18. Chen's Mathematic Shell Model of Nuclides

In overall, there are multi-correlations among α , 2π and nuclides. It seems there should be a mathematical shell model of nuclides, in which the core is 2π formulas and the middle layer is 2π -e formulas and the outer layer is Chen's formulas of α (**Fig. 9**, ϕ is explained in **Section 21**). The nucleon numbers, stability and abundance of nuclides are regulated by these formulas, especially by their integer factors.



Chen's Mathematic Shell Model of Nuclides

Dr. Gang Chen (2020/1/12-13, 3/1)

Fig. 9

19. Ideal Extended Elements

In the deduction of Chen's formulas of the fine-structure constant, it was reasonably assumed the factors in them related to nucleon numbers of nuclides, and it seems this assumption is quite correct. So by somewhat correlation and decoding methodology, all 119th to 170th ideal extended elements were predicted (**Table 7**). In addition, nuclides can even relate to naked 2π 's approximate rational numbers (2π formulas). Some typical examples of correlations of ideal extended elements with formulas of α and 2π are listed as follows.

Example 1: Correlations of 100, 121, 125, 126, 157, 257, 169, *et al.*

$$\alpha_{1-9} = \frac{47}{3^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{10}{9}\right)^{19}} 112 + \frac{1}{4 \cdot 17}} \frac{1}{2(8 \cdot 9 \cdot 37 - 1) + \frac{3 \cdot 17}{157}}$$

$$\alpha_{1-50} = \frac{2 \cdot 257}{100 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{14 \cdot 13}{181}\right)^{3 \cdot 11^2}} 112 + \frac{1}{29 \cdot 61 + \frac{157}{16 \cdot 11}}}$$

$$\alpha_{2-24} = \frac{2^2 \cdot 6 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{63}{62}\right)^{125}}}{5 \cdot 37} \frac{1}{112 - \frac{1}{257} + \frac{1}{10 \cdot (12 \cdot 13 \cdot 83 + 1) + \frac{23}{81}}}$$

$$2\pi \approx \frac{4 \cdot 157}{100} \quad 2\pi \approx \frac{16 \cdot 3 \cdot 7 \cdot 11 \cdot 17}{100} \quad 2\pi \approx \frac{4 \cdot 11}{7} \quad 2\pi \approx \frac{17^2}{2 \cdot 23} \quad 2\pi \approx \frac{30 \cdot 31}{4 \cdot 37} \quad 2\pi \approx \frac{4 \cdot 5 \cdot 71}{2 \cdot 113}$$

${}^{100}_{44}\text{Ru}_{56}$ ${}^{168,169}_{68,69}\text{Tm}_{100}$ ${}^{257}_{100}\text{Fm}_{157}^*$ ${}^{302}_{121=11^2}\text{Ch}_{181}^{ie}$ ${}^{24 \cdot 13 \cdot 2 \cdot 157}_{125,126}\text{Ch}_{117,4 \cdot 47}^{ie}$ ${}^{2 \cdot 11 \cdot 17}_{148=4 \cdot 37}\text{Ch}_{226}^{ie}$ ${}^{400,402}_{157,158}\text{Ch}_{3 \cdot 81,4 \cdot 61}^{ie}$ ${}^{6 \cdot 71}_{169}\text{Ch}_{257}^{ie}$

Example 2: Correlations of 83, 126, 84, 125, 209, 112, 173, 285, 115 and 137

$$\alpha_{1-16} = \frac{83}{4^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{17}{16}\right)^{33}} 112 + \frac{1}{28} - \frac{1}{6 \cdot (18 \cdot 41 + 1) + \frac{173}{2 \cdot (2 \cdot 75 - 1)}}$$

$$\alpha_{1-25} = \frac{3 \cdot 43}{5^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{35}{34}\right)^{3 \cdot 23}} 112 + \frac{1}{11 \cdot 19} - \frac{1}{13^2(2 \cdot 136 - 1) + \frac{11}{25}}$$

$$\alpha_{1-32} = \frac{15 \cdot 11}{2 \cdot 2^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{41}{40}\right)^{81}} 112 + \frac{1}{25 \cdot 29} - \frac{1}{19 \cdot 23}$$

$$\alpha_{2-10} = \frac{10 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{5 \cdot 21}{8 \cdot 13}\right)^{11 \cdot 19}}}{77} \frac{1}{112 - \frac{1}{3 \cdot 14 \cdot 19} + \frac{1}{14 \cdot (4 \cdot 27 \cdot (2 \cdot 15 \cdot 19 + 1) - 1)}}$$

$$\alpha_{2-32} = \frac{2 \cdot 4^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{42}{41}\right)^{83}}}{13 \cdot 19} \frac{1}{112 - \frac{1}{11 \cdot 13} + \frac{1}{6 \cdot 37 \cdot (5 \cdot 210 - 1) + \frac{10}{11}}}$$

$$\alpha_{1-13} = \frac{67}{13 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{47}{2 \cdot 23}\right)^{3 \cdot 31}} 112 + \frac{1}{137} - \frac{1}{6(2 \cdot 27 \cdot 59 + 1) + \frac{9}{50}}$$

$$\alpha_{1-17} = \frac{2^2 \cdot 22}{17 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{21}{20}\right)^{41}} 112 + \frac{1}{137} - \frac{1}{2 \cdot 19 \cdot 23 \cdot 59 - \frac{30}{100}}$$

$$\alpha_{1-27} = \frac{139}{27 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{67}{66}\right)^{7 \cdot 19}} 112 + \frac{1}{11 \cdot 47 + \frac{18}{23} + \frac{1}{6 \cdot 23 \cdot 137}}$$

$$\alpha_{2-18} = \frac{18 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{38}{37}\right)^{75}}}{139} \frac{1}{112 - \frac{1}{2 \cdot 47} + \frac{1}{2 \cdot 31 \cdot (16 \cdot 17 - 1) + \frac{83}{137}}}$$

$$2\pi \approx \frac{16 \cdot 3 \cdot 7 \cdot 11 \cdot 17}{100} \approx \frac{4 \cdot 11}{7} \approx \frac{17^2}{2 \cdot 23} \quad \begin{matrix} 8-17, 137, 6-23 \\ 56 \end{matrix} Ba_{80,81,82} \quad \begin{matrix} 11-19 \\ 83 \end{matrix} D_{126}^* \quad \begin{matrix} 209 \\ 84 \end{matrix} Po_{125}^* \quad \begin{matrix} 210 \\ 85 \end{matrix} At_{125}^* \quad \begin{matrix} 15-19 \\ 112 \end{matrix} Cn_{173}^* \quad \begin{matrix} 2-173 \\ 137 \end{matrix} Fy_{209}^{ie}$$

Table 7. Correlations of Ideal Extended Elements (IEE) with Formulas of α and 2π .

IEE	Page	α	2π
¹¹³ Nh ₁₇₁	10 19 21 28 29 31	$\alpha_c^2 \alpha_{1-5,7} \alpha_{2-22,23,31,37,38,253}$	$2\pi \approx 4 \times 355/226$
¹¹⁴ Fl ₁₇₅	19 23 28 31	$\alpha_{1-11,133,155} \alpha_{2-37,38}$	$2\pi \approx 17^2/46$
¹¹⁵ Mc ₁₇₃	20 21 25 31	$\alpha_{1-1,16,23} \alpha_{2-5}$	$2\pi \approx 17^2/46$
¹¹⁶ Lv ₁₇₇ ¹¹⁷ Ts ₁₇₇	10 20 22 27 31	$\alpha_{1-13,59} \alpha_{2-23} 1/\alpha_c^2$	$2\pi \approx 622/99$
¹¹⁸ Og ₁₇₆	20 22 23 27	$\alpha_{1-17,20,50,59,133} \alpha_{2-19,269}$	$2\pi \approx 44/7$
¹¹⁹⁻¹²² Ch ₁₇₉₋₁₈₂	21-23 28 31 37 39	$\alpha_{1-23,29,50,170} \alpha_{2-37} \alpha_{p/2} c_{au}$	$2\pi \approx 44/7$ et al.
¹²³ Ch _{183/185}	19 20 21 25 28	$\alpha_{1-4,11,16,17,31} \alpha_{2-1,4,5,9,32}$	$2\pi \approx 333/53 \approx 465/74$ et al

The relationships between elements and ideal extended elements (the frontier of elements) and an overall picture of them were depicted as above.


21. Some Supplements

Supplement 1:

$$2\pi \approx \frac{16 \cdot 3 \cdot 7 \cdot 11 \cdot 17}{100^2} = \frac{3 \cdot 7 \cdot 44 \cdot 68}{100^2} = \frac{3 \cdot 112 \cdot 11 \cdot 17}{100^2} = \frac{2 \cdot 168 \cdot 11 \cdot 17}{100^2} = \frac{2 \cdot 3 \cdot 7 \cdot 11 \cdot 136}{100^2} = \dots = 6.2832$$

Refer to Section 16; Supplements: ${}_{16}^{32,33,34}O_{16,17,18}$ and some of the follows


$$2\pi \approx \frac{9 \cdot 7 \cdot 127 \cdot (4 \cdot 13 \cdot 151 + 1)}{10^7} = \frac{63 \cdot 127 \cdot (52 \cdot 151 + 1)}{10 \cdot 100^3} = \frac{63 \cdot 127 \cdot (2 \cdot 3 \cdot 7 \cdot 11 \cdot 17 - 1)}{10 \cdot (8 \cdot 125)^2} = \dots = 6.2831853$$



2020/2/11-12


Supplement 2:

$$2\pi \approx \frac{509}{81} = \frac{4 \cdot 127 + 1}{9^2} \quad 2\pi \approx \frac{201}{32} = \frac{3 \cdot 67}{32} \quad 2\pi \approx \frac{333}{53} = \frac{9 \cdot 37}{53}$$



$$2\pi \approx \frac{622}{99} = \frac{4 \cdot (310 + 1)}{9 \cdot 22} \quad 2\pi \approx \frac{465}{74} = \frac{30 \cdot 31}{4 \cdot 37} \quad 2\pi \approx \frac{44}{7} = \frac{2 \cdot 22}{7}$$

$$2\pi \approx \frac{245}{39} = \frac{5 \cdot 7^2}{3 \cdot 13} \quad 2\pi \approx \frac{289}{46} = \frac{17^2}{2 \cdot 23} \quad 2\pi \approx \frac{377}{60} = \frac{13 \cdot 29}{4 \cdot 3 \cdot 5} \quad 2\pi \approx \frac{628}{100} = \frac{4 \cdot 157}{100}$$



2020/2/12


Supplement 3:

Table 8. Relationships of factors in α_{1-7} and α_{2-13} with primordial nuclides (2020/2/16-17).

Nuclides	${}_3Li_4$	${}_{29}Cu_{34}$	${}_{31}Ga_{40}$	${}_{64}Gd_{92}$	${}_{75}Re_{112}$
A	7	65=5×13	71	156=12×13	187=11×17
PN before	5	70	78	209	252
PN all	285	285	285	285	285
Ratios	1/57	14/57	26/95	11/15	84/95

- 3, 29, 31, 64, 75 and 112 are factors in α_1 and α_2 .
- PN: primordial nuclides; PN all: usually regarded as 286.
- Nucleon number 285 of ${}_{112}Cn_{173}$ would relate to PN all, or PN all should be 285 rather than 286, and ${}^{235}U$ should not be a primordial nuclide.
- ${}^{235}U_{143}$ should not be a primordial nuclide, its relative stability (but not much stable) should come from relative stable nucleon numbers 92=96-4 and 143=11×13, so number of PN would become 285 from 286.

Supplement 4: Correlations of factors in α (α_{1-7} , α_{2-13} and α_{1-50}) and nuclides



Relationships between Formulas of α (α_{1-7} , α_{2-13} and α_{1-50}) and Nuclides
Dr. Gang Chen, 2020 / 2 / 18–19

In this scheme there are several important clues based on factors in the formulas of α_{1-7} , α_{2-13} and α_{1-50} such as 6 (36, 48, 138, 144, *et al*), 7 (56, 84, 112, 126, 166-168, 210, 252), 10 (30, 50, 70, 100, 120, 130, 170, 200, 210, 220, 250, 310, 330, 400, 420), 11 (44, 88, 121, 134, 176, 187, 209, 220, 330, 363), 13 (26, 143, 169, 221, 364), 29 (87,145, 348), 25 (75, 100, 125, 200, 250, 400), 31 (93 124 186 310, 372), 61 (122, 244), 64 (136 et al), 137 (68, 69, 136,138), 139, 157 (314), 257, *et al*. And these clues correlate each others. These relationships are strong proofs that Chen’s formulas of the fine-structure constant are correct, otherwise so many coincidences couldn’t be explained.

In addition, numbers 7, 13 and 50 in α_{1-7} , α_{2-13} and α_{1-50} may have the following relationships: $(13+7)(13-7) = 50+70=120$ and ${}_{50}\text{Sn}_{70}$. And Sn is special, it has the most stable nuclides (up to 10) among which ${}_{50}\text{Sn}_{70}$ has the most relative abundance.

Supplement 5: Other two formulas of the fine-structure constant

$$\alpha_{1-9/11} = \frac{9}{11 + \frac{1}{84} - \frac{1}{11 \cdot 17 \cdot 53 + \frac{2 \cdot 191}{3 \cdot 157}}} \frac{1}{112 + \frac{1}{75^2}} = 1/137.035999037435$$

$$\alpha_{2-20/25} = \frac{20 + \frac{1}{2} - \frac{1}{14} + \frac{1}{251} - \frac{1}{8 \cdot (12 \cdot 43 \cdot 227 + 1) - \frac{8}{37}}}{25} \frac{1}{112 - \frac{1}{3 \cdot 29 \cdot 64}} = 1/137.035999111818$$

${}_{19}\text{F}_{10}$ ${}_{23}\text{Na}_{12}$ ${}_{35,37}\text{Cl}_{18,20}$ ${}_{53}\text{Cr}_{29}$ ${}_{75}\text{As}_{42}$ ${}_{84}\text{Kr}_{48}$ ${}_{95}\text{Mo}_{53}$ ${}_{127}\text{I}_{74}$ ${}_{129}\text{Xe}_{75}$ ${}_{144}\text{Nd}_{84}$
 ${}_{157}\text{Gd}_{93}$ ${}_{185,117}\text{Re}_{110,112}$ ${}_{191}\text{Y}_{114}$ ${}_{119}\text{Po}_{125}$ ${}_{257}\text{Fm}_{157}$ ${}_{7,37}\text{No}_{157}$ ${}_{6,53}\text{Ch}_{191}$ ${}_{400}\text{Ch}_{392}$
 ${}_{40,43,44,48}\text{Ca}_{20,23,24,28}$ ${}_{55}\text{Mn}_{30}$ ${}_{85,87}\text{Rb}_{48,50}$ ${}_{90,96}\text{Zr}_{50,56}$ ${}_{116,120,124}\text{Sn}_{66,70,74}$ ${}_{99}\text{Tc}_{56}$ ${}_{106,111,112,116}\text{Cd}_{58,63,64,68}$
 ${}_{191,193}\text{Y}_{114,4,29}$ ${}_{200}\text{Hg}_{120}$ ${}_{6,37}\text{Rn}_{136}$ ${}_{223}\text{Fr}_{136}$ ${}_{227}\text{Ac}_{138}$ ${}_{251}\text{Cf}_{153}$ ${}_{11,29}\text{Ch}_{127}$ ${}_{376}\text{Ch}_{149}$ ${}_{22,19}\text{Ch}_{167}$ ${}_{251}\text{Ch}_{251}$

Supplement 6: Other formulas of the speed of light c_{au}

$$c_{au} = \frac{1}{\sqrt{\alpha_{1-9/11}\alpha_{2-20/25}}}$$

$$= \sqrt{\frac{(11 + \frac{1}{84} - \frac{1}{11 \cdot 17 \cdot 53 + \frac{2 \cdot 191}{3 \cdot 157}})(112 + \frac{1}{75^2}) \cdot 25 \cdot (112 - \frac{1}{3 \cdot 29 \cdot 64})}{9 \cdot (20 + \frac{1}{2} - \frac{1}{14} + \frac{1}{251} - \frac{1}{8 \cdot (12 \cdot 43 \cdot 227 + 1) - \frac{8}{37}})}}$$

$$= \frac{5}{3} \sqrt{\frac{(11 + \frac{1}{84} - \frac{1}{11 \cdot 17 \cdot 53 + \frac{2 \cdot 191}{3 \cdot 157}})(112^2 - \frac{7 \cdot 19}{2^2 \cdot 3^2 \cdot 25^2 \cdot 29} - \frac{1}{2^6 \cdot 3^3 \cdot 25^2 \cdot 29})}{20 + \frac{1}{2} - \frac{1}{14} + \frac{1}{251} - \frac{1}{8 \cdot (12 \cdot 43 \cdot 227 + 1) - \frac{8}{37}}}}$$

$$= \frac{5}{3} \sqrt{\frac{(11 + \frac{1}{84} - \frac{1}{11 \cdot 17 \cdot 53 + \frac{2 \cdot 191}{3 \cdot 157}})(112^2 - \frac{1}{6 \cdot 17 \cdot 47 + \frac{2}{3}})}{20 + \frac{1}{2} - \frac{1}{14} + \frac{1}{251} - \frac{1}{8 \cdot (12 \cdot 43 \cdot 227 + 1) - \frac{8}{37}}}}$$

$$= \sqrt{137.035999037435 \times 137.035999111818} = 137.035999074627$$

²³ ₁₁ Cr ₁₂	²⁴⁻²⁶ ₁₂ Mg ₁₂₋₁₄	²⁸⁻³⁰ ₁₄ Si ₁₄₋₁₆	^{35,37} ₁₇ Cl _{18,20}	³⁹⁻⁴¹ ₁₉ K ₂₀₋₂₂	⁴⁷ ₂₂ Ti ₂₅	⁵³ ₂₄ Cr ₂₉	⁵⁵ ₂₅ Mn ₃₀	^{63,65} ₂₉ Cu _{34,36}	⁸⁴ ₃₆ Kr
⁸⁴ ₃₇ Rb [*]	^{85,87} ₃₇ Rb _{48,50}	^{90-92,94,96} ₄₀ Zr _{50-52,54,56}	⁹⁹ ₄₃ Tc [*]	^{107,109} ₄₇ Ag _{60,62}	¹²⁷ ₅₃ I ₇₄	⁷⁻¹⁹ ₅₅ Cs ₇₈	⁵⁻²⁹ ₆₁ Pm [*]	¹⁵⁷ ₆₄ Gd ₉₃	
^{3,53} ₆₅ Tb ₉₄	^{5,37,11,17} ₇₅ Re _{110,112}	^{191,193} ₇₇ Y _{114,4-29}	¹¹⁻¹⁹ ₈₄ Po [*]	²²³ ₈₇ Fr [*]	²²⁷ ₈₉ Ac [*]	²⁴⁴ ₉₄ Pu [*]	²⁵¹ ₉₈ Cf [*]	²⁵⁷ ₁₀₀ Fm [*]	
⁶⁻⁴³ ₁₀₁ Md [*]	^{7,37} ₁₀₂ No [*]	^{6,53,11-29,320} ₁₂₇ Ch ^{ie}	^{32,11,4-89} _{140,141} Ch ^{ie}	^{3,112} ₁₃₃ Ch ^{ie}	⁸⁻⁴⁷ ₁₄₉ Ch ^{ie}	⁴⁰⁰ ₁₅₇ Ch ^{ie}	²²¹⁹ ₁₆₇ Ch ^{ie}	²⁵¹ ₁₆₇ Ch ^{ie}	

2020 / 2 / 21 - 22

$$c_{au} = \frac{25 \cdot 112}{3} \sqrt{\frac{37}{11 \cdot 12 \cdot 13} - \frac{1}{2 \cdot 17 \cdot 41 \cdot 163 + \frac{47}{6 \cdot 31}}} = 137.035999074628$$

²³ ₁₁ Na ₁₂	^{24,25} ₁₂ Mg _{12,13}	^{35,37} ₁₇ Cl _{18,20}	⁵⁵ ₂₅ Mn ₃₀	^{69,71} ₃₁ Ga _{38,40}	^{74,77,78,82} ₃₄ Se _{40,43,44,48}	⁸⁴ ₃₇ Rb [*]	^{85,87} ₃₇ Rb _{48,50}	⁹³ ₄₁ Nb ₅₂
¹⁰⁰ ₄₄ Ru ₅₆	^{107,109} ₄₇ Ag _{60,62}	¹¹² ₄₈ Cd ₆₄	^{112,114-120,122,124} ₅₀ Sn _{62,64-70,72,74}	^{144,147,148,150,154} ₆₂ Sm _{82,85,86,88,92}	^{157,158} ₆₄ Gd _{93,94}			
¹⁶³ ₆₆ Dy ₉₇	¹⁶⁸ ₆₈ Er ₁₀₀	¹⁶⁹ ₆₉ Tm ₁₀₀	^{5,37,11,17} ₇₅ Re _{110,112}	^{204,206-1613} ₈₂ Pb _{122,124-126}	²³⁷ ₉₃ Np [*]	²⁴⁷ ₉₇ Bk [*]	²⁵⁷ ₁₀₀ Fm [*]	²⁶⁸ ₁₀₅ Db [*]
²⁶⁹ ₁₀₆ Sg [*]	²⁷⁰ ₁₀₇ Bh [*]	²⁸⁵ ₁₁₂ Cn [*]	³¹⁰ ₁₂₄ Ch ^{ie}	³³⁴ ₁₃₂ Ch ^{ie}	^{3,112,2813} ₁₄₃ Ch ^{ie}	³⁷⁸ ₁₅₀ Ch ^{ie}	³⁹⁴ ₁₅₆ Ch ^{ie}	⁴¹⁰ ₁₆₃ Ch ^{ie}

Note: $112 \times 5/3 \approx 187 = 11 \times 17$, $112 \times 25/3 \approx 5 \times 11 \times 17$

2020 / 2 / 24

$$c_{au} = \frac{25 \cdot 112}{3} \sqrt{\frac{1}{47} + \frac{1}{40 \cdot 89} - \frac{1}{16 \cdot 9 \cdot (2 \cdot 21 \cdot 31 \cdot 43 + 1) + \frac{3}{4}}} = 137.035999074627$$

⁴⁵ ₂₁ Ti ₂₄	⁴⁷ ₂₂ Ti ₂₅	⁵⁵ ₂₅ Mn ₃₀	^{64,66,70} ₃₀ Zn _{34,36,40}	^{69,71} ₃₁ Ga _{38,40}	⁷² ₃₂ Ge ₄₀	^{78,80,83,84,86} ₃₆ Kr _{42,44,47,48,50}	⁸⁹ ₃₉ Y ₅₀	^{90,94,96} ₄₀ Zr _{50,54,56}
^{92,94-98,100} ₄₂ Mo _{50,52-56,58}	^{98,99} ₄₃ Tc [*]	^{107,109} ₄₇ Ag _{60,62}	¹¹² ₄₈ Cd ₆₄	^{144,147,148,150,152} ₆₂ Sm _{82,85,86,88,90}	^{151,153} ₆₃ Eu _{88,90}	¹⁷⁸ ₇₂ Hf ₁₀₆		
^{185,187} ₇₅ Re _{110,112}	²²² ₈₆ Rn [*]	²²⁷ ₈₉ Ac [*]	²³⁷ ₉₃ Np [*]	²⁴⁴ ₉₄ Pu [*]	^{326,328} ₁₂₉ Ch ^{ie}	³⁶⁶ ₁₄₄ Ch ^{ie}	⁹⁻⁴² ₁₅₀ Ch ^{ie}	

$$c_{au} = \frac{25 \cdot 112}{3} \sqrt{\frac{1}{46} - \frac{1}{55 \cdot 100} + \frac{1}{9 \cdot 25 \cdot 13 \cdot (20 \cdot 293 + 1) - \frac{4}{23}}} = 137.035999074627$$

^{50,51} ₂₃ Mn _{27,28}	⁵⁵ ₂₅ Mn ₃₀	⁸⁹ ₃₉ Y ₅₀	^{99,100} ₄₄ Ru _{55,56}	^{106,110} ₄₆ Pd _{60,64}	¹¹⁷ ₅₀ Sn ₆₇	¹³³ ₅₅ Cs ₇₈	¹⁶⁹ ₆₉ Tm ₁₀₀	^{185,187} ₇₅ Re _{110,112}	¹⁹⁵ ₇₈ Pd ₁₁₇
^{5,47,238} ₉₂ U [*]	²⁵⁷ ₁₀₀ Fm [*]	²⁸⁵ ₁₁₂ Cn [*]	²⁹³ ₁₁₆ Lv ^{ie}	²⁹⁴ ₁₁₇ Ts ^{ie}	⁴⁰⁰ ₁₅₇ Ch ^{ie}	⁴²⁶ ₁₆₉ Ch ^{ie}			

2020 / 2 / 25

Supplement 7: Comparison of formulas of 1, N, e, 2π , $\pi/2$, ϕ , α , α_c , c_{au} and $\alpha_{p/c}$

$$1 = 4\gamma_c + \frac{4\gamma_1}{1(1+1)} + \frac{4\gamma_2}{2(2+1)} + \frac{4\gamma_3}{3(3+1)} + \dots$$

$$= |B| \frac{\pi}{2} + \sum_{n=1}^{\infty} \frac{|B_{2n}| (\pi/2)^{2n}}{(2n)!} = \sum_{n=1}^{\infty} \frac{|B_{2n}| \pi^{2n}}{(2n)!} = -|B| \frac{3\pi}{2} + \sum_{n=1}^{\infty} \frac{|B_{2n}| (3\pi/2)^{2n}}{(2n)!}$$

$$N \sim -\frac{3}{2}|B| + \sum_{n=1}^N \frac{|B_{2n}| (2\pi)^{2n}}{2(2n)!}$$

$$e = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \dots$$

$$2\pi = \left(\frac{e}{e^{\gamma_c}}\right)^2 = e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \frac{e^2}{\left(\frac{4}{3}\right)^7} \dots, \quad \frac{\pi}{2} = \left(\frac{e}{e^{\gamma_s}}\right)^2 = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots$$

$$2\pi \approx \frac{4 \cdot 157}{100} \approx \frac{9 \cdot 37}{53} \approx \frac{4 \cdot 5 \cdot 71}{15^2 + 1} \approx \dots, \quad \frac{\pi}{2} \approx \frac{157}{25} \approx \frac{9(9+1/4)}{53} \approx \frac{5 \cdot 71}{15^2 + 1} \approx \dots$$

$$\phi_1 = \frac{\sqrt{5}-1}{2} = 0.618\dots, \quad \phi_2 = -\frac{\sqrt{5}+1}{2} = -1.618\dots$$

$$\sqrt{\frac{\sqrt{5}+1}{2} + 2} - \frac{\sqrt{5}+1}{2} = \frac{e^{-\frac{2\pi}{5}}}{1 + \frac{e^{-2\pi}}{1 + \frac{e^{-4\pi}}{1 + \frac{e^{-6\pi}}{1 + \dots}}}} \quad (\text{Ramanujan Formula})$$

$$\alpha_1 = \frac{6^2}{7 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{113}{112}\right)^{225}}} \frac{1}{112 + \frac{1}{75^2}} = 1/137.035999037435$$

$$\alpha_2 = \frac{13 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{9 \cdot 31}{278}\right)^{557}}}{10^2} \frac{1}{112 - \frac{1}{3 \cdot 29 \cdot 64}} = 1/137.0359991118181$$

$$\frac{1}{\alpha_c^2} = \frac{1}{\alpha_1 \alpha_2} = 112 \times \left(168 - \frac{1}{3} + \frac{1}{12 \cdot 47} - \frac{1}{6 \cdot 29 \cdot 53 \cdot 59 - 79/47}\right)$$

$$= 137.035999074627^2 = 18778.865042381$$

$$c_{au} = \frac{1}{\alpha_c} = \frac{1}{\sqrt{\alpha_1 \alpha_2}} = \frac{5}{3} \sqrt{\frac{7 (2\pi)_{112}}{13 (2\pi)_{278}} \left(112^2 - \frac{1}{30^2 \cdot 5} + \frac{1}{60^2 \cdot 15} - \frac{1}{120^2 \cdot 15 \cdot 29}\right)}$$

$$= \frac{25 \cdot 112}{3} \sqrt{\frac{1}{47} + \frac{1}{40 \cdot 89} - \frac{1}{16 \cdot 9 \cdot (2 \cdot 21 \cdot 31 \cdot 43 + 1)} + \frac{3}{4}}$$


$$= \frac{25 \cdot 112}{3} \sqrt{\frac{1}{46} - \frac{1}{55 \cdot 100} + \frac{1}{9 \cdot 25 \cdot 13 \cdot (20 \cdot 293 + 1)} - \frac{4}{23}} = \dots = 137.035999074627$$

$$\frac{1}{\alpha_{p/c}^2} = \frac{1}{\alpha_{p/1} \alpha_{p/2}} = 252.040872632515^2 = 63524.60147736 \quad (\text{Supposed})$$

Ramanujan's formulas, so ϕ should connect to nuclides. And Fibonacci sequences are the integer presentations of ϕ , so Fibonacci sequences should connect to nuclides.


These connections are described as follows.

Fibonacci Sequence p_1 : 1 8 9 17 26 43 69 112




Note: $\begin{matrix} 56 \\ 26 \end{matrix} \text{Fe}_{30}$ $\begin{matrix} 100 \\ 43 \end{matrix} \text{Tc}_{56}^*$ $\begin{matrix} 169 \\ 69 \end{matrix} \text{Tm}_{100}$, relay of the numbers 56 and 100.

Fibonacci Sequence p_2 : 1 10 11 21 32 53 85 138




Note: $\begin{matrix} 74 \\ 32 \end{matrix} \text{Ge}_{42}$ $\begin{matrix} 127 \\ 53 \end{matrix} \text{I}_{74}$ $\begin{matrix} 210 \\ 85 \end{matrix} \text{At}_{125}^*$ $\begin{matrix} 12\cdot 29 \\ 138 \end{matrix} \text{Fy}_{210}^{ie}$, relay of the numbers 29, 74 and 210.

Fibonacci Sequence n_1 : 0 5 5 10 15 25 40 65 105 170




Note: $\begin{matrix} 170 \\ 68=4\cdot 17 \end{matrix} \text{Er}_{102=6\cdot 17}$ $\begin{matrix} 170,171 \\ 70 \end{matrix} \text{Yb}_{100=4\cdot 25,101}$ $\begin{matrix} 420 \\ 170 \end{matrix} \text{Ch}_{2\cdot 105}^{ie}$, relay of 21, 71 and 113.

Fibonacci Sequence n_2 : 0 4 4 8 12 20 32 52 84 136



Note: relay of the numbers 24, 41, 73, 125 and 209.

Fibonacci Sequence e : -1 9 8 17 25 42 67 109 176 285



Note: $\begin{matrix} 117 \\ 50=2\cdot 25 \end{matrix} \text{Sn}_{67}$, Fibonacci Sequence e is less relevant to specific nuclides.

Numbers of primordial nuclides before $\begin{matrix} 6 \\ 5 \end{matrix} \text{C}$ $\begin{matrix} 11 \\ 10 \end{matrix} \text{B}_6$ $\begin{matrix} 10 \\ 14 \end{matrix} \text{Ne}$ $\begin{matrix} 14 \\ 20 \end{matrix} \text{Si}$ $\begin{matrix} 42 \\ 28 \end{matrix} \text{Ca}_{22}$ $\begin{matrix} 61 \\ 28 \end{matrix} \text{Ni}_{33}$

$\begin{matrix} 94 \\ 40 \end{matrix} \text{Zr}_{54}$ $\begin{matrix} 56 \\ 56 \end{matrix} \text{Ba}$ and $\begin{matrix} 112 \\ 112 \end{matrix} \text{Cn}^*$ are 9 8 17 25 42 67 109 176 and 285

2020 / 2 / 27 - 28, 3 / 3 (add Fibonacci Sequence p_2 and n_2)

As stated in **Section 4**, the mathematic expression of chirality is $\pm 2\pi$. There are 10 fingers in a pair of human hands, and there are 14 finger segments in a single hand. It was assumed by us that the numbers 10/20 and 28/56 stand for a pair of "hands" in the world of nuclides. So nuclides $\begin{matrix} 10 \\ 10 \end{matrix} \text{Ne}$ $\begin{matrix} 14 \\ 14 \end{matrix} \text{Si}$ $\begin{matrix} 20 \\ 20 \end{matrix} \text{Ca}$ $\begin{matrix} 28 \\ 28 \end{matrix} \text{Ni}$ $\begin{matrix} 40 \\ 40 \end{matrix} \text{Zr}$ $\begin{matrix} 56 \\ 56 \end{matrix} \text{Ba}$ and $\begin{matrix} 112 \\ 112 \end{matrix} \text{Cn}$ stand for two pairs of "hands" emerging gradually, and the numbers of primordial nuclides just before them are the numbers of Fibonacci Sequence e . This means chirality or $\pm 2\pi$ is the inner essence and ϕ or Fibonacci sequence is the outer expression of nuclides.

1014	56	112
${}_{5}\text{B}_6$ ${}_{10}\text{Ne}_{17}$ ${}_{14}\text{Si}_{25}$ ${}_{20}\text{Ca}_{42}$ ${}_{28}\text{Ni}_{67}$ ${}_{40}\text{Zr}_{109}$ ${}_{56}\text{Ba}_{176}$ ${}_{112}\text{Cn}_{285}$		
8	176	285

Natural End


Primordial Nuclides

Primordial nuclides (PN) before Ba take about 0.618 part of all (176/285)
 Numbers of PN before ${}_{6}\text{C}$ ${}_{5}\text{B}_6$ ${}_{10}\text{Ne}$ ${}_{14}\text{Si}$ ${}_{20}\text{Ca}_{22}$ ${}_{28}\text{Ni}_{33}$ ${}_{40}\text{Zr}_{54}$ ${}_{56}\text{Ba}$ and ${}_{112}\text{Cn}$ are
 9 8 17 25 42 67 109 176 and 285 which is Fibonacci Sequence e

Primordial Nuclides and Fibonacci Sequence e

Dr. Gang Chen (2018/1-3, 2020/2/29-3/1)

Fig. 12



The Integrated Picture of Nuclides and Fibonacci Sequences

The Nuclide Picture was taken from Wikipedia

Dr. Gang Chen (2018/1-3, 2020/3/1-3)

Fig. 13

Why are there two pairs of “hands” in the world of nuclides? This should be because a pair of “hands” takes the right “hand” as priority and the other pair takes left “hand” as priority. So, 5 10 14 20 28 40 56 112 could be defined to be Numbers of Chirality, they are connected to Fibonacci Sequence e in nuclides (**Fig 13**).

22. Discussion and Conclusion

Regarding the fine-structure constant, Richard Feynman said: “is it related to π or perhaps to the base of natural logarithms?”⁴ Our answer is that it relate to $2\pi-e$ formula. He also deduced that the maximum element should be the 137th element Fynmanium (Fy) based on the analyses of the electron line velocity of his ideal hydrogen-like atoms. Our answer is that the natural end of the elements is the 112th element Copernicium (Cn^{*}), but the elements could have some ideal extensions, and above all, the fine-structure constant does relate to elements.

So, based on the analyses of ideal and real natural maximum element, Chen’s Chirality and Poetry Model of Atomic Nuclei⁷ and $2\pi-e$ formula^{6,7,8}, we deduced two series of Chen’s formulas of the fine-structure constant which gave two values $\alpha_1=1/137.035999037435$ and $\alpha_2=1/137.035999111818$. The factors in the formulas are much coincident to nucleon numbers of some nuclides, this means the formulas should be correct (too many coincidences mean too few possibilities to be wrong, or too many coincidences imply science). And we indicate the reason of $\alpha \approx 1/137.036$ is that it’s almost the equal ratio factor between 112 and 168 (more precisely 168-1/3) which are the key stable numbers (magic numbers) in Chen’s Chirality and Poetry Model of Atomic Nuclei⁷.

With Chen’s formulas of the fine-structure constant, we predicted the nucleon numbers of all 119th to 170th ideal extended elements; we theoretically or mathematically calculated the speed of light in atomic units, i.e., $c_{au}=1/\alpha_c=1/(\alpha_1\alpha_2)^{1/2}=137.035999074627$; we deduced a concise Schrödinger-Chen equation of hydrogen atom which included α_1/α_2 factor; in analogy to α and its formulas, α_p (the second fine-structure constant) and its formulas were hypothesized, and the proton charge radius r_p was supposed to be 0.833027203 fm; in the end we discovered that the approximate rational numbers of 2π marvelously and directly related to nuclides. Based on these, a mathematic shell model of elements was established and a picture

of elements and ideal extended elements was depicted.

In their relations to nuclides, 2π formulas can only be certain approximate rational numbers and 2π -e formulas in Chen's formulas of the fine-structure constant can only take certain k values. So we also believe the two values of the fine-structure constant should be rational numbers with definite digits rather than irrational numbers with infinite digits, and actually the fine-structure constant is transformed to nucleon numbers of 136, 137 and 138 in the world of nuclides.

In a recent paper¹¹, physicist Nicolas Gisin commented that in 1920s there once was a debate between mathematicians David Hilbert and Luitzen Egbertus Jan Brouwer. Hilbert was promoting formalized mathematics, in which every real number with its infinite series of digits is a completed individual object. On the other side the Luitzen Egbertus Jan Brouwer was defending the view that each point on the line should be represented as a never-ending process that develops in time, a view known as intuitionistic mathematics. Hilbert and his supporters clearly won the debate. In consequence, formalized mathematics has been adopted as the language of physics. In the end of his paper, Nicolas Gisin said: "Physics can be as successful if built on intuitionistic mathematics, even if this breaks its marriage to determinism. Contrary to usual expectations, I bet that the next physical theory will not be even more abstract than quantum field theory, but might well be closer to human experience."

In this paper we adopted mathematical language like intuitionistic mathematics, but we go ahead even more. The formulas of 2π , 2π -e and the fine-structure constant consist of integer factors and relate to nucleon numbers of nuclides, and hence correlate with each others. So in this paper we may use super-intuitionistic mathematics or decoding methodology with features of multi-correlations of integer factors or rational numbers which relate to nucleon numbers of nuclides, and it seems it is the real language in the world of nuclides. As we know an atomic nucleus is a N -body system and chaos should be its real state, so it seems N -body chaos returns to integers. In overall, Leopold Kronecker's famous saying "God made the integers, all else is the work of man" should be correct in the world of nuclides or even in other fields of the real world. It seems an irrational number can only be a rational number to play roles in the real world.

"God is a pure mathematician!" declared British astronomer Sir James

Jeans(1877-1946). The physical Universe does seem to be organised around elegant mathematical relationships³. The fine-structure constant may be the most important number in physics. As it is dimensionless, it could be called the proportional ruler of the nature or the bridge of mathematics and physics. And we have successfully given reasonable and precise formulas of it. In some sense, we explain the bridge between mathematics and physics, or we may realize the unification of mathematics and physics. It seems we prove the saying “God is a pure mathematician”. At least, it seems that good mathematics means good physics, and some pure mathematical numbers do have scientific meanings.

References

1. Wikipedia. The fine-structure constant.
2. R. H. Parker *et al.*, *Science* **360**, 191-195 (2018).
3. P. Davies, *Cosmos* 67 - Feb-Mar 2016.
4. R. P. Feynman, *QED: The Strange Theory of Light and Matter*, Princeton University Press, ISBN 0-691-08388-6, p. 129 (1985).
5. B. Palais, *the Mathematical Intelligencer* 03, 2001, Vol. 23, pp. 7-8.
6. G. Chen and T. Chen, Copyright Registration, Chen’s Periodic Table of Elements and Natural Group Theory, *GuoZuoDengZi-2018-L-00472808*.
7. G. Chen, T. Chen and T. Chen, Copyright Registration, Chirality and Poetry Model of Atomic Nuclei, *GuoZuoDengZi-2018-L-00421847*.
8. G. Chen, T. Chen and T. Chen, Copyright Registration, Chen’s Theory of the Fine-structure Constant, *GuoZuoDengZi-2018-L-00547467*.
9. N. Bezginov *et al.*, *Science* November 06, 2019, Vol. 365, pp. 1007-1012.
10. W. Xiong *et al.*, *Nature* November 06 2019, Vol. 575, pp. 147-150.
11. N. Gisin, *Nature Physics*, <http://www.nature.com/articles/s41567-019-0748-5>.

Acknowledgements

The main author Dr. Gang Chen studied in the Department of Chemistry at Sichuan University from 1983 to 1987 (B. Sc.), in Institute of Chemistry of the Academy of Sciences of China from 1987-1990 (M. Sc. under the supervision of Prof. Rongben Zhang), in the Hong Kong Polytechnic University from 1999 to 2004 (PhD and research assistant under the supervision of Prof. Albert Sun-Chi Chan) and in Kyoto University from 2004 to 2005 (postdoctoral research

under the supervision of Prof. Tamio Hayashi).

Leeman Chemical (HK) Co., Ltd. and Jiangshu Leeman Chemical Co., Ltd. gave Dr. Gang Chen a full time employment from May 2005 to July 2017.

Yichang Huifu Silicon Material Co., Ltd. has been giving Dr. Gang Chen a part-time employment since Dec. 2018. Special thanks to Dr. Yuelin Wang of the company for his appreciation and financial support.

Thank Mr. Qiyu Yang who is a famous Chinese traditional poet and an amateur researcher in mathematics. He gave me his book “The Story of Infinity” in 2010, and inspired by this, I started to write my new theory about infinity sets and one of my discoveries was 2π -e formula (in 2013).

Thank Prof. Raymond Honfu Chan of Department of Mathematics of the Chinese University of Hong Kong who discussed with me about 2π -e formula through about a dozen emails in 2013.

Thank Prof. Shiqing Zhang of College of Mathematics of Sichuan University who encourages me to write, open and publish papers in several years.

Thank Ramanujan, the famous Indian mathematician. His story, the picture of his Goddess and the film about him gave me some inspirations in this work.

The main author Dr. Gang Chen is deeply grateful to the above universities, institutions and companies, and to all teachers, classmates, friends and colleagues for their appreciation, encouragement and help to this work.

Appendix I: Research History

Items	Page	Discover/Create	Revise/Supplement
2π -e formula	3	2013/4-12	
Formulas related to 2π -e formula	4	2013/4-12	
Preliminary applications of 2π -e formula and its related formulas	6	2013/4-12	
Chen’s Periodic Table of Elements and Natural Group Theory	6	2014-2017/12	
Chirality and Poetry Model of Atomic Nuclei	6	2017/12-2018/3	
Chen’s theory of the fine-structure constant	6	2018/4-6	2018/7-2020/1
Original Inspiration for Formulas of the Fine-structure Constant	6	2018/4/12	
Logical deduction of Chen’s formulas of the fine-structure constant	6 7	2018/4/12-24	
α_1 (α_{1-7})	6 7	2018/4/12	2018/4/20 (+1/75 ²)

$\alpha_2 (\alpha_{2-13})$	7	2018/4/24	2018/9/18-20 (280→278 <i>et al.</i>)
Calculation tables and diagrams of α	8	2018/4/12-24	2018/9/18-20
$\alpha_{1-(3/2)}$	9	2019/4/25	
$\alpha_{2-(3/2)}$	9	2019/4/25	
α_c^2	10	2018/6/8-9, 9/18-19; 2019/4/17-19	
$1/\alpha_c^2$	10	2019/12/14	
112/137≈137/168 <i>et al.</i>	11	2018/4-6	
$^{136,137,138}\text{Fy}_{208,209,210}$	11	2019/12-2020/1	
$^{125,126}\text{Ch}$, $^{144-149}\text{Ch}$, $^{153,154}\text{Ch}$, $^{155,156}\text{Ch}$, ^{157}Ch , ^{163}Ch , $^{164-168}\text{Ch}$, ^{169}Ch	11	2019/12-2020/1	
c_{au} formulas	12	2019/12/16	2020/1/5-8
The special 29 and 75 factors	12-15	2019/4/22-24	
α_1/α_2 in Schrödinger Equation of Hydrogen Atom	16	2018/4-6	2019/12/13
α_1/α_2	16	2019/8/28-29	
The two kinds of general formulas of α	17	2019/6/27	2019/7/2-3
Parameters and results of α_{1-m}	18	2019/7/2	
α_{1-1}	19	2019/7/2	
α_{1-2}	19	2019/6/26	
α_{1-3}	19	2019/5/26	2019/6/26
α_{1-4}	19	2019/6/27	
α_{1-5}	19	2019/6/27	
α_{1-6}	19	2019/6/27	
$\alpha_{1-7} (\alpha_1)$	19	2018/4/12	2018/4/20 (+1/75 ²)
α_{1-9}	20	2019/6/28	
α_{1-11}	20	2019/6/29	
α_{1-13}	20	2019/6/29	
α_{1-16}	20	2019/6/29-30	
α_{1-17}	20	2019/6/30	2019/10/29
α_{1-19}	20	2019/7/1	
α_{1-20}	20	2018/4-6	2019/6/26
α_{1-22}	21	2019/5/25	2019/12/12
α_{1-23}	21	2019/7/4	
α_{1-25}	21	2019/7/4	
α_{1-27}	21	2019/5/25-26	
α_{1-29}	21	2019/5/25	
α_{1-31}	21	2019/7/4	
α_{1-32}	21	2019/7/5	
α_{1-33}	22	2019/7/5	
α_{1-34}	22	2019/5/24	
α_{1-36}	22	2019/5/24	

α_{1-43}	22	2019/12/15	
α_{1-50}	22	2018/4-6	2019/6/26
α_{1-59}	22	2019/5/25	
α_{1-81}	22	2019/5/25-26	
α_{1-96}	23	2019/5/25-26	2019/12/29
α_{1-103}	23	2019/12/15	
α_{1-133}	23	2019/5/26	2019/12/29
α_{1-140}	23	2019/5/26	
α_{1-155}	23	2019/12/15	
α_{1-170}	23	2019/7/2	2019/7/9
<hr/>			
Parameters and results of $\alpha_{2-m'}$	24	2019/7/3	
α_{2-1}	25	2018/4-6	2019/5/26-27
α_{2-4}	25	2019/5/28	
α_{2-5}	25	2019/6/22	
α_{2-6}	25	2019/6/23	
α_{2-7}	25	2019/5/24	
α_{2-9}	25	2019/6/21	
α_{2-10}	26	2019/5/28	
α_{2-11}	26	2019/6/22	
$\alpha_{2-13} (\alpha_2)$	26	2018/4/24	2018/9/18-20 (280→278 <i>et al.</i>)
α_{2-15}	26	2019/6/19	
α_{2-17}	26	2019/6/24	
α_{2-18}	26	2019/6/24	
α_{2-19}	27	2019/6/24	
α_{2-23}	27	2019/6/23	
α_{2-24}	27	2019/6/23	
α_{2-25}	27	2019/6/24	
α_{2-27}	27	2019/6/21	
α_{2-29}	27	2019/6/20	
α_{2-31}	28	2019/7/6	
α_{2-32}	28	2019/7/6	
α_{2-33}	28	2019/7/6	
α_{2-36}	28	2019/6/25	
α_{2-37}	28	2019/7/7	
α_{2-38}	28	2019/7/7	
α_{2-125}	29	2019/5/25	2019/12/20
α_{2-253}	29	2019/7/3	2019/7/9
α_{2-269}	29	2019/7/7	2019/12/19
<hr/>			
$a_0/r_e, r_e; a_0/r_p, r_p$	30	2019/12/19-23	
<hr/>			
$\alpha_{p/1}$	30	2020/1/2	
<hr/>			
$\alpha_{p/2}$	30 31	2020/1/2-3	
<hr/>			

${}_{164}\text{Ch}_{252}$	31	2020/1/3	
Direct relationships of 2π with nuclides	31	2020/1/8-10	
Some correlations of formulas of 2π and α	32 33	2020/1/11-13	
Chen's Mathematic Shell Model of Nuclides	33	2020/1/12-13	
${}_{130,131}\text{Ch}_{200,201}$	20 22 26 27	2020/1/26-28	
${}_{119,120,121}\text{Ch}_{179,180,181}$	21 28 31	2020/1/28-29	2020/2/5 (add 121)
${}_{128,129}\text{Ch}_{198,197/199}$	23 26 31	2020/1/28-29	2020/1/31
${}_{139}\text{Ch}_{209}$	21 26 28	2020/1/29	
${}_{132,133}\text{Ch}_{202,203}$	21 23	2020/1/31	
${}_{169}\text{Ch}_{257}$	11 22 27	2020/1/29-30	
${}_{157}\text{Ch}_{243}$	11 20 21 22 31	2019/1/8	
${}_{158}\text{Ch}_{243}$	11 20 22 25 30 31	2019/1/31	
${}_{169}\text{Ch}_{257}$	11 22 26 27	2020/1/29-30	
${}_{134,135}\text{Ch}_{206,205}$	20	2020/1/31	
${}_{127}\text{Ch}_{191,193}$	27	2020/1/31	2020/2/1 (add 191)
${}_{150,151,152}\text{Fy}_{228,229,230}$	19 28	2020/2/2	
${}_{170}\text{Ch}_{250}$	23 26	2020/2/3	
Chen's Picture of Elements and Ideal Extended Elements	41	2018/1-3 2020/2/2-5	2020/2/12, 16, 17, 19, 22-24
$2\pi=62831853/10^7$	37	2020/2/11-12	
${}_{124}\text{Ch}_{186}$	37	2020/2/12	
Table 8	37	2020/2/16-17	
${}_{143}\text{Ch}_{220,221}$	36	2020/2/17	
Supplement 4	38	2020/2/18-21	
${}_{122}\text{Ch}_{182}$	22	2020/2/19	
Supplement 5: $\alpha_{1-9/11}$ $\alpha_{2-20/25}$	38	2020/2/21	
${}_{127}\text{Ch}_{192}$ ${}_{167}\text{Ch}_{251}$	38	2020/2/21	
Other Formulas of c_{au}	39	2020/2/21-22, 24-25	
${}_{140,141}\text{Ch}_{212,215}$	39	2020/2/22	
${}_{123}\text{Ch}_{183,185}$	36	2020/2/23	
${}_{159/161,160}\text{Ch}_{245,246}$	36	2020/2/23	
${}_{165}\text{Ch}_{255}$	36	2020/2/23	
${}_{162}\text{Ch}_{246}$	36	2020/2/23	
Supplement 7	40	2020/2/25-26	
Supplement 8	41	2020/2/26	
Supplement 9	42	2020/2/27-3/3	
Preparing this paper	1-50	2019/12/1-2020/3/3	