

# Garo to English School Dictionary and the Graphical law

Anindya Kumar Biswas\*

*Department of Physics;*

*North-Eastern Hill University,*

*Mawkynroh-Umshing, Shillong-793022.*

(Dated: September 23, 2025)

## Abstract

We study a Garo to English School Dictionary. We draw the natural logarithm of the number of entries, normalised, starting with a letter vs the natural logarithm of the rank of the letter, normalised. We conclude that the Dictionary can be characterised by  $BP(4, \beta H = 0.02)$  i.e. a magnetisation curve for the Bethe-Peierls approximation of the Ising model with four nearest neighbours with  $\beta H = 0.02$ .  $\beta$  is  $\frac{1}{k_B T}$  where, T is temperature, H is external magnetic field and  $k_B$  is the Boltzmann constant.

---

\* anindya@nehu.ac.in

## I. INTRODUCTION

Garo people comes from Garo Hills of Meghalaya, India. The people are referred to as Garos, their language is known as Garo. "Bachi?" meaning where, "cha.a ringa?" meaning to eat and drink, "da.al nama?" meaning "are you fine today?", are the commonest way of conversations among them. A common man smiles as he talks. They laugh profusely. There are twelve clans. Am.beng, A.we, etc; four surnames; plenty of middle names, each referring to one ancestral place in the Garo Hills. Each clan has their respective places. A.we comes from Rishibelpara. The dialects vary as one goes from one region to another. The Rishibelpara dialect is the "official" version. In this language, "Balgito" means a lily, "Miktoksi" means a white flowering small tree, "boka" means white, "salanti" means everyday, "salaram" means the east, "janera" means a mirror, "grit" means a sugarcane, "matcha" means a tiger, "matchu" means a bull.

In this article, we try to do a thorough study of magnetic field pattern behind a dictionary of the Garo language,[1]. We have started considering magnetic field pattern in [2], in the languages we converse with. We have studied there, a set of natural languages, [2] and have found existence of a magnetisation curve under each language. We have termed this phenomenon as graphical law. Then, we moved on to investigate into, [3], dictionaries of five disciplines of knowledge and found existence of a curve magnetisation under each discipline. This was followed by finding of the graphical law behind the bengali language,[4] and the basque language[5]. This was pursued by finding of the graphical law behind the Romanian language, [6], five more disciplines of knowledge, [7], Onsager core of Abor-Miri, Mising languages,[8], Onsager Core of Romanised Bengali language,[9], the graphical law behind the Little Oxford English Dictionary, [10], the Oxford Dictionary of Social Work and Social Care, [11] and the Visayan-English Dictionary, [12], respectively.

In our first paper, [2], we have studied the Garo to English School Dicionary,[1]. There we took resort to average counting i.e. finding an average number of words par page and multiplying by the number of pages corresponding to a letter we obtained the number of words starting with a letter. We deduced that the dictionary,[1], is characterised by  $BP(4,\beta H=0.01)$ . Here, in this paper we leave behind the approximate method. We count thoroughly, one by one each word. Moreover, we augment the analysis. But the conclusion is very close to that of [2]. We conclude here, that the dictionary can be characterised by

BP(4, $\beta H=0.02$ ). It is desirable that this analysis is to be followed for other Garo language dictionaries to find out whether the same conclusion one reaches to for all of those.

The planning of the paper is as follows. We give an introduction to the standard curves of magnetisation of Ising model in the section II. In the section III, we describe analysis of the entries of the Garo language, [1]. Sections IV, V are Acknowledgement and Bibliography respectively.

## II. MAGNETISATION

### A. Bragg-Williams approximation

Let us consider a coin. Let us toss it many times. Probability of getting head or, tale is half i.e. we will get head and tale equal number of times. If we attach value one to head, minus one to tale, the average value we obtain, after many tossing is zero. Instead let us consider a one-sided loaded coin, say on the head side. The probability of getting head is more than one half, getting tale is less than one-half. Average value, in this case, after many tossing we obtain is non-zero, the precise number depends on the loading. The loaded coin is like ferromagnet, the unloaded coin is like paramagnet, at zero external magnetic field. Average value we obtain is like magnetisation, loading is like coupling among the spins of the ferromagnetic units. Outcome of single coin toss is random, but average value we get after long sequence of tossing is fixed. This is long-range order. But if we take a small sequence of tossing, say, three consecutive tossing, the average value we obtain is not fixed, can be anything. There is no short-range order.

Let us consider a row of spins, one can imagine them as spears which can be vertically up or, down. Assume there is a long-range order with probability to get a spin up is two third. That would mean when we consider a long sequence of spins, two third of those are with spin up. Moreover, assign with each up spin a value one and a down spin a value minus one. Then total spin we obtain is one third. This value is referred to as the value of long-range order parameter. Now consider a short-range order existing which is identical with the long-range order. That would mean if we pick up any three consecutive spins, two will be up, one down. Bragg-Williams approximation means short-range order is identical with long-range order, applied to a lattice of spins, in general. Row of spins is a lattice of one

dimension.

Now let us imagine an arbitrary lattice, with each up spin assigned a value one and a down spin a value minus one, with an unspecified long-range order parameter defined as above by  $L = \frac{1}{N}\sum_i \sigma_i$ , where  $\sigma_i$  is  $i$ -th spin,  $N$  being total number of spins.  $L$  can vary from minus one to one.  $N = N_+ + N_-$ , where  $N_+$  is the number of up spins,  $N_-$  is the number of down spins.  $L = \frac{1}{N}(N_+ - N_-)$ . As a result,  $N_+ = \frac{N}{2}(1 + L)$  and  $N_- = \frac{N}{2}(1 - L)$ . Magnetisation or, net magnetic moment,  $M$  is  $\mu\sum_i \sigma_i$  or,  $\mu(N_+ - N_-)$  or,  $\mu NL$ ,  $M_{max} = \mu N$ .  $\frac{M}{M_{max}} = L$ .  $\frac{M}{M_{max}}$  is referred to as reduced magnetisation. Moreover, the Ising Hamiltonian,[13], for the lattice of spins, setting  $\mu$  to one, is  $-\epsilon\sum_{n.n}\sigma_i\sigma_j - H\sum_i \sigma_i$ , where n.n refers to nearest neighbour pairs. The difference  $\Delta E$  of energy if we flip an up spin to down spin is, [14],  $2\epsilon\gamma\bar{\sigma} + 2H$ , where  $\gamma$  is the number of nearest neighbours of a spin. According to Boltzmann principle,  $\frac{N_-}{N_+}$  equals  $exp(-\frac{\Delta E}{k_B T})$ , [15]. In the Bragg-Williams approximation,[16],  $\bar{\sigma} = L$ , considered in the thermal average sense. Consequently,

$$\ln \frac{1+L}{1-L} = 2 \frac{\gamma\epsilon L + H}{k_B T} = 2 \frac{L + \frac{H}{\gamma\epsilon}}{\frac{T}{\gamma\epsilon/k_B}} = 2 \frac{L + c}{\frac{T}{T_c}} \quad (1)$$

where,  $c = \frac{H}{\gamma\epsilon}$ ,  $T_c = \gamma\epsilon/k_B$ , [17].  $\frac{T}{T_c}$  is referred to as reduced temperature.

Plot of  $L$  vs  $\frac{T}{T_c}$  or, reduced magnetisation vs. reduced temperature is used as reference curve. In the presence of magnetic field,  $c \neq 0$ , the curve bulges outward. Bragg-Williams is a Mean Field approximation. This approximation holds when number of neighbours interacting with a site is very large, reducing the importance of local fluctuation or, local order, making the long-range order or, average degree of freedom as the only degree of freedom of the lattice. To have a feeling how this approximation leads to matching between experimental and Ising model prediction one can refer to FIG.12.12 of [14]. W. L. Bragg was a professor of Hans Bethe. Rudlof Peierls was a friend of Hans Bethe. At the suggestion of W. L. Bragg, Rudlof Peierls following Hans Bethe improved the approximation scheme, applying quasi-chemical method.

## **B. Bethe-peierls approximation in presence of four nearest neighbours, in absence of external magnetic field**

In the approximation scheme which is improvement over the Bragg-Williams, [13],[14],[15],[16],[17], due to Bethe-Peierls, [18], reduced magnetisation varies with reduced temperature, for  $\gamma$

neighbours, in absence of external magnetic field, as

$$\frac{\ln \frac{\gamma}{\gamma-2}}{\ln \frac{factor-1}{factor^{\frac{\gamma-1}{\gamma}} - factor^{\frac{1}{\gamma}}}} = \frac{T}{T_c}; factor = \frac{\frac{M}{M_{max}} + 1}{1 - \frac{M}{M_{max}}}. \quad (2)$$

$\ln \frac{\gamma}{\gamma-2}$  for four nearest neighbours i.e. for  $\gamma = 4$  is 0.693. For a snapshot of different kind of magnetisation curves for magnetic materials the reader is urged to give a google search "reduced magnetisation vs reduced temperature curve". In the following, we describe datas generated from the equation(1) and the equation(2) in the table, I, and curves of magnetisation plotted on the basis of those datas. BW stands for reduced temperature in Bragg-Williams approximation, calculated from the equation(1). BP(4) represents reduced temperature in the Bethe-Peierls approximation, for four nearest neighbours, computed from the equation(2). The data set is used to plot fig.1. Empty spaces in the table, I, mean corresponding point pairs were not used for plotting a line.

reduced temperature, $\frac{T}{T_c}$				$\frac{M}{M_{max}}$ ,
BW(c=0)	BW(c=0.005)	BW(c=0.01)	BP(4, $\beta H = 0$ )	reduced magnetisation
0	0	0	0	1
0.435	0.437	0.439	0.563	0.978
0.439	0.441	0.443	0.568	0.977
0.491	0.493	0.495	0.624	0.961
0.501	0.504	0.507	0.630	0.957
0.514	0.517	0.519	0.648	0.952
0.559	0.562	0.565	0.654	0.931
0.566	0.569	0.573	0.7	0.927
0.584	0.587	0.590	0.7	0.917
0.601	0.604	0.607	0.722	0.907
0.607	0.610	0.613	0.729	0.903
0.653	0.658	0.661	0.770	0.869
0.659	0.663	0.666	0.773	0.865
0.669	0.674	0.678	0.784	0.856
0.679	0.684	0.688	0.792	0.847
0.701	0.705	0.709	0.807	0.828
0.723	0.728	0.732	0.828	0.805
0.732	0.736	0.743	0.832	0.796
0.753	0.758	0.766	0.845	0.772
0.779	0.784	0.788	0.864	0.740
0.838	0.844	0.853	0.911	0.651
0.850	0.858	0.864	0.911	0.628
0.870	0.877	0.885	0.923	0.592
0.883	0.891	0.899	0.928	0.564
0.899	0.908	0.918		0.527
0.905	0.914	0.926	0.941	0.513
0.944	0.956	0.968	0.965	0.400
		0.985		0.350
		0.998		0.310
0.969	0.985		0.965	0.300
	0.998			0.250
0.987			1	0.200
0.997			1	0.100
1			1	0

TABLE I. Datas for Reduced temperature[ for the Bragg-Williams approximation, in the absence (BW(c=0)) and in the presence (BW(c=0.005), BW(c=0.01)) of magnetic field,  $c = 0$ ,  $c = \frac{H}{\gamma\epsilon} = 0.005$ ,  $c = \frac{H}{\gamma\epsilon} = 0.01$  respectively and in the Bethe-Peierls approximation, BP(4, $\beta H=0$ ), in the absence of magnetic field, for four nearest neighbours] vs reduced magnetisation. Reduced temperature data set( say, data set BW(c=0)) is drawn along the x-axis and the corresponding Reduced magnetisation data set is drawn along the y-axis. In gnuplot the command is plot ".dat" using 1:2 with line; 1 standing for x-axis and 2 standing for y-axis datas.[For example, for drawing BW(c=0), ".dat" file, say denoted as "0.dat", contains BW(c=0) data set in first column and reduced magnetisation data set in second column. Moreover, after (0.944,0.400), next pair of points will be (0.969,0.300), then (0.987,0.200), ...and so on in the "0.dat" file.]




FIG. 1. Reduced magnetisation vs reduced temperature curves, for the Bragg-Williams approximation, in the absence (BW( $c=0$ )) and in the presence (BW( $c=0.005$ ), BW( $c=0.01$ )) of magnetic field,  $c = 0$ ,  $c = \frac{H}{\gamma\epsilon} = 0.005$ ,  $c = \frac{H}{\gamma\epsilon} = 0.01$ , outwards; and in the Bethe-Peierls approximation, BP(4, $\beta H=0$ ), in the absence of magnetic field, for four nearest neighbours (outer in the top).

### C. Bethe-peierls approximation in presence of four nearest neighbours, in presence of external magnetic field

In the Bethe-Peierls approximation scheme , [18], reduced magnetisation varies with reduced temperature, for  $\gamma$  neighbours, in presence of external magnetic field, as

$$\frac{\ln \frac{\gamma}{\gamma-2}}{\ln \frac{e^{-\frac{2\beta H}{\gamma}} \text{factor}^{\frac{\gamma-1}{\gamma}}}{e^{-\frac{2\beta H}{\gamma}} \text{factor}^{\frac{\gamma-1}{\gamma}} - e^{-\frac{2\beta H}{\gamma}} \text{factor}^{\frac{1}{\gamma}}}} = \frac{T}{T_c}; \text{factor} = \frac{\frac{M}{M_{max}} + 1}{1 - \frac{M}{M_{max}}}. \quad (3)$$

Derivation of this formula ala [18] is given in the appendix of [7].

$\ln \frac{\gamma}{\gamma-2}$  for four nearest neighbours i.e. for  $\gamma = 4$  is 0.693. For four neighbours,

$$\frac{0.693}{\ln \frac{e^{-\frac{2\beta H}{\gamma}} \text{factor}^{\frac{\gamma-1}{\gamma}}}{e^{-\frac{2\beta H}{\gamma}} \text{factor}^{\frac{\gamma-1}{\gamma}} - e^{-\frac{2\beta H}{\gamma}} \text{factor}^{\frac{1}{\gamma}}}} = \frac{T}{T_c}; \text{factor} = \frac{\frac{M}{M_{max}} + 1}{1 - \frac{M}{M_{max}}}. \quad (4)$$

In the following, we describe datas in the table, II, generated from the equation(4) and curves of magnetisation plotted on the basis of those datas. BP(m=0.03) stands for reduced temperature in Bethe-Peierls approximation, for four nearest neighbours, in presence of a variable external magnetic field, H, such that  $\beta H = 0.06$ . calculated from the equation(4). BP(m=0.025) stands for reduced temperature in Bethe-Peierls approximation, for four nearest neighbours, in presence of a variable external magnetic field, H, such that  $\beta H = 0.05$ . calculated from the equation(4). BP(m=0.02) stands for reduced temperature in Bethe-Peierls approximation, for four nearest neighbours, in presence of a variable external magnetic field, H, such that  $\beta H = 0.04$ . calculated from the equation(4). BP(m=0.01) stands for reduced temperature in Bethe-Peierls approximation, for four nearest neighbours, in presence of a variable external magnetic field, H, such that  $\beta H = 0.02$ . calculated from the equation(4). BP(m=0.005) stands for reduced temperature in Bethe-Peierls approximation, for four nearest neighbours, in presence of a variable external magnetic field, H, such that  $\beta H = 0.01$ . calculated from the equation(4). The data set is used to plot fig.2. Empty spaces in the table, II, mean corresponding point pairs were not used for plotting a line.




FIG. 2. Reduced magnetisation vs reduced temperature curves for Bethe-Peierls approximation in presence of little external magnetic fields, for four nearest neighbours, with  $\beta H = 2m$ .

reduced temperature, $\frac{T}{T_c}$					$\frac{M}{M_{max}}$ ,
BP(m=0.03)	BP(m=0.025)	BP(m=0.02)	BP(m=0.01)	BP(m=0.005)	reduced magnetisation
0	0	0	0	0	1
0.583	0.580	0.577	0.572	0.569	0.978
0.587	0.584	0.581	0.575	0.572	0.977
0.647	0.643	0.639	0.632	0.628	0.961
0.657	0.653	0.649	0.641	0.637	0.957
0.671	0.667		0.654	0.650	0.952
	0.716			0.696	0.931
0.723	0.718	0.713	0.702	0.697	0.927
0.743	0.737	0.731	0.720	0.714	0.917
0.762	0.756	0.749	0.737	0.731	0.907
0.770	0.764	0.757	0.745	0.738	0.903
0.816	0.808	0.800	0.785	0.778	0.869
0.821	0.813	0.805	0.789	0.782	0.865
0.832	0.823	0.815	0.799	0.791	0.856
0.841	0.833	0.824	0.807	0.799	0.847
0.863	0.853	0.844	0.826	0.817	0.828
0.887	0.876	0.866	0.846	0.836	0.805
0.895	0.884	0.873	0.852	0.842	0.796
0.916	0.904	0.892	0.869	0.858	0.772
0.940	0.926	0.914	0.888	0.876	0.740
	0.929			0.877	0.735
	0.936			0.883	0.730
	0.944			0.889	0.720
	0.945				0.710
	0.955			0.897	0.700
	0.963			0.903	0.690
	0.973			0.910	0.680
				0.909	0.670
	0.993			0.925	0.650
		0.976	0.942		0.651
	1.00				0.640
		0.983	0.946	0.928	0.628
		1.00	0.963	0.943	0.592
			0.972	0.951	0.564
			0.990	0.967	0.527
				0.964	0.513
			1.00		0.500
				1.00	0.400
					0.300
					0.200
					0.100
					0

TABLE II. Bethe-Peierls approx. in presence of little external magnetic fields

### D. Onsager solution

At a temperature  $T$ , below a certain temperature called phase transition temperature,  $T_c$ , for the two dimensional Ising model in absence of external magnetic field i.e. for  $H$  equal to zero, the exact, unapproximated, Onsager solution gives reduced magnetisation as a function of reduced temperature as, [19], [20], [21], [18],

$$\frac{M}{M_{max}} = [1 - (\sinh \frac{0.8813736}{\frac{T}{T_c}})^{-4}]^{1/8}.$$

Graphically, the Onsager solution appears as in fig.3.




FIG. 3. Reduced magnetisation vs reduced temperature curves for exact solution of two dimensional Ising model, due to Onsager, in absence of external magnetic field

letter	A	B	C	D	E	F	G	H	I	J	K	L	M
number	1208	1456	725	683	105	0	1220	80	110	697	798	52	1261
splitting	1192+16	1435+21	695+30	653+30	105+0	0	1205+15	80+0	108+2	682+15	774+24	51+1	1234+27
letter	N	O	P	Q	R	S	T	U	V	W	X	Y	Z
number	562	266	618	0	1414	1374	528	128	0	428	0	0	0
splitting	530+32	265+1	611+7	0	1372+42	1340+34	520+8	116+12	0	422+6	0	0	0

TABLE III. Entries of the Garo to English Dictionary: the first row represents letters of the english alphabet in the serial order, the second row is the respective number of entries, the third row describes the splitting of entries.

### III. METHOD OF STUDY AND RESULTS

The Garo language written in English alphabet is composed of twenty letters. We count all the entries in the dictionary, [1], one by one from the beginning to the end, starting with different letters. This has been done in two steps for the dictionary. First, we have counted all entries initiating with A form the section for the letter A. The number is one thousand one hundred ninety two. Second, we have enlisted all entries initiating with A form the sections for the letters B, D,...,Z. Then we have removed from the list entries already appearing in the section belonging to A. Then we have counted the number of the entries in that list. The number is sixteen. As a result total number of words beginning with A is one thousand two hundred and eight. This exercise was then followed for B,C,..Z. The result is the following table, III. Highest number of entries, one thousand four hundred fifty six, starts with the letter B followed by words numbering one thousand four hundred fourteen beginning with R, one thousand three hundred seventy four with the letter S etc. To visualise we plot the number of entries against the respective letters in the figure fig.4. For the purpose of exploring graphical law, we assort the letters according to the number of words, in the descending order, denoted by  $f$  and the respective rank, [22], denoted by  $k$ .  $k$  is a positive integer starting from one. Moreover, we attach a limiting rank,  $k_{lim}$ , and a limiting number of words. The limiting rank is maximum rank plus one, here it is twenty




FIG. 4. Vertical axis is number of entries of the Garo to English school Dictionary,[1]. Horizontal axis is the letters of the English alphabet. Letters are represented by the sequence number in the alphabet.

one and the limiting number of words is one. As a result both  $\frac{\ln f}{\ln f_{max}}$  and  $\frac{\ln k}{\ln k_{lim}}$  varies from zero to one. Then we tabulate in the adjoining table, IV, and plot  $\frac{\ln f}{\ln f_{max}}$  against  $\frac{\ln k}{\ln k_{lim}}$  in the figure fig.5.

We then ignore the letter with the highest number of words, tabulate in the adjoining table, IV, and redo the plot, normalising the  $\ln f$ s with next-to-maximum  $\ln f_{nextmax}$ , and starting from  $k = 2$  in the figure fig.6. Normalising the  $\ln f$ s with next-to-next-to-maximum  $\ln f_{nextnextmax}$ , we tabulate in the adjoining table, IV, and starting from  $k = 3$  we draw in the figure fig.7. Normalising the  $\ln f$ s with next-to-next-to-next-to-maximum  $\ln f_{nextnextnextmax}$  we record in the adjoining table, IV, and plot starting from  $k = 4$  in the figure fig.8. Normalising the  $\ln f$ s with 4n-maximum  $\ln f_{4n-max}$  we record in the adjoining table, IV, and plot starting from  $k = 5$  in the figure fig.9. Normalising the  $\ln f$ s with 5n-maximum  $\ln f_{5n-max}$  we record in the adjoining table, IV, and plot starting from  $k = 6$  in the figure fig.10, with 6n-maximum  $\ln f_{6n-max}$  we record in the adjoining table, IV, and plot starting from  $k = 7$  in the figure fig.11.

k	lnk	lnk/lnk <sub>lim</sub>	f	lnf	lnf/lnf <sub>max</sub>	lnf/lnf <sub>nmax</sub>	lnf/lnf <sub>nnmax</sub>	lnf/lnf <sub>nnnmax</sub>	lnf/lnf <sub>nnnnmax</sub>	lnf/lnf <sub>nnnnnmax</sub>	lnf/lnf <sub>nnnnnnmax</sub>
1	0	0	1456	7.283	1	Blank	Blank	Blank	Blank	Blank	Blank
2	0.69	0.228	1414	7.254	0.996	1	Blank	Blank	Blank	Blank	Blank
3	1.10	0.361	1374	7.225	0.992	0.996	1	Blank	Blank	Blank	Blank
4	1.39	0.455	1261	7.140	0.980	0.984	0.988	1	Blank	Blank	Blank
5	1.61	0.528	1220	7.107	0.976	0.980	0.984	0.995	1	Blank	Blank
6	1.79	0.589	1208	7.097	0.974	0.978	0.982	0.994	0.999	1	Blank
7	1.95	0.639	798	6.682	0.917	0.921	0.925	0.936	0.940	0.942	1
8	2.08	0.683	725	6.586	0.904	0.908	0.912	0.922	0.922	0.928	0.986
9	2.20	0.722	697	6.547	0.899	0.903	0.906	0.917	0.921	0.923	0.980
10	2.30	0.756	683	6.526	0.896	0.900	0.903	0.914	0.918	0.920	0.977
11	2.40	0.788	618	6.426	0.882	0.886	0.889	0.900	0.904	0.905	0.962
12	2.48	0.816	562	6.332	0.869	0.873	0.876	0.887	0.891	0.892	0.948
13	2.56	0.842	528	6.269	0.861	0.864	0.868	0.878	0.882	0.883	0.938
14	2.64	0.867	428	6.059	0.832	0.835	0.839	0.849	0.853	0.854	0.907
15	2.71	0.889	266	5.583	0.767	0.770	0.773	0.782	0.786	0.787	0.836
16	2.77	0.911	128	4.852	0.666	0.669	0.672	0.680	0.683	0.684	0.726
17	2.83	0.930	110	4.700	0.645	0.648	0.651	0.658	0.661	0.662	0.703
18	2.89	0.949	105	4.654	0.639	0.642	0.644	0.652	0.655	0.656	0.696
19	2.94	0.967	80	4.382	0.602	0.604	0.607	0.614	0.617	0.617	0.656
20	3.00	0.984	52	3.951	0.542	0.545	0.547	0.553	0.556	0.557	0.591
21	3.05	1	1	0	0	0	0	0	0	0	0

TABLE IV. Entries of the Garo to English Dictionary: ranking, natural logarithm, normalisations




FIG. 5. Vertical axis is  $\frac{\ln f}{\ln f_{max}}$  and horizontal axis is  $\frac{\ln k}{\ln k_{lim}}$ . The + points represent the entries of the Garo language with the fit curve being Bethe-Peierls curve in presence of four nearest neighbours and little magnetic field,  $m = 0.01$  or,  $\beta H = 0.02$ . The uppermost curve is the Onsager solution.




FIG. 6. Vertical axis is  $\frac{\ln f}{\ln f_{next-max}}$  and horizontal axis is  $\frac{\ln k}{\ln k_{lim}}$ . The + points represent the entries of the Garo language with the fit curve being Bethe-Peierls curve in presence of four nearest neighbours and little magnetic field,  $m = 0.01$  or,  $\beta H = 0.02$ . The uppermost curve is the Onsager solution.




FIG. 7. Vertical axis is  $\frac{\ln f}{\ln f_{nn-max}}$  and horizontal axis is  $\frac{\ln k}{\ln k_{lim}}$ . The + points represent the entries of the Garo language with the fit curve being Bethe-Peierls curve in presence of four nearest neighbours and little magnetic field,  $m = 0.01$  or,  $\beta H = 0.02$ . The uppermost curve is the Onsager solution.




FIG. 8. Vertical axis is  $\frac{\ln f}{\ln f_{nnn-max}}$  and horizontal axis is  $\frac{\ln k}{\ln k_{lim}}$ . The + points represent the entries of the Garo language with the fit curve being Bethe-Peierls curve in presence of four nearest neighbours and little magnetic field,  $m = 0.02$  or,  $\beta H = 0.04$ . The uppermost curve is the Onsager solution.




FIG. 9. Vertical axis is  $\frac{\ln f}{\ln f_{4n-max}}$  and horizontal axis is  $\frac{\ln k}{\ln k_{lim}}$ . The + points represent the entries of the Garo language with the fit curve being Bethe-Peierls curve in presence of four nearest neighbours and little magnetic field,  $m = 0.02$  or,  $\beta H = 0.04$ . The uppermost curve is the Onsager solution.




FIG. 10. Vertical axis is  $\frac{\ln f}{\ln f_{5n-max}}$  and horizontal axis is  $\frac{\ln k}{\ln k_{lim}}$ . The + points represent the entries of the Garo language with the fit curve being Bethe-Peierls curve in presence of four nearest neighbours and little magnetic field,  $m = 0.02$  or,  $\beta H = 0.04$ . The uppermost curve is the Onsager solution.




FIG. 11. Vertical axis is  $\frac{\ln f}{\ln f_{nnnnn-\max}}$  and horizontal axis is  $\frac{\ln k}{\ln k_{lim}}$ . The + points represent the entries of the english language with the fit curve being Bethe-Peierls curve in presence of four nearest neighbours and little magnetic field,  $m = 0.05$  or,  $\beta H = 0.1$ . The uppermost curve is the Onsager solution. The points of the Garo language do not go over to Onsager's solution i.e. the Garo language as viewed through this dictionary does not have Onsager core.

1. *conclusion*

From the figures (fig.5-fig.11), we observe that behind the entries of the dictionary, [1], there is a magnetisation curve,  $BP(4, \beta H = 0.02)$ , in the Bethe-Peierls approximation with four nearest neighbours, in presence of little magnetic field,  $\beta H = 0.02$ .

Moreover, the associated correspondence with the Ising model is,

$$\frac{\ln f}{\ln f_{\text{maximum}}} \longleftrightarrow \frac{M}{M_{\text{max}}},$$

and

$$\ln k \longleftrightarrow T.$$

$k$  corresponds to temperature in an exponential scale, [23]. As temperature decreases, i.e.  $\ln k$  decreases,  $f$  increases. The letters which are recording higher entries compared to those which have lesser entries are at lower temperature. As the Garo language expands, the letters which get enriched more and more, fall at lower and lower temperatures. This is a manifestation of cooling effect as was first observed in [24] in another way.

#### IV. ACKNOWLEDGEMENT

We have used gnuplot for drawing the figures. The author would like to thank people whom he has come across on the way and students in the M.Sc classes, who have illuminated him about language, traditions, places of the Garo Hills.

#### V. BIBLIOGRAPHY

---

- [1] C. M. Sangma, School Dictionary, Garo to English, 2010, D. J. Publication, Ringrey, Tura-794001, West Garo Hills(Meghalaya), India.
- [2] Anindya Kumar Biswas, "Graphical Law beneath each written natural language", arXiv:1307.6235v3[physics.gen-ph]. A preliminary study of words of dictionaries of twenty six languages, more accurate study of words of dictionary of Chinese usage and all parts of speech of dictionary of Lakher(Mara) language and of verbs, adverbs and adjectives of dictionaries of six languages are included.
- [3] Anindya Kumar Biswas, "A discipline of knowledge and the graphical law", IJARPS Volume 1(4), p 21, 2014; viXra: 1908:0090[Linguistics].
- [4] Anindya Kumar Biswas, "Bengali language and Graphical law ", viXra: 1908:0090[Linguistics].
- [5] Anindya Kumar Biswas, "Basque language and the Graphical Law", viXra: 1908:0414[Linguistics].
- [6] Anindya Kumar Biswas, "Romanian language, the Graphical Law and More ", viXra: 1909:0071[Linguistics].
- [7] Anindya Kumar Biswas, "Discipline of knowledge and the graphical law, part II", viXra:1912.0243 [Condensed Matter],International Journal of Arts Humanities and Social Sciences Studies Volume 5 Issue 2 February 2020.
- [8] Anindya Kumar Biswas, "Onsager Core of Abor-Miri and Mising Languages", viXra: 2003.0343[Condensed Matter].
- [9] Anindya Kumar Biswas, "Bengali language, Romanisation and Onsager Core", viXra:

- 2003.0563[Linguistics].
- [10] Anindya Kumar Biswas, "Little Oxford English Dictionary and the Graphical Law", viXra: 2008.0041[Linguistics].
- [11] Anindya Kumar Biswas, "Oxford Dictionary Of Social Work and Social Care and the graphical law", viXra: 2008.0077[Condensed Matter].
- [12] Anindya Kumar Biswas, "Visayan-English Dictionary and the graphical law", viXra: 2009.0014[Linguistics].
- [13] E. Ising, Z.Physik 31,253(1925).
- [14] R. K. Pathria, Statistical Mechanics, p400-403, 1993 reprint, Pergamon Press,© 1972 R. K. Pathria.
- [15] C. Kittel, Introduction to Solid State Physics, p. 438, Fifth edition, thirteenth Wiley Eastern Reprint, May 1994, Wiley Eastern Limited, New Delhi, India.
- [16] W. L. Bragg and E. J. Williams, Proc. Roy. Soc. A, vol.145, p. 699(1934);
- [17] P. M. Chaikin and T. C. Lubensky, Principles of Condensed Matter Physics, p. 148, first edition, Cambridge University Press India Pvt. Ltd, New Delhi.
- [18] Kerson Huang, Statistical Mechanics, second edition, John Wiley and Sons(Asia) Pte Ltd.
- [19] S. M. Bhattacharjee and A. Khare, "Fifty Years of the Exact solution of the Two-dimensional Ising Model by Onsager", arXiv:cond-mat/9511003v2.
- [20] L. Onsager, Nuovo Cim. Supp.6(1949)261.
- [21] C. N. Yang, Phys. Rev. 85, 809(1952).
- [22] A. M. Gun, M. K. Gupta and B. Dasgupta, Fundamentals of Statistics Vol 1, Chapter 12, eighth edition, 2012, The World Press Private Limited, Kolkata.
- [23] Sonntag, Borgnakke and Van Wylen, Fundamentals of Thermodynamics, p206-207, fifth edition, John Wiley and Sons Inc.
- [24] Alexander M. Petersen, Joel N. Tenenbaum, Shlomo Havlin, H. Eugene Stanley, and Matjaž Perc, "Languages cool as they expand: Allometric scaling and the decreasing need for new words", Sci. Rep.2(2012) 943, arXiv:1212.2616v1. and references therein.