An elementary proof of $0.999 \cdots = 1$

O. Kurwa

April 23, 2021

1 Abstract

One of the properties distinguishing irrational and rational numbers is the uniqueness (or the lack thereof) of their decimal representations. For example, the numbers π and 1 can be used as specimens of this phenomenon, as π has precisely one expression as decimals, but $1=1.0=1.00=\ldots$ In this paper, we provide an elementary proof for the fact that $0.999\ldots$ is also a decimal representation of 1, using the Lebesgue measure.

2 Preliminaries

We will refer to these well-known [1] facts throughout this paper.

Lemma 2.1. The following are true:

- (1) Given any interval $A \subseteq \mathbb{R}$, its Lebesgue measure satisfies m(A) = 0 iff A has only countably many points.
- (2) The only intervals with countably many points are degenerate intervals (singletons) and the empty set. In particular, any non-vacuous open interval has positive measure. (3) If A is not degenerate, then $m(A) = \max(A) \min(A)$.

3 The main result

We are now ready to prove the

Theorem 3.1. $0.999 \cdots = 1$.

Proof. We will argue by contradiction. Assume that $0.999 \cdots < 1$. There is then an open interval $\emptyset \neq A \subseteq (0.999 \ldots, 1)$. By Lemma 2.1(1), m(A) > 0. Let $(a_n)_{n \in \mathbb{N}}$ be the sequence

$$a_0 = 0, a_{n+1} = a_n + 9 \cdot 10^{-(n+1)}.$$

Clearly, $a_n \to 0.999\dots$ and $1 - a_n = 10^{-n} \to 0$ when $n \to \infty$. Defining $A_n = (a_n, 1)$, we obtain a sequence

$$(0,1)\supset (a_1,1)\supset (a_2,1)\supset \cdots$$

of open intervals, such that $A \subseteq \bigcap_{n \in \mathbb{N}} (a_n, 1)$. However,

$$m(\bigcap_{n\in\mathbb{N}}(a_n,1)) = \lim_{n\to\infty}m((a_n,1)) = \lim_{n\to\infty}10^{-n} = 0,$$

by 2.1(3). It follows from 2.1(2) that the intersection consists of at most one element, but then $0 \le |A| \le 1$, contradicting m(A) > 0. Thus, we conclude that $0.999 \cdots = 1$.

4 References

[1] Obviously true.