

Concise Urdu to English Dictionary and The Graphical Law

Anindya Kumar Biswas*

Department of Physics;

North-Eastern Hill University,

Mawkynroh-Umshing, Shillong-793022.

(Dated: September 4, 2025)

Abstract

We study the entries of the Concise Urdu to English Dictionary compiled by M. Zaman, Naved Akhtar, 2015. We draw the natural logarithm of the number of entries, normalised, starting with a letter vs the natural logarithm of the rank of the letter, normalised. We conclude that the Dictionary can be characterised by BP(4, $\beta H = 0.04$) i.e. a magnetisation curve for the Bethe-Peierls approximation of the Ising model with four nearest neighbours with $\beta H = 0.04$. β is $\frac{1}{k_B T}$ where, T is temperature, H is external magnetic field and k_B is the tiny Boltzmann constant.

* anindya@nehu.ac.in

alif	be	pe	te(1)	te(2)	se	jim	che	he(1)	khe	dal	dāl	zāl	re(1)	re(2)	ze	zhe	sin
386+2080	2000	1368	1373	242	63	828	773	465	549	776	189	90	673	0	265	20	1003
shin	sad	zad	toe	zoe	ain	ghain	fe	qaf	kaf	gaf	lam	mlm	nun	vao	he(2)		ye
437	253	90	203	50	409	187	325	348	1100	582	409	3053	502	274	588	0	235

TABLE I. Concise Urdu to English Dictionary entries: the odd rows represent letters of the Urdu alphabet,[1], in the serial order; the even rows represent the number of entries of the Concise Urdu to English Dictionary[1].




FIG. 1. The vertical axis is the number of entries of the Concise Urdu to English Dictionary[1]. The horizontal axis is the letters of the Urdu alphabet. Letters are represented by the sequence number in the alphabet as it appears in the dictionary, [1].

I. INTRODUCTION

Urdu is the language of large number of people, in particular in the Indian subcontinent. We study a Urdu to English Dictionary. This is the Concise Urdu to English Dictionary compiled by M. Zaman, Naved Akhtar, 2015, [1]. We count one by one all the entries, in this Dictionary,[1]. The result is the table, I. To visualise we plot the number of entries against the respective letters in the dictionary sequence,[1], in the adjoining figure, fig.1.

Looking for the Graphical Law in this dictionary, we proceed narrating the development. We have started considering magnetic field pattern in [2], in the languages we converse with. We

have studied there, a set of natural languages, [2] and have found existence of a magnetisation curve under each language. We have termed this phenomenon as the Graphical Law. Then, we moved on to investigate into, [3], dictionaries of five disciplines of knowledge and found existence of a curve magnetisation under each discipline. This was followed by finding of the graphical law in references from [4] to [71].

The planning of the paper is as follows. In the next section, we describe the Graphical Law analysis of the entries of the Concise Urdu to English Dictionary[1]. The section III, we give an introduction to the standard curves of magnetisation of Ising model. The section IV is Acknowledgment. The last section is Bibliography.

II. THE GRAPHICAL LAW ANALYSIS

For the purpose of exploring graphical law, we assort the letters according to the number of entries, in the descending order, denoted by f and the respective rank, [72], denoted by k . k is a positive integer starting from one. Moreover, the minimum non-zero number of entries is twenty. Hence, we attach a limiting entry number one. The limiting rank is maximum rank plus one, here it is thirty three. As a result both $\frac{\ln f}{\ln f_{max}}$ and $\frac{\ln k}{\ln k_{lim}}$ varies from zero to one. Then we tabulate in the adjoining table,II, and plot $\frac{\ln f}{\ln f_{max}}$ against $\frac{\ln k}{\ln k_{lim}}$ in the figure fig.2. We then ignore the letter with the highest of entries, tabulate in the adjoining table,II,and redo the plot, normalising the $\ln f$ s with $\ln f_{n-max}$, and starting from $k = 2$ in the figure fig.3. Normalising the $\ln f$ s with $\ln f_{2n-max}$, we tabulate in the adjoining table,II, and starting from $k = 3$ we draw in the figure fig.4. Normalising the $\ln f$ s with $\ln f_{3n-max}$ we record in the adjoining table,II, and plot starting from $k = 4$ in the figure fig.5. In this way we obtain up to the figure fig.7.

k	lnk	lnk/ lnk_{lim}	f	lnf	lnf/ lnf_{max}	lnf/ lnf_{n-max}	lnf/ lnf_{2n-max}	lnf/ lnf_{3n-max}	lnf/ lnf_{4n-max}	lnf/ lnf_{5n-max}
1	0	0	3053	8.024	1	Blank	Blank	Blank	Blank	Blank
2	0.69	0.197	2466	7.810	0.973	1	Blank	Blank	Blank	Blank
3	1.10	0.314	2000	7.601	0.947	0.973	1	Blank	Blank	Blank
4	1.39	0.397	1373	7.225	0.900	0.925	0.951	1	Blank	Blank
5	1.61	0.460	1368	7.221	0.900	0.925	0.950	0.999	1	Blank
6	1.79	0.511	1100	7.003	0.873	0.897	0.921	0.969	0.970	1
7	1.95	0.557	1003	6.911	0.861	0.885	0.909	0.957	0.957	0.987
8	2.08	0.594	828	6.719	0.837	0.860	0.884	0.930	0.930	0.959
9	2.20	0.629	776	6.654	0.829	0.852	0.875	0.921	0.921	0.950
10	2.30	0.657	773	6.650	0.829	0.851	0.875	0.920	0.921	0.950
11	2.40	0.686	673	6.512	0.812	0.834	0.857	0.901	0.902	0.930
12	2.48	0.709	588	6.377	0.795	0.817	0.839	0.883	0.883	0.911
13	2.56	0.731	582	6.366	0.793	0.815	0.838	0.881	0.882	0.909
14	2.64	0.754	549	6.308	0.786	0.808	0.830	0.873	0.874	0.901
15	2.71	0.774	502	6.219	0.775	0.796	0.818	0.861	0.861	0.888
16	2.77	0.791	465	6.142	0.765	0.786	0.808	0.850	0.851	0.877
17	2.83	0.809	437	6.080	0.758	0.778	0.800	0.842	0.842	0.868
18	2.89	0.826	409	6.014	0.750	0.770	0.791	0.832	0.833	0.859
19	2.94	0.840	348	5.852	0.729	0.749	0.770	0.810	0.810	0.836
20	3.00	0.857	325	5.784	0.721	0.741	0.761	0.801	0.801	0.826
21	3.04	0.869	274	5.613	0.700	0.719	0.738	0.777	0.777	0.802
22	3.09	0.883	265	5.580	0.695	0.714	0.734	0.772	0.773	0.797
23	3.14	0.897	253	5.533	0.690	0.708	0.728	0.766	0.766	0.790
24	3.18	0.909	242	5.489	0.684	0.703	0.722	0.760	0.760	0.784
25	3.22	0.920	235	5.460	0.680	0.699	0.718	0.756	0.756	0.780
26	3.26	0.931	203	5.313	0.662	0.680	0.699	0.735	0.736	0.759
27	3.30	0.943	189	5.242	0.653	0.671	0.690	0.726	0.726	0.749
28	3.33	0.951	187	5.231	0.652	0.670	0.688	0.724	0.724	0.747
29	3.37	0.963	90	4.500	0.561	0.576	0.592	0.623	0.623	0.643
30	3.40	0.971	63	4.143	0.516	0.530	0.545	0.573	0.574	0.592
31	3.43	0.980	50	3.912	0.488	0.501	0.515	0.541	0.542	0.559
32	3.47	0.991	20	2.996	0.373	0.384	0.394	0.415	0.415	0.428
33	3.50	1	1	0	0	0	0	0	0	0

TABLE II. Urdu to English Dictionary Entries: ranking,natural logarithm, normalisations




FIG. 2. The vertical axis is $\frac{\ln f}{\ln f_{max}}$ and the horizontal axis is $\frac{\ln k}{\ln k_{lim}}$. The + points represent the entries of the Concise Urdu to English Dictionary with the fit curve, BW($c=0.01$), being the Bragg-Williams curve in the presence of external magnetic field, $c = \frac{H}{\gamma\epsilon} = 0.01$.




FIG. 3. The vertical axis is $\frac{\ln f}{\ln f_{n-max}}$ and the horizontal axis is $\frac{\ln k}{\ln k_{lim}}$. The + points represent the entries of the Concise Urdu to English Dictionary with the fit curve, BP($4, \beta H = 0.01$), being the Bethe-Peierls curve in the presence of four nearest neighbours and external magnetic field, $m = 0.005$ or, $\beta H = 0.01$.




FIG. 4. The vertical axis is $\frac{\ln f}{\ln f_{2n-max}}$ and the horizontal axis is $\frac{\ln k}{\ln k_{lim}}$. The + points represent the entries of the Concise Urdu to English Dictionary with the fit curve, $BP(4, \beta H = 0.01)$, being the Bethe-Peierls curve in the presence of four nearest neighbours and external magnetic field, $m = 0.005$ or, $\beta H = 0.01$.




FIG. 5. The vertical axis is $\frac{\ln f}{\ln f_{3n-max}}$ and the horizontal axis is $\frac{\ln k}{\ln k_{lim}}$. The + points represent the entries of the Concise Urdu to English Dictionary with the fit curve, $BP(4, \beta H = 0.04)$, being the Bethe-Peierls curve in the presence of four nearest neighbours and external magnetic field, $m = 0.02$ or, $\beta H = 0.04$.




FIG. 6. The vertical axis is $\frac{\ln f}{\ln f_{4n-max}}$ and the horizontal axis is $\frac{\ln k}{\ln k_{lim}}$. The + points represent the entries of the Concise Urdu to English Dictionary with the fit curve, $BP(4, \beta H = 0.04)$, being the Bethe-Peierls curve in the presence of four nearest neighbours and external magnetic field, $m = 0.02$ or, $\beta H = 0.04$.




FIG. 7. The vertical axis is $\frac{\ln f}{\ln f_{5n-max}}$ and the horizontal axis is $\frac{\ln k}{\ln k_{lim}}$. The + points represent the entries of the Concise Urdu to English Dictionary with the fit curve, $BP(4, \beta H = 0.04)$, being the Bethe-Peierls curve in the presence of four nearest neighbours and external magnetic field, $m = 0.02$ or, $\beta H = 0.04$.

A. conclusion

From the figures (fig.2-fig.7), we observe that there is a curve of magnetisation, behind the entries of the Concise Urdu to English Dictionary,[1]. This is the magnetisation curve in the Bethe-Peierls approximation of the Ising model, in the presence of four nearest neighbours and external magnetic field, $BP(4, \beta H = 0.04)$.

Moreover, the associated correspondence is,

$$\frac{\ln f}{\ln f_{5n-max}} \longleftrightarrow \frac{M}{M_{max}},$$
$$\ln k \longleftrightarrow T.$$

k corresponds to temperature in an exponential scale, [79].

Though dispersion does reduce significantly over successive normalisations, probably, Spin-Glass magnetisation curve is well suited to describe the dictionary.

III. APENDIX: MAGNETISATION

A. Bragg-Williams approximation

Let us consider a coin. Let us toss it many times. Probability of getting head or, tale is half i.e. we will get head and tale equal number of times. If we attach value one to head, minus one to tale, the average value we obtain, after many tossing is zero. Instead let us consider a one-sided loaded coin, say on the head side. The probability of getting head is more than one half, getting tale is less than one-half. Average value, in this case, after many tossing we obtain is non-zero, the precise number depends on the loading. The loaded coin is like ferromagnet, the unloaded coin is like para magnet, at zero external magnetic field. Average value we obtain is like magnetisation, loading is like coupling among the spins of the ferromagnetic units. Outcome of single coin toss is random, but average value we get after long sequence of tossing is fixed. This is long-range order. But if we take a small sequence of tossing, say, three consecutive tossing, the average value we obtain is not fixed, can be anything. There is no short-range order.

Let us consider a row of spins, one can imagine them as spears which can be vertically up or, down. Assume there is a long-range order with probability to get a spin up is two third. That would mean when we consider a long sequence of spins, two third of those are with spin up. Moreover, assign with each up spin a value one and a down spin a value minus one. Then total spin we obtain is one third. This value is referred to as the value of long-range order parameter. Now consider a short-range order existing which is identical with the long-range order. That would mean if we pick up any three consecutive spins, two will be up, one down. Bragg-Williams approximation means short-range order is identical with long-range order, applied to a lattice of spins, in general. Row of spins is a lattice of one dimension.

Now let us imagine an arbitrary lattice, with each up spin assigned a value one and a down spin a value minus one, with an unspecified long-range order parameter defined as above by $L = \frac{1}{N}\sum_i\sigma_i$, where σ_i is i-th spin, N being total number of spins. L can vary from minus one to one. $N = N_+ + N_-$, where N_+ is the number of up spins, N_- is the number of down spins. $L = \frac{1}{N}(N_+ - N_-)$. As a result, $N_+ = \frac{N}{2}(1 + L)$ and $N_- = \frac{N}{2}(1 - L)$. Magnetisation or, net magnetic moment , M is $\mu\sum_i\sigma_i$ or, $\mu(N_+ - N_-)$ or, μNL , $M_{max} = \mu N$. $\frac{M}{M_{max}} = L$. $\frac{M}{M_{max}}$ is

referred to as reduced magnetisation. Moreover, the Ising Hamiltonian,[73], for the lattice of spins, setting μ to one, is $-\epsilon \sum_{n.n} \sigma_i \sigma_j - H \sum_i \sigma_i$, where n.n refers to nearest neighbour pairs. The difference ΔE of energy if we flip an up spin to down spin is, [74], $2\epsilon\gamma\bar{\sigma} + 2H$, where γ is the number of nearest neighbours of a spin. According to Boltzmann principle, $\frac{N_-}{N_+}$ equals $exp(-\frac{\Delta E}{k_B T})$, [75]. In the Bragg-Williams approximation,[76], $\bar{\sigma} = L$, considered in the thermal average sense. Consequently,

$$\ln \frac{1+L}{1-L} = 2 \frac{\gamma\epsilon L + H}{k_B T} = 2 \frac{L + \frac{H}{\gamma\epsilon}}{\frac{T}{\gamma\epsilon/k_B}} = 2 \frac{L + c}{\frac{T}{T_c}} \quad (1)$$

where, $c = \frac{H}{\gamma\epsilon}$, $T_c = \gamma\epsilon/k_B$, [77]. $\frac{T}{T_c}$ is referred to as reduced temperature.

Plot of L vs $\frac{T}{T_c}$ or, reduced magnetisation vs. reduced temperature is used as reference curve. In the presence of magnetic field, $c \neq 0$, the curve bulges outward. Bragg-Williams is a Mean Field approximation. This approximation holds when number of neighbours interacting with a site is very large, reducing the importance of local fluctuation or, local order, making the long-range order or, average degree of freedom as the only degree of freedom of the lattice. To have a feeling how this approximation leads to matching between experimental and Ising model prediction one can refer to FIG.12.12 of [74]. W. L. Bragg was a professor of Hans Bethe. Rudolf Peierls was a friend of Hans Bethe. At the suggestion of W. L. Bragg, Rudolf Peierls following Hans Bethe improved the approximation scheme, applying quasi-chemical method.

B. Bethe-peierls approximation in presence of four nearest neighbours, in absence of external magnetic field

In the approximation scheme which is improvement over the Bragg-Williams, [73],[74],[75],[76],[77], due to Bethe-Peierls, [78], reduced magnetisation varies with reduced temperature, for γ neighbours, in absence of external magnetic field, as

$$\frac{\ln \frac{\gamma}{\gamma-2}}{\ln \frac{factor-1}{factor^{\frac{\gamma-1}{\gamma}} - factor^{\frac{1}{\gamma}}}} = \frac{T}{T_c}; factor = \frac{\frac{M}{M_{max}} + 1}{1 - \frac{M}{M_{max}}} \quad (2)$$

$\ln \frac{\gamma}{\gamma-2}$ for four nearest neighbours i.e. for $\gamma = 4$ is 0.693. For a snapshot of different kind of magnetisation curves for magnetic materials the reader is urged to give a google search "reduced magnetisation vs reduced temperature curve". In the following, we describe




FIG. 8. Reduced magnetisation vs reduced temperature curves, for the Bragg-Williams approximation, in the absence (BW($c=0$)) and in the presence (BW($c=0.005$), BW($c=0.01$)) of magnetic field, $c = 0$, $c = \frac{H}{\gamma\epsilon} = 0.005$, $c = \frac{H}{\gamma\epsilon} = 0.01$, outwards; and in the Bethe-Peierls approximation, BP(4, $\beta H=0$), in the absence of magnetic field, for four nearest neighbours (outer in the top).

data s generated from the equation(1) and the equation(2) in the table, III, and curves of magnetisation plotted on the basis of those data s. BW stands for reduced temperature in Bragg-Williams approximation, calculated from the equation(1). BP(4) represents reduced temperature in the Bethe-Peierls approximation, for four nearest neighbours, computed from the equation(2). The data set is used to plot fig.8. Empty spaces in the table, III, mean corresponding point pairs were not used for plotting a line.

reduced temperature, $\frac{T}{T_c}$				$\frac{M}{M_{max}}$,
BW(c=0)	BW(c=0.005)	BW(c=0.01)	BP(4, $\beta H = 0$)	reduced magnetisation
0	0	0	0	1
0.435	0.437	0.439	0.563	0.978
0.439	0.441	0.443	0.568	0.977
0.491	0.493	0.495	0.624	0.961
0.501	0.504	0.507	0.630	0.957
0.514	0.517	0.519	0.648	0.952
0.559	0.562	0.565	0.654	0.931
0.566	0.569	0.573	0.7	0.927
0.584	0.587	0.590	0.7	0.917
0.601	0.604	0.607	0.722	0.907
0.607	0.610	0.613	0.729	0.903
0.653	0.658	0.661	0.770	0.869
0.659	0.663	0.666	0.773	0.865
0.669	0.674	0.678	0.784	0.856
0.679	0.684	0.688	0.792	0.847
0.701	0.705	0.709	0.807	0.828
0.723	0.728	0.732	0.828	0.805
0.732	0.736	0.743	0.832	0.796
0.753	0.758	0.766	0.845	0.772
0.779	0.784	0.788	0.864	0.740
0.838	0.844	0.853	0.911	0.651
0.850	0.858	0.864	0.911	0.628
0.870	0.877	0.885	0.923	0.592
0.883	0.891	0.899	0.928	0.564
0.899	0.908	0.918		0.527
0.905	0.914	0.926	0.941	0.513
0.944	0.956	0.968	0.965	0.400
		0.985		0.350
		0.998		0.310
0.969	0.985		0.965	0.300
	0.998			0.250
0.987			1	0.200
0.997			1	0.100
1			1	0

TABLE III. Datas for Reduced temperature[for the Bragg-Williams approximation, in the absence (BW(c=0)) and in the presence (BW(c=0.005), BW(c=0.01)) of magnetic field, $c = 0$, $c = \frac{H}{\gamma\epsilon} = 0.005$, $c = \frac{H}{\gamma\epsilon} = 0.01$ respectively and in the Bethe-Peierls approximation, BP(4, $\beta H=0$), in the absence of magnetic field, for four nearest neighbours] vs reduced magnetisation. Reduced temperature data set(say, data set BW(c=0)) is drawn along the x-axis and the corresponding Reduced magnetisation data set is drawn along the y-axis. In gnuplot the command is plot ".dat" using 1:2 with line; 1 standing for x-axis and 2 standing for y-axis datas.[For example, for drawing BW(c=0), ".dat" file, say denoted as "0.dat", contains BW(c=0) data set in first column and reduced magnetisation data set in second column. Moreover, after (0.944,0.400), next pair of points will be (0.969,0.300), then (0.987,0.200), .and so on in the "0.dat" file.]

C. Bethe-peierls approximation in presence of four nearest neighbours, in the presence of external magnetic field

In the Bethe-Peierls approximation scheme , [78], reduced magnetisation varies with reduced temperature, for γ neighbours, in presence of external magnetic field, as

$$\frac{\ln \frac{\gamma}{\gamma-2}}{\ln \frac{e^{-\frac{2\beta H}{\gamma}} \text{factor}^{\frac{\gamma-1}{\gamma}} - e^{-\frac{2\beta H}{\gamma}} \text{factor}^{\frac{1}{\gamma}}}{\text{factor} - 1}} = \frac{T}{T_c}; \text{factor} = \frac{\frac{M}{M_{max}} + 1}{1 - \frac{M}{M_{max}}}. \quad (3)$$

Derivation of this formula ala [78] is given in the appendix of [7].

$\ln \frac{\gamma}{\gamma-2}$ for four nearest neighbours i.e. for $\gamma = 4$ is 0.693. For four neighbours,

$$\frac{0.693}{\ln \frac{e^{-\frac{2\beta H}{\gamma}} \text{factor}^{\frac{\gamma-1}{\gamma}} - e^{-\frac{2\beta H}{\gamma}} \text{factor}^{\frac{1}{\gamma}}}{\text{factor} - 1}} = \frac{T}{T_c}; \text{factor} = \frac{\frac{M}{M_{max}} + 1}{1 - \frac{M}{M_{max}}}. \quad (4)$$

In the following, we describe datas in the table, IV, generated from the equation(4) and curves of magnetisation plotted on the basis of those datas. BP(m=0.03) stands for reduced temperature in Bethe-Peierls approximation, for four nearest neighbours, in presence of a variable external magnetic field, H, such that $\beta H = 0.06$. calculated from the equation(4). BP(m=0.025) stands for reduced temperature in Bethe-Peierls approximation, for four nearest neighbours, in presence of a variable external magnetic field, H, such that $\beta H = 0.05$. calculated from the equation(4). BP(m=0.02) stands for reduced temperature in Bethe-Peierls approximation, for four nearest neighbours, in presence of a variable external magnetic field, H, such that $\beta H = 0.04$. calculated from the equation(4). BP(m=0.01) stands for reduced temperature in Bethe-Peierls approximation, for four nearest neighbours, in presence of a variable external magnetic field, H, such that $\beta H = 0.02$. calculated from the equation(4). BP(m=0.005) stands for reduced temperature in Bethe-Peierls approximation, for four nearest neighbours, in presence of a variable external magnetic field, H, such that $\beta H = 0.01$. calculated from the equation(4). The data set is used to plot fig.9. Empty spaces in the table, IV, mean corresponding point pairs were not used for plotting a line.

reduced temperature, $\frac{k_B T}{J}$					$\frac{M}{M_{max}}$
BP(m=0.03)	BP(m=0.025)	BP(m=0.02)	BP(m=0.01)	BP(m=0.005)	reduced magnetisation
0	0	0	0	0	1
0.583	0.580	0.577	0.572	0.569	0.978
0.587	0.584	0.581	0.575	0.572	0.977
0.647	0.643	0.639	0.632	0.628	0.961
0.657	0.653	0.649	0.641	0.637	0.957
0.671	0.667		0.654	0.650	0.952
	0.716			0.696	0.931
0.723	0.718	0.713	0.702	0.697	0.927
0.743	0.737	0.731	0.720	0.714	0.917
0.762	0.756	0.749	0.737	0.731	0.907
0.770	0.764	0.757	0.745	0.738	0.903
0.816	0.808	0.800	0.785	0.778	0.869
0.821	0.813	0.805	0.789	0.782	0.865
0.832	0.823	0.815	0.799	0.791	0.856
0.841	0.833	0.824	0.807	0.799	0.847
0.863	0.853	0.844	0.826	0.817	0.828
0.887	0.876	0.866	0.846	0.836	0.805
0.895	0.884	0.873	0.852	0.842	0.796
0.916	0.904	0.892	0.869	0.858	0.772
0.940	0.926	0.914	0.888	0.876	0.740
	0.929			0.877	0.735
	0.936			0.883	0.730
	0.944			0.889	0.720
	0.945				0.710
	0.955			0.897	0.700
	0.963			0.903	0.690
	0.973			0.910	0.680
				0.909	0.670
	0.993			0.925	0.650
		0.976	0.942		0.651
	1.00				0.640
		0.983	0.946	0.928	0.628
		1.00	0.963	0.943	0.592
			0.972	0.951	0.564
			0.990	0.967	0.527
				0.964	0.513
			1.00		0.500
				1.00	0.400
					0.300
					0.200
					0.100
					0

TABLE IV. Bethe-Peierls approx. in presence of little external magnetic fields




FIG. 9. Reduced magnetisation vs reduced temperature curves for Bethe-Peierls approximation in presence of little external magnetic fields, for four nearest neighbours, with $\beta H = 2m$.

IV. ACKNOWLEDGMENT

We have used gnuplot for plotting the figures in this paper.

-
- [1] Concise Twenty First Century Dictionary, Urdu to English, compiled by M.Zaman, Naved Akhtar, 2015, Educational Publishing House, 3108, Vakil Street, Kucha Pandit, Lal Kuan, Delhi-6(India). ISBN 81-86232-51-6.
 - [2] Anindya Kumar Biswas, "Graphical Law beneath each written natural language", arXiv:1307.6235v3[physics.gen-ph]. A preliminary study of words of dictionaries of twenty six languages, more accurate study of words of dictionary of Chinese usage and all parts of speech of dictionary of Lakher(Mara) language and of verbs, adverbs and adjectives of dictionaries of six languages are included.
 - [3] Anindya Kumar Biswas, "A discipline of knowledge and the graphical law", IJARPS Volume 1(4), p 21, 2014; viXra: 1908:0090[Linguistics].
 - [4] Anindya Kumar Biswas, "Bengali language and Graphical law", viXra: 1908:0090[Linguistics].
 - [5] Anindya Kumar Biswas, "Basque language and the Graphical Law", viXra: 1908:0414[Linguistics].
 - [6] Anindya Kumar Biswas, "Romanian language, the Graphical Law and More", viXra: 1909:0071[Linguistics].
 - [7] Anindya Kumar Biswas, "Discipline of knowledge and the graphical law, part II", viXra:1912.0243 [Condensed Matter],International Journal of Arts Humanities and Social Sciences Studies Volume 5 Issue 2 February 2020.
 - [8] Anindya Kumar Biswas, "Onsager Core of Abor-Miri and Mising Languages", viXra: 2003.0343[Condensed Matter].
 - [9] Anindya Kumar Biswas, "Bengali language, Romanisation and Onsager Core", viXra: 2003.0563[Linguistics].
 - [10] Anindya Kumar Biswas, "Little Oxford English Dictionary and the Graphical Law", viXra: 2008.0041[Linguistics].
 - [11] Anindya Kumar Biswas, "Oxford Dictionary Of Social Work and Social Care and the Graphical law", viXra: 2008.0077[Condensed Matter].

- [12] Anindya Kumar Biswas, "Visayan-English Dictionary and the Graphical law", viXra: 2009.0014[Linguistics].
- [13] Anindya Kumar Biswas, "Garo to English School Dictionary and the Graphical law", viXra: 2009.0056[Condensed Matter].
- [14] Anindya Kumar Biswas, "Mursi-English-Amharic Dictionary and the Graphical law", viXra: 2009.0100[Linguistics].
- [15] Anindya Kumar Biswas, "Names of Minor Planets and the Graphical law", viXra: 2009.0158[History and Philosophy of Physics].
- [16] Anindya Kumar Biswas, "A Dictionary of Tibetan and English and the Graphical law", viXra: 2010.0237[Condensed Matter].
- [17] Anindya Kumar Biswas, "Khasi English Dictionary and the Graphical law", viXra: 2011.0011[Linguistics].
- [18] Anindya Kumar Biswas, "Turkmen-English Dictionary and the Graphical law", viXra: 2011.0069[Linguistics].
- [19] Anindya Kumar Biswas, " Webster's Universal Spanish-English Dictionary, the Graphical law and A Dictionary of Geography of Oxford University Press", viXra: 2103.0175[Condensed Matter].
- [20] Anindya Kumar Biswas, "A Dictionary of Modern Italian, the Graphical law and Dictionary of Law and Administration, 2000, National Law Development Foundation", viXra: 2107.0171[Condensed Matter].
- [21] Anindya Kumar Biswas, "Langenscheidt's German-English English-German Dictionary and the Graphical law", viXra: 2107.0179[Linguistics].
- [22] Anindya Kumar Biswas, "Essential Dutch dictionary by G. Quist and D. Strik, the Graphical law Classification", viXra: 2108.0040[Linguistics].
- [23] Anindya Kumar Biswas, "Swahili, a lingua franca, Swahili-English Dictionary by C. W. Rechenbach and the Graphical law", viXra: 2108.0101[Linguistics].
- [24] Anindya Kumar Biswas, "The French, Larousse Dictionnaire De Poche and the Graphical law", viXra: 2109.0080[Linguistics].
- [25] Anindya Kumar Biswas, "An Arabic dictionary: "al-Mujam al-wáfi" or, "adhunik arabi-bangla abhidhan" and the Onsager's solution", viXra: 2109.0119[Condensed Matter].

- [26] Anindya Kumar Biswas, "Langenscheidt Taschenwörterbuch Deutsch-Englisch / Englisch-Deutsch, Völlige Neubearbeitung and the Graphical law", viXra: 2109.0141[Linguistics].
- [27] Anindya Kumar Biswas, Bawansuk Lyngkhoi, "The Graphical law behind the NTC's Hebrew and English Dictionary by Arie Comey and Naomi Tsur", viXra: 2109.0164[Linguistics].
- [28] Anindya Kumar Biswas, "Oxford Dictionary Of Media and Communication and the Graphical law", viXra: 2109.0202[Social Science].
- [29] Anindya Kumar Biswas, "Oxford Concise Dictionary Of Mathematics, Penguin Dictionary Of Mathematics and the Graphical law", viXra: 2112.0054[Social Science].
- [30] Anindya Kumar Biswas, "An Arabic dictionary: "al-Mujam al-wáfi" or, "adhunik arabi-bangla abhidhan" and the Onsager's solution Second part", viXra: 2201.0021[Condensed Matter].
- [31] Anindya Kumar Biswas, "The Penguin Dictionary Of Sociology and the Graphical law", viXra: 2201.0046[Social Science].
- [32] Anindya Kumar Biswas, "The Concise Oxford Dictionary Of Politics and the Graphical law", viXra: 2201.0069[Social Science].
- [33] Anindya Kumar Biswas, "A Dictionary Of Critical Theory by Ian Buchanan and the Graphical law", viXra: 2201.0136[Social Science].
- [34] Anindya Kumar Biswas, "The Penguin Dictionary Of Economics and the Graphical law", viXra: 2201.0169[Economics and Finance].
- [35] Anindya Kumar Biswas, "The Concise Gojri-English Dictionary by Dr. Rafeeq Anjum and the Graphical law", viXra: 2201.0205[Linguistics].
- [36] Anindya Kumar Biswas, "A Dictionary of the Kachin Language by Rev.O.Hanson and the Graphical law" ("A Dictionary of the Kachin Language by Rev.o.Hanson and the Graphical law", viXra: 2202.0030[Linguistics]).
- [37] Anindya Kumar Biswas, "A Dictionary Of World History by Edmund Wright and the Graphical law", viXra: 2202.0130[History and Philosophy of Physics].
- [38] Anindya Kumar Biswas, "Ekagi-Dutch-English-Indonesian Dictionary by J. Steltenpool and the Onsager's solution", viXra: 2202.0157[Condensed Matter].
- [39] Anindya Kumar Biswas, "A Dictionary of Plant Sciences by Michael Allaby and the Graphical law", viXra: 2203.0011[Mind Science].
- [40] Anindya Kumar Biswas, "Along the side of the Onsager's solution, the Ekagi language", viXra: 2205.0065[Condensed Matter].

- [41] Anindya Kumar Biswas, "Along the side of the Onsager's solution, the Ekagi language-Part Three", viXra: 2205.0137[Condensed Matter].
- [42] Anindya Kumar Biswas, "Oxford Dictionary of Biology by Robert S. Hine and the Graphical law", viXra: 2207.0089[Physics of Biology].
- [43] Anindya Kumar Biswas, "A Dictionary of the Mikir Language by G. D. Walker and the Graphical law", viXra: 2207.0165[Linguistics].
- [44] Anindya Kumar Biswas, "A Dictionary of Zoology by Michael Allaby and the Graphical law", viXra: 2208.0075[Physics of Biology].
- [45] Anindya Kumar Biswas, "Dictionary of all Scriptures and Myths by G. A. Gaskell and the Graphical law", viXra: 2208.0093[Religion and Spiritualism].
- [46] Anindya Kumar Biswas, "Dictionary of Culinary Terms by Philippe Pilibossian and the Graphical law", viXra: 2211.0061[Social Sciences].
- [47] Anindya Kumar Biswas, "A Greek and English Lexicon by H.G.Liddle et al simplified by Didier Fontaine and the Graphical law", viXra: 2211.0087[Linguistics].
- [48] Anindya Kumar Biswas, "Learner's Mongol-English Dictionary and the Graphical law", viXra: 2211.0101[Linguistics].
- [49] Anindya Kumar Biswas, "Complete Bulgarian-English Dictionary and the Graphical law", viXra: 2212.0009[Linguistics].
- [50] Anindya Kumar Biswas, "A Dictionary of Sindhi Literature by Dr. Motilal Jotwani and the Graphical Law", viXra: 2212.0015[Social Sciences].
- [51] Anindya Kumar Biswas, "Penguin Dictionary of Physics, the Fourth Edition, by John Cullerne, and the Graphical law", viXra: 2212.0072[History and Philosophy of Physics].
- [52] Anindya Kumar Biswas, "Oxford Dictionary of Chemistry, the seventh edition and the Graphical Law", viXra: 2212.0113[Chemistry].
- [53] Anindya Kumar Biswas, "A Burmese-English Dictionary, Part I-Part V, by J. A. Stewart and C. W. Dunn et al, head words and the Graphical Law", viXra: 2212.0127[Linguistics].
- [54] Anindya Kumar Biswas, "The Graphical Law behind the head words of Dictionary Kannada and English written by W. Reeve, revised, corrected and enlarged by Daniel Sanderson", viXra: 2212.0185[Linguistics].
- [55] Anindya Kumar Biswas, "Sanchayita and the Graphical Law", viXra: 2301.0075[Social Science].

- [56] Anindya Kumar Biswas, "Samsad Bangla Abhidan and The Graphical Law", viXra: 2302.0026[Linguistics].
- [57] Anindya Kumar Biswas, "Bangiya Sabdakosh and The Graphical Law", viXra: 2302.0060[Linguistics].
- [58] Anindya Kumar Biswas, "Samsad Bengali-English Dictionary and The Graphical Law ", viXra: 2304.0047[Linguistics].
- [59] Anindya Kumar Biswas, "Rudyard Kipling's Verse and the Graphical Law", viXra: 2304.0207[Social Science].
- [60] Anindya Kumar Biswas, "W. B. Yeats, The Poems and the Graphical Law", viXra: 2305.0008[Social Science].
- [61] Anindya Kumar Biswas, "The Penguin Encyclopedia of Places by W. G. Moore and the Graphical Law", viXra: 2305.0147[Archaeology].
- [62] Anindya Kumar Biswas, "The Poems of Tennyson and the Graphical Law", viXra: 2305.0157[Social Science].
- [63] Anindya Kumar Biswas, "Khasi-Jaintia Jaid(Surnames) and the Graphical law", viXra:2307.0135[Social Science].
- [64] Anindya Kumar Biswas, "Age, Amplitude of accommodation and the Graphical law", viXra:2311.0110[Physics of Biology].
- [65] Anindya Kumar Biswas, "Dictionary of Ayurveda by Dr. Ravindra Sharma and the Graphical law", viXra:2401.0030[General Science and Philosophy].
- [66] Anindya Kumar Biswas, "The Practical Sanskrit-English Dictionary by Vaman Shivram Apte and The Graphical Law", viXra:2402.0041[Linguistics].
- [67] Anindya Kumar Biswas, "The Langenscheidt's Pocket Russian Dictionary and The Graphical Law", viXra:2402.0049[Linguistics]
- [68] Anindya Kumar Biswas, "The Scholar Dictionary Portuguese and The Graphical Law", viXra:2402.0044[Linguistics]
- [69] Anindya Kumar Biswas, "The Langenscheidt's Pocket Japanese Dictionary and the Onsager's solution", viXra:2402.0052[Condensed Matter]
- [70] Anindya Kumar Biswas, "Langenscheidt's Pocket Chinese Dictionary and The Graphical Law", viXra:2403.0066[Linguistics]

- [71] Anindya Kumar Biswas, "Oxford Hindi-English Dictionary and The Graphical Law", viXra:2403.0129[Linguistics]
- [72] A. M. Gun, M. K. Gupta and B. Dasgupta, Fundamentals of Statistics Vol 1, Chapter 12, eighth edition, 2012, The World Press Private Limited, Kolkata.
- [73] E. Ising, Z.Physik 31,253(1925).
- [74] R. K. Pathria, Statistical Mechanics, p400-403, 1993 reprint, Pergamon Press,© 1972 R. K. Pathria.
- [75] C. Kittel, Introduction to Solid State Physics, p. 438, Fifth edition, thirteenth Wiley Eastern Reprint, May 1994, Wiley Eastern Limited, New Delhi, India.
- [76] W. L. Bragg and E. J. Williams, Proc. Roy. Soc. A, vol.145, p. 699(1934);
- [77] P. M. Chaikin and T. C. Lubensky, Principles of Condensed Matter Physics, p. 148, first edition, Cambridge University Press India Pvt. Ltd, New Delhi.
- [78] Kerson Huang, Statistical Mechanics, second edition, John Wiley and Sons(Asia) Pte Ltd.
- [79] Sonntag, Borgnakke and Van Wylen, Fundamentals of Thermodynamics, p206-207, fifth edition, John Wiley and Sons Inc.