Supplemented Newtonian Gravitational Equation

Istvan Polgar

Independent Researcher Email: polgar.istvan@proton.me

Abstract

Gravitational equation $F = G_N \frac{m_1 m_2}{r^2} + G_p \frac{m_1 m_2}{r}$, and new theory based on the effect of gravitational fields to each other. The calculated attractive force F eliminates the need for dark matter.

Introduction

According to Newton's law of universal gravitation[1], the magnitude of the attractive force is evenly distributed around a point-like mass in the form of a sphere. But according to this theory, this is only true if we do not count on the effect of gravitational fields on each other. If the attractive forces are represented by vectors, then the paths of vectors are curved by the gravitational fields in the direction of each other. And the effect of this increases with distance. In other words, the curvature of space-time has a different effect on the path of light than the path of gravity, because the path of light depends on time and always follows the fastest way[2].




Figure 1: The distance between straight and curved path on the gravitational circle increases the Newtonian gravitational force. The dot line of curved path follows Einstein's curvature of space-time $\propto \frac{1}{r^2}$.

$$\frac{\beta}{\alpha} = \frac{F_N + F_p}{F_N}$$
$$F = F_N + F_p$$
$$F_p = G_p \frac{m_1 m_2 r}{r^2} = G_p \frac{m_1 m_2}{r} = \left(\frac{8\pi}{c^4} r\right) F_N$$

- F_N [N] is the Newtonian force;
- F_p [N] is the force of path curvature effect;
- F [N] is the force between the masses;
- m_1 [kg] is the first mass;
- m_2 [kg] is the second mass;
- r [m] is the distance between the centers of masses;
- c = 299792.458 [km/s] is the speed of light;
- $G_N = 6.67430 \times 10^{-11} \left[N(\frac{m}{kg})^2 \right]$ is the Newtonian gravitational constant[3, 4];
- $G_p = \frac{8\pi G_N}{c^4} \approx 2.07665 \times 10^{-31} [N(\frac{m}{kg^2})]$ is the constant of gravitational path curvature effect. Value is calculated based on Einstein's gravitational constant[5] and the rotational velocity[6, 7] of an average disc-shaped galaxy;

The F_p and G_p value requires empirical and observable evidence. The current standard uncertainty of G_N is $U = 0.00015 \times 10^{-11} [N(\frac{m}{kg})^2][4]$, so the value of F_p cannot be measured within $\frac{U}{G_p} \approx 0.75[ly]$ by known methods. The calculated attractive force F eliminates the need for dark matter[8, 9].




Figure 2: Simplified galaxy rotation speed comparison: Newtonian vs Gravitational Path Curvature. Mass = 1.5×10^{41} kg.

Conclusion

Gravity is not the curvature of space-time because the curvature of spacetime affects gravity. Any theory based on spherically symmetric space-time and gravity needs to be revised. For example gravitational lensing, because the energy of light causes a space-time curvature, and the path of light depends on gravity, which is affected by this.

Acknowledgement

This research used *GeoGebra* and *GNU Octave* software and information from *WikipediA*.

References

- I. Bernard Cohen. The first english version of newton's hypotheses non fingo. *Isis*, 53(3):379–388, 1962.
- [2] Arto Annila. Least-time paths of light: Least-time paths of light. Monthly Notices of the Royal Astronomical Society, 416(4):2944–2948, July 2011.
- [3] Jens H. Gundlach and Stephen M. Merkowitz. Measurement of newton's constant using a torsion balance with angular acceleration feed-

back. *Phys.Rev.Lett.* 85 (2000) 2869-2872, 85(14):2869-2872, October 2000.

- [4] Peter Mohr, David Newell, Barry Taylor, and Eite Tiesinga. Codata recommended values of the fundamental physical constants: 2022. August 2024.
- [5] Jorge Pullin. General relativity. an introduction to the theory of the gravitational field (hans stephani). SIAM Rev., 33(4):674–675, 1991.
- [6] Daeun Jeong, Ho Seong Hwang, Haeun Chung, and Yongmin Yoon. Diverse rotation curves of galaxies in a simulated universe: The observed dependence on stellar mass and morphology reproduced. *The Astrophysical Journal*, 982(1):11, 2025.
- [7] Francesco Sylos Labini. Generalized rotation curves of the milky way from the gaia dr3 data set: Constraints on mass models. *The Astrophysical Journal*, 976(2):185, 2024.
- [8] Samurović S. The newtonian and mond dynamical models of ngc 5128: Investigation of the dark matter contribution. Serbian Astronomical Journal, 2016(192):9–20, 2016.
- [9] John Herbert Marr. Entropy and mass distribution in disc galaxies. Galaxies, 8(1):12, 2 2020.