

Gravitational Collapse as Analog of Continuous Spacetime Dimensions

Ervin Goldfain

Global Institute for Research, Education and Scholarship (GIRES), USA

E-mail ervinggoldfain@gmail.com

Abstract

Starting from the Newtonian potential of a two-body system, we recently pointed out that a spacetime endowed with continuous dimensions can be interpreted as analog of classical gravitation in four dimensions. Here we extend the analysis to relativistic two-body systems and suggest that the formation of Black Holes echoes the behavior of continuous spacetime dimensions in primordial cosmology.

Key words: continuous spacetime dimensions, relativistic two-body systems, gravitational collapse, Schwarzschild metric, Cantor Dust.

According to [1], far above the low-energy sector of field theory, the temporal component of the non-relativistic metric $g_{00}(\mu)$ is linearly related to the continuous deviation from four space-time dimensions $\varepsilon(\mu)$ as in

$$g_{00}(\mu) \approx 1 - 2 \frac{m^2(\mu)}{M_{pl}^2} \approx 1 - 2\varepsilon(\mu) \quad (1a)$$

in which

$$\varepsilon(\mu) = 4 - D(\mu) \quad (1b)$$

Here, μ is the observation scale, $m(\mu)$ stands for mass, M_{pl} is the Planck mass and the Newtonian potential is given by

$$\varphi_N(\mu) \approx \frac{1}{2} [g_{00}(\mu) - 1] \quad (2)$$

It is known that General Relativity (GR) does not allow a straightforward extrapolation of the two-body gravitational potential (2) to curved spacetime. In this instance, one appeals to several *effective models* such as the Post-Newtonian (PN) and Effective One Body (EOB) approximations. These are typically applied for treating two-body problems (for instance, binary black holes or neutron star binaries) in a full relativistic context [2 - 5].

With reference to EOB, the two-body problem with masses m_1 and m_2 is mapped onto:

- a) A test particle of reduced mass $m_R = m_1 m_2 / (m_1 + m_2)$,
- b) Total mass of the system $M = m_1 + m_2$,
- c) A symmetric mass ratio $\eta = m_R / M$.

The effective EOB line element is written in Schwarzschild-like coordinates as

$$ds_{\text{eff}}^2 = -A(r)dt^2 + B(r)dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2) \quad (3)$$

where the metric functions $A(r)$ and $B(r)$ contain all the relativistic corrections to the Newtonian metric. Specifically, the counterpart of (1a) is

$$A(r) = 1 - 2u + 2\eta u^3 + a_4(\eta)u^4 + \dots \quad (4)$$

with the Newtonian potential encoded in

$$-2u = -\frac{2G_N M}{r} = -\frac{2M}{M_{Pl}^2 r} \quad (5)$$

and nonlinear corrections included in the series $\sum_n a_n(\eta)u^n$, with ($n \geq 2$).

Likewise, the metric function $B(r)$ assumes the form

$$B(r) = \frac{1}{1-2u} + b_2(\eta)u^2 + b_3(\eta)u^3 + \dots \quad (6)$$

All relativistic terms can be safely ignored if $M = m_1 + m_2 \ll M_{Pl}$ or the radial coordinate approaches infinity (matching the flat manifold condition), i.e. $r \rightarrow \infty, u \rightarrow 0$.

With reference to [1], it is apparent that (5) recovers the expression of minimal dimensional deviation $\varepsilon \ll 1$ in the limit $m_1 = m_2 = m \ll M_{Pl}$ and by taking the radial coordinate to be comparable with the Compton wavelength, that is,

$$\boxed{u \rightarrow 0 \Leftrightarrow \varepsilon = O(m^2/M_{Pl}^2) \rightarrow 0} \quad (7)$$

It is also apparent that demanding the metric (3) to become singular at $A(r)=0, B(r) \rightarrow \infty$ leads to the well-known expression of the *Schwarzschild radius*, namely,

$$1 - 2u = 0 \Rightarrow r_G = 2G_N M = 4m/M_{Pl}^2 \quad (8)$$

Let the Schwarzschild radius (8) be again commensurate with the Compton wavelength, $r_G = O(\lambda_c) = O(m^{-1})$. By (1b), (8) yields

$$\boxed{\frac{4m^2}{M_{Pl}^2} \approx 4\varepsilon \approx 1} \quad (9)$$

Condition (9) signals a *regime of high fractality* $\varepsilon = O(1)$, a complex dynamic setting where there is a continuous *dimensional reduction* from the ordinary four-spacetime of classical and quantum physics [6].

Several key conclusions may be drawn from the combined use of (7), (9) and ref. [1], namely,

1. As emphasized in [1], (7) sets the stage for the gravitational interpretation of *Cantor Dust* (CD), under the assumption that CD acts like a large-scale cluster of ultralight Dark Matter (DM) particles such as self-interacting bosons, axions, fuzzy condensates or 3D anyons.
2. (9) hints that gravitational collapse and the formation of Black Holes is analogous to the process of *dimensional condensation* and the emergence of Cantor Dust.
3. (9) also hints that dimensional condensation of DM is analogous to DM formation via *primordial Black Holes*.

References

1. Goldfain, E., (2025), On the Gravitational Analog of Continuous Spacetime Dimensions, preprint <http://dx.doi.org/10.13140/RG.2.2.24427.25127/2>
2. Buonanno, A. & Damour, T. (1999), *Phys. Rev. D* **59**, 084006
3. Damour, T. (2010), *Int. J. Mod. Phys. A* **23**, 1130

4. Damour, T., Jaranowski, P., Schäfer, G. (2000), *Phys. Rev. D* **62**, 084011
5. Nagar, A. & Akcay, S. et al. (2021), EOB waveform models for GW detectors
6. Goldfain, E., (2023, Dimensional Reduction as Source of Cosmological Anomalies, preprint <https://doi.org/10.32388/RX6BTE>