

Modeling the nS1/2 Lamb shifts in atomic hydrogen as magnetic monopole phenomena.

Hans Peter Good
Sargans, Switzerland
e-mail: hp.good at catv.rol.ch

July 8, 2025

Abstract

Relativistic quantum mechanics fails to provide an explanation for the Lamb shift. It is proposed that a point-like magnetic monopole associated with the proton interacts with the magnetic dipole moment of the electron, causing Zeeman energy shifts of the nS1/2 levels. If the 1S1/2–2S1/2 transition in atomic hydrogen is taken as a reference, the model predicts 2 922 743 278.669 71 MHz for the 1S1/2–3S1/2 transition, and 770 649 350.463 33 MHz for the 2S1/2–8S1/2 interval without the application of the constants m_e , h , or c defined by CODATA.

Keywords: Relativistic quantum mechanics, Lamb shift, magnetic monopole, atomic hydrogen, proton structure.

In [1], the author presented a self-consistent mathematical formalism to characterize the atomic hydrogen spectrum, revealing that, considering the margins of error, the spectroscopic fine structure constant α aligns with $\alpha_{geom} \equiv 2^{-6}\pi^{\frac{2}{3}} \approx 0.00\ 72\ 843$ ($\alpha_{geom}^{-1} \approx 137.28$). Throughout the text, including the tables and formulas that follow, the fine structure constant α is an exact parameter, and the symbol α always denotes the number constant α_{geom} .

The most recent data from Table 1 was used to supply the algorithm outlined in [1]. Only the measured values and uncertainties listed in Table 1 were used to estimate the parameters in Table 2. Other data, such as the electron mass m_e , the Planck constant h , or the speed of light in vacuum c , have no impact. They are embedded within the scaling factor γ_{zero} , which also adjusts for the recoil correction coming from nuclear motion.

Relativistic quantum mechanics fails to provide an explanation for the Lamb shift. It is interpreted through quantum field theory, which constitutes a fundamentally different paradigm. Assuming that a point-like magnetic monopole μ of the proton interacts with the magnetic dipole moment of the electron μ_e , causing Zeeman energy shifts of the nS1/2 levels due to the proton magnetic monopole field. If the magnetic field is attributed to a magnetic charge and originates from a scalar potential, meaning the source of magnetism is a type of spherical bar magnet, then the factor $(-1/2)$ must be taken into account [4]. Electrons that are confined circulate due to the monopole field and create an induced magnetic field that opposes the monopole field, as per Lenz's law, indicating the presence of local diamagnetic shielding characterized by the factor $\gamma_{diamag_shielding}$. With all these aspects in mind, it is theorized that the dimensionless Lamb shift constant $B(\gamma_{zero})$ equals

$$B(\gamma_{zero})^{theo} = \frac{\mu}{\mu_B} \cdot \left(-\frac{1}{2}\right) \cdot \gamma_{diamag_shielding} \cdot (2\alpha)^2 \quad (1)$$

where μ signifies an intrinsic magnetic moment of the proton, which needs to be defined so that the difference from the precisely estimated value $B(\gamma_{zero})$ in Table 2 is as small as possible. Simple numerical testing [2:p.175] gives

$$\frac{\mu}{\mu_B} = \frac{5}{\left(\frac{\lambda_{e_bar}}{L_2(L)} + 3\right)} \approx 0.023025 \quad (2)$$

$$\frac{\lambda_{e_bar}}{L_2(L)} = 2^{16} \pi^{-5}$$

and

$$\gamma_{diamag_shielding} \equiv \frac{1}{3} \left(1 + 2\sqrt{1 - (3\alpha)^2}\right) \approx 1 - \frac{(3\alpha)^2}{3} \quad (3)$$

being consistent with the relativistic adjustment to the magnetic moment for a central electric potential of charge $Z = 3$ and a uniform magnetic field of arbitrary strength [5: p. 213, eq. 47.2]. The energy building block $\frac{1}{L_2(L)}$ in the energy unit $\frac{1}{\lambda_{e_bar}}$, where λ_{e_bar} corresponds to the reduced Compton wavelength of the electron, is also connected to the saturation density of matter established through elastic electron scattering experiments [6]. Furthermore, the phenomenological expression (1) is consistent with the concept that symmetries in quantum theory result in simple mathematical equations that include integers or fractions of integers.

According to the compact algebraic formula (1), the calculated value $B(\gamma_{zero})^{theo}$ is about **-2.443 049 ppm**, which is comfortably within the error limits of the experimental result **-2.443 049(17) ppm** presented in Table 2. When the input A7 of Table 1 is used as a “metrological” reference to determine the scaling factor γ_{zero} , the theoretical dimensionless value $B(\gamma_{zero})^{theo}$ predicts, according to formula (1), **2 922 743 278.669 71 MHz** for the 1S1/2–3S1/2 interval, and **770 649 350.463 33 MHz** for the 2S1/2–8S1/2 transition. For consistency verification, it is essential to experimentally test and confirm these conjectures with accurate optical frequency measurements.

Table 1: Theoretical (exact) and experimental (stochastic) input data.

	value	error	unit	reference	
α	$2^{-6} \pi^{-\frac{2}{3}}$	exact		[2:p.42]	
1S1/2–2S1/2	2 466 061 413.187 03	0.000 01	MHz	Table X, A6	[3]
1S1/2–2S1/2	2 466 061 413.187 01	0.000 01	⋮	Table X, A7	[3]
1S1/2–3S1/2	2 922 743 278.659	0.017	⋮	Table X, A8	[3]
1S1/2–3S1/2	2 922 743 278.678	0.013	⋮	Table X, A22	[3]
1S1/2–3S1/2	2 922 743 278.671 5	0.002 6	MHz	Table X, A23	[3]

Table 2: Output values obtained from the input data (Table 1) by means of formula (1) provided in [1]. The least squares solutions $A(\gamma = 1)$ and $B(\gamma = 1)$ were calculated by setting $m_e = h = c = 1$. The parameter $B(\gamma_{zero})$ is given by $B(\gamma_{zero}) = B(\gamma = 1) / \gamma_{zero}$ with $\gamma_{zero} = 1 + A(\gamma = 1)$. The statistical analysis involved 1000 data sets (Table 1) whose five experimental transitions were randomly varied within the error margins.

parameter	mean value	error (1 σ , random)	unit	rel. std. uncert.
$B(\gamma_{zero})$	-2.443 049	0.000 017	ppm	7.1×10^{-6}
Lamb shift $\mathcal{L}(1S1/2)$	8 032.976	0.057	MHz	7.1×10^{-6}
Ionization	3 288 086 857.911	0.009	MHz	2.9×10^{-12}
mean abs. dev.	0.004 8	0.002 4	MHz	
γ_{zero}	$1.239 353 \times 10^{20}$			2.0×10^{-11}

References

- [1] The Spectra of Hydrogen and Deuterium Interpreted with an Alternative Fine Structure Constant Compared to the Codata Recommended Value.
Hans Peter Good viXra 2411.0170 (2025)
- [2] On the Origin of Natural Constants.
Hans Peter Good, De Gruyter Berlin (2018)
- [3] CODATA recommended values of the fundamental physical constants (2018).
E Tiesinga et al Rev Mod Phys **93** 025010-1 (2021)
- [4] The nature of intrinsic magnetic dipole moments: what can the famous 21 cm astrophysical spectral line of atomic hydrogen tell us about the nature of magnetic dipoles?
J D Jackson, CERN 77-17 Lecture (1977)
- [5] Quantum Mechanics of One- and Two-Electron Atoms.
H A Bethe, E E Salpeter, Springer-Verlag Berlin (1957)
- [6] Calculation of the Nuclear Saturation Density.
Hans Peter Good viXra 2112.0111 (2024)